

ON THE CHOQUET AND BISHOP-DE LEEUW THEOREMS

SILVIU TELEMAN

*National Institute for Scientific and Technical Creation,
Bucharest, România*

In the theory of integral representation of points in compact convex sets (Choquet theory), there are two fundamental theorems, which assert that the maximal Radon probability measures on such (sub) sets (of Hausdorff locally convex topological real vector spaces) are pseudoconcentrated on their sets of extreme points. Maximality is meant here either in the sense of the Bishop-de Leeuw preorder relation, to which the Bishop-de Leeuw theorem belongs (see [3], Theorem 5.3), or in the sense of the Choquet order relation, to which the Choquet theorem belongs (see [5]; [7], Theorem T32; [8], Ch. 4). An immediate consequence of these two theorems is the Choquet-Bishop-de Leeuw theorem, which yields the representability of any point in the given compact convex set by a boundary integral (see [8], Ch. 4).

Chronologically, Choquet first proved a theorem of this kind, for metrizable compact convex sets, in 1956 (see [4]). In 1959 Bishop and de Leeuw, by using a certain preorder relation (which will be recalled in what follows) in the set of all Radon probability measures, on arbitrary compact convex subsets of Hausdorff locally convex topological real vector spaces, proved that any measure which is maximal for the preorder relation in question is pseudoconcentrated on the set of all extreme points of the given compact convex set.

By establishing the existence of such maximal measures, they thus obtained an extension of Choquet's theorem (see [3], Theorem 5.6).

In 1960, Choquet, by using another order relation, showed that any maximal measure (with respect to his order relation) is pseudoconcentrated on the set of all extreme points of the given compact convex set. Again, by proving the existence of such maximal measures, Choquet obtained another proof of Bishop's and de Leeuw's extension of Choquet's theorem, to the effect that any point of the given compact convex set K is represented by a Radon probability measure, which is pseudoconcentrated on the set of all extreme points of K . This result is known as the Choquet-Bishop-de Leeuw theorem (see [8], Ch. 4).

Of course, both the Bishop-de Leeuw theorem and the Choquet theorem are stronger than the Choquet-Bishop-de Leeuw theorem, in that they assert that any maximal measure, either with respect to the Bishop-de Leeuw preorder relation,

or with respect to the Choquet order relation, is pseudoconcentrated on the set of extreme points.

Since the two (pre) order relations are, in general, different (see [9], p. 289), these two theorems are different.

The Bishop-de Leeuw preorder relation is very useful in connection with the central and irreducible disintegration of the representations of C^* -algebras (see [9], [10], [11], [12]). However, for the orthogonal Radon probability measures, defined on the state space of a C^* -algebra, the two order relations coincide (see [9], Theorem 13).

The aim of this note is to give a generalization of the theorems of Choquet and Bishop-de Leeuw. In this manner we shall obtain a new and unitary proof for both the theorems of Bishop-de Leeuw and Choquet. This will be achieved by means of a slight extension of H. Bauer's Minimum Principle. The exposition is almost self-contained.

1

Let E be a Hausdorff locally convex topological real vector space and $K \subset E$ a compact convex subset of E .

By $C(K; \mathbb{R})$ we shall denote the algebra of all continuous real functions defined on K ; by $A(K; \mathbb{R})$ we shall denote the real vector space of all continuous affine real functions defined on K ; by $S(K; \mathbb{R})$ we denote the convex sup-cone of all convex continuous real functions which are defined on K . It is well known that $S(K; \mathbb{R}) - S(K; \mathbb{R})$ is a vector sublattice of $C(K; \mathbb{R})$, uniformly dense in virtue of the Stone approximation theorem (see [8], Ch. 4). It is obvious that $h \in A(K; \mathbb{R}) \Rightarrow h^2 \in S(K; \mathbb{R})$.

Let $\mathcal{M}_+^1(K)$ be the convex set of all Radon probability measures on K ; for any $\mu \in \mathcal{M}_+^1(K)$ there exists a uniquely determined $b(\mu) \in K$ such that

$$h(b(\mu)) = \int_K h(x) d\mu(x), \quad \forall h \in A(K; \mathbb{R});$$

$b(\mu)$ is called the *barycenter* of μ (see [8], Ch. 1).

Two measures $\mu, \nu \in \mathcal{M}_+^1(K)$ are said to be *equivalent* if $b(\mu) = b(\nu)$. One denotes this relation by $\mu \sim \nu$. A point $x \in K$ is said to be *represented* by $\mu \in \mathcal{M}_+^1(K)$ if $x = b(\mu)$. It is obvious that any $x \in K$ is represented by the Dirac measure ε_x at x .

The fundamental theorems of Choquet and Bishop-de Leeuw give existential solutions to the following problem: given $x \in K$, represent it by a measure $\mu \in \mathcal{M}_+^1(K)$ whose support is as close to $\text{ex}K$ as possible (here $\text{ex}K$ denotes the set of all extreme points of K).

We consider the following relations on $\mathcal{M}_+^1(K)$:

- (a) $\mu < \nu$ iff $\mu(f) \leq \nu(f)$, for any $f \in S(K; \mathbb{R})$;
- (b) $\mu \ll \nu$ iff $\mu(h^2) \leq \nu(h^2)$, for any $h \in A(K; \mathbb{R})$.

The first is due to Choquet (see [5], [7], [8]), and the second to Bishop and de Leeuw (see [3]).

Obviously, $\mu < \nu$ implies $\mu \ll \nu$ for any $\mu, \nu \in \mathcal{M}_+^1(K)$. Both relations are reflexive and transitive; the first is also antisymmetric, hence an order relation; this follows from the fact that " $\mu < \nu$ and $\nu < \mu$ " implies $\mu(f) = \nu(f)$ for any $f \in S(K; \mathbb{R})$. Consequently, the same equality holds for any $f \in S(K; \mathbb{R}) - S(K; \mathbb{R})$. Stone's approximation theorem now implies that $\mu = \nu$.

Since $A(K; \mathbb{R}) = -A(K; \mathbb{R}) \subset S(K; \mathbb{R})$, it is easy to see that $\mu < \nu \Rightarrow \mu \sim \nu$. Since $\pm 1 \in A(K; \mathbb{R})$, an elementary argument shows that $\mu \ll \nu \Rightarrow \mu \sim \nu$.

An element $\mu \in \mathcal{M}_+^1(K)$ is maximal for the Choquet order relation if $\nu \in \mathcal{M}_+^1(K)$ and $\mu < \nu$ implies $\mu = \nu$. Similarly, an element $\mu \in \mathcal{M}_+^1(K)$ is maximal for the Bishop-de Leeuw preorder relation if $\nu \in \mathcal{M}_+^1(K)$ and $\mu \ll \nu$ implies $\mu(h^2) = \nu(h^2)$, $\forall h \in A(K; \mathbb{R})$.

In order to get the generalization of the theorems of Choquet and Bishop-de Leeuw which we have in view, we shall consider an arbitrary subset $S \subset S(K; \mathbb{R})$ such that

$$\{h^2; h \in A(K; \mathbb{R})\} \subset S.$$

We shall define a preorder relation in $\mathcal{M}_+^1(K)$ by defining

$$\mu \triangleleft \nu \Leftrightarrow \mu(f) \leq \nu(f), \quad \forall f \in S, \quad \forall \mu, \nu \in \mathcal{M}_+^1(K).$$

Of course, the relation is reflexive and transitive, and $\mu < \nu \Rightarrow \mu \triangleleft \nu$, $\mu \triangleleft \nu \Rightarrow \mu \ll \nu$, $\forall \mu, \nu \in \mathcal{M}_+^1(K)$. It follows that $\mu \triangleleft \nu \Rightarrow \mu \sim \nu$, $\forall \mu, \nu \in \mathcal{M}_+^1(K)$. A measure $\mu \in \mathcal{M}_+^1(K)$ is *maximal* for the preorder relation \triangleleft iff $\nu \in \mathcal{M}_+^1(K)$ and $\mu \triangleleft \nu \Rightarrow \mu(f) = \nu(f)$, $\forall f \in S$. In this case we shall say that μ is (\triangleleft)-*maximal*.

2

Let $\mathcal{B}(K)$ be the σ -algebra of all Borel measurable subsets of K and $\mathcal{B}_0(K)$ the σ -algebra of all Baire measurable subsets of K . We recall that $\mathcal{B}(K)$ is the σ -algebra generated by all open (or, equivalently, closed) subsets of K , whereas $\mathcal{B}_0(K)$ is the smallest σ -algebra of subsets of K such that all functions $f \in C(K; \mathbb{R})$ are measurable. Obviously, one has the inclusion $\mathcal{B}_0(K) \subset \mathcal{B}(K)$. If K is metrizable, then the equality holds.

A measure $\mu \in \mathcal{M}_+^1(K)$ is said to be *pseudoconcentrated* on $\text{ex}K$ if

- (*) $\mu(U) = 0$ for any $U \in \mathcal{B}_0(K)$ such that $U \cap (\text{ex}K) = \emptyset$.

Let $\mathcal{A}_0(\text{ex}K)$ be the σ -algebra of all subsets of $\text{ex}K$ which are traces on $\text{ex}K$ of the Baire measurable subsets of K

$$\mathcal{A}_0(\text{ex}K) = \{U \cap (\text{ex}K); U \in \mathcal{B}_0(K)\}.$$

If the measure $\mu \in \mathcal{M}_+^1(K)$ is pseudoconcentrated on $\text{ex}K$, then one can define a probability measure $\check{\mu}$ on $\mathcal{A}_0(\text{ex}K)$ by the formula

$$\check{\mu}(U \cap (\text{ex}K)) = \mu(U), \quad \forall U \in \mathcal{B}_0(K).$$

Property (*) ensures the correctness of the definition.

It is easy to prove now that if $f: K \rightarrow \mathbb{R}$ is a Baire measurable function (i.e., if it is $\mathcal{B}_0(K)$ -measurable), then $f|_{\text{ex}K}$ is $\mathcal{B}_0(\text{ex}K)$ -measurable. The importance of measures which are pseudoconcentrated on $\text{ex}K$ is shown by the following:

LEMMA 1. Let $\mu \in \mathcal{M}_+^1(K)$ be pseudoconcentrated on $\text{ex}K$ and $f: K \rightarrow \mathbb{R}$ a bounded Baire measurable function. Then

$$\int_K f d\mu = \int_{\text{ex}K} f d\check{\mu}.$$

Proof. For any $\varepsilon > 0$ one can find a partition $\{U_1, U_2, \dots, U_n\}$ of K consisting of Baire measurable subsets of K , and real numbers $a_i \in \mathbb{R}$, $i = 1, 2, \dots, n$, such that

$$\left| f(x) - \sum_{i=1}^n a_i \chi_{U_i}(x) \right| < \varepsilon, \quad \forall x \in K.$$

We then have

$$\left| \int_K f d\mu - \sum_{i=1}^n a_i \mu(U_i) \right| < \varepsilon$$

and

$$\left| \int_{\text{ex}K} f d\check{\mu} - \sum_{i=1}^n a_i \check{\mu}(U_i \cap (\text{ex}K)) \right| < \varepsilon.$$

By taking into account the definition of $\check{\mu}$, we now easily get

$$\left| \int_K f d\mu - \int_{\text{ex}K} f d\check{\mu} \right| < 2\varepsilon,$$

for any $\varepsilon > 0$. The lemma is proved.

A simple compactness argument shows that the (\blacktriangleleft) -ordered set $\mathcal{M}_+^1(K)$ is inductive. Hence Zorn's lemma implies that for any $\mu \in \mathcal{M}_+^1(K)$ there exists a (\blacktriangleleft) -maximal measure $\nu \in \mathcal{M}_+^1(K)$ such that $\mu \blacktriangleleft \nu$. In particular, this is true for the order relation $<$ (see [8], Ch. 4), as well as for the preorder relation \ll (see [3], p. 307).

3

The following lemma is a slight extension of Proposition 4.2 from [8] (which is identical with part (b) of the lemma). The proof is adapted from that of Proposition 4.2 from [8].

We first recall that for any $f \in C(K; \mathbb{R})$ one defines its upper semicontinuous concave hull \bar{f} by

$$\bar{f}(x) = \inf_{\substack{h \in A(K; \mathbb{R}) \\ h \geq f}} h(x).$$

It is easy to prove that (see [8], Ch. 3):

- (a) \bar{f} is concave, bounded and upper semicontinuous for any $f \in C(K; \mathbb{R})$;
- (b) $f \leq \bar{f}$ and $\bar{f} = f \Leftrightarrow f$ is concave, $\forall f \in C(K; \mathbb{R})$;
- (c) if $f, g \in C(K; \mathbb{R})$, then $(f+g)^- \leq \bar{f} + \bar{g}$ and $|\bar{f} - \bar{g}| \leq \|f - g\|$; $(f+g)^- = \bar{f} + \bar{g}$, for $g \in A(K; \mathbb{R})$; $(rf)^- = r\bar{f}$, for any $r \in \mathbb{R}_+$ and $f \in C(K; \mathbb{R})$;
- (d) if $f \leq f'$, where f' is concave and upper semicontinuous, then $\bar{f} \leq f'$.

LEMMA 2. (a) If $\mu \in \mathcal{M}_+^1(K)$ is any (\blacktriangleleft) -maximal measure, then $\mu(\bar{f}) = \mu(f)$ for any $f \in S$; in particular, $\mu((h^2)^-) = \mu(h^2)$, $\forall h \in A(K; \mathbb{R})$.

(b) If $\mu \in \mathcal{M}_+^1(K)$ is any $(<)$ -maximal measure, then $\mu(\bar{f}) = \mu(f)$, for any $f \in C(K; \mathbb{R})$.

Proof. Let $f_0 \in C(K; \mathbb{R})$. We define a real linear functional $L: \mathbb{R}f_0 \rightarrow \mathbb{R}$ by the formula

$$L(af_0) = a\mu(\bar{f}_0), \quad a \in \mathbb{R}.$$

We also define a positively homogeneous sublinear functional $p: C(K; \mathbb{R}) \rightarrow \mathbb{R}$ by the formula

$$p(f) = \mu(\bar{f}), \quad f \in C(K; \mathbb{R}).$$

For $a \in \mathbb{R}$, $a \geq 0$, we obviously have $L(af_0) = p(af_0)$. For $a < 0$, from

$$0 = (af_0 - af_0)^- \leq (af_0)^- + (-af_0)^- = (af_0)^- - a\bar{f}_0,$$

we infer that

$$L(af_0) = a\mu(\bar{f}_0) = \mu(a\bar{f}_0) \leq \mu((af_0)^-) = p(af_0).$$

The Hahn-Banach theorem now implies that there exists a linear functional $L': C(K; \mathbb{R}) \rightarrow \mathbb{R}$ such that

$$L'|_{\mathbb{R}f_0} = L \quad \text{and} \quad L'(f) \leq p(f), \quad \forall f \in C(K; \mathbb{R}).$$

If $g \in C(K; \mathbb{R})$, $g \leq 0$, we have $g \leq 0$ and, therefore,

$$L'(g) \leq p(g) = \mu(\bar{g}) \leq 0.$$

This shows that $L' \geq 0$ as a linear functional on $C(K; \mathbb{R})$ and, therefore, there exists a positive Radon measure ν on K such that

$$L'(f) = \nu(f), \quad \forall f \in C(K; \mathbb{R}).$$

If $f \in C(K; \mathbb{R})$ is convex, then $-f$ is concave and, therefore, $(-f)^- = -f$; it follows that

$$\mu(f) = -\mu(-f) = -\mu((-f)^-) = -p(-f) \leq -L'(-f) = L'(f) = \nu(f),$$

and, consequently, we have $\mu \prec \nu$; hence we also have $\mu \blacktriangleleft \nu$, i.e., $\mu(f) \leq \nu(f)$, $\forall f \in S$.

(a) If μ is (\blacktriangleleft) -maximal, we infer that $\mu(f) = \nu(f)$, $\forall f \in S$. In particular, we have (if $f_0 \in S$):

$$\mu(f_0) = \nu(f_0) = L'(f_0) = L(f_0) = \mu(\bar{f}_0).$$

(b) If μ is $(<)$ -maximal, then $\mu = \nu$. It follows that

$$\mu(f_0) = \nu(f_0) = L'(f_0) = L(f_0) = \mu(\bar{f}_0).$$

The lemma is proved.

For any $f \in C(K; \mathbb{R})$ let us denote $S_f = \{x \in K; \bar{f}(x) = f(x)\}$. Obviously, S_f is a G_δ -subset of K .

PROPOSITION 1 (H. Bauer). For any $x \in K$ we have

$$x \in \text{ex}K \Leftrightarrow \text{there is a unique } \mu \in \mathcal{M}_+^1(K) \text{ such that } b(\mu) = x.$$

Proof (see [8], Ch. 1). If $x \in \text{ex}K$ and $\mu \in \mathcal{M}_+^1(K)$ is such that $b(\mu) = x$, we have $\text{supp } \mu = \{x\}$. Indeed, it is sufficient to prove that $\mu(D) = 0$ for any compact set $D \subset K \setminus \{x\}$. If this is not true, then there exists a compact set $D \subset K \setminus \{x\}$ and a point $y \in D$ such that $\mu(D \cap U) > 0$ for any neighbourhood U of y . Let us choose a compact convex set U , such that $x \notin U$ and define $K_0 = \overline{\text{co}}(D \cap U)$. Then $x \notin K_0$ and $\mu(K_0) > 0$. Let us define $\mu_1 = (\mu(K_0))^{-1}(\chi_{K_0}\mu)$. We obviously have $b(\mu_1) \in K_0$. If $\mu(K_0) = 1$, then $\mu_1 = \mu$, hence $x = b(\mu) = b(\mu_1) \in K_0$, a contradiction. Consequently, we have $\mu(K_0) < 1$. Let $\mu_2 = (1 - \mu(K_0))^{-1}(\chi_{K \setminus K_0}\mu)$. Then $b(\mu_2) \in K$ and $\mu = \mu(K_1)\mu_1 + (1 - \mu(K_1))\mu_2$; this implies that

$$x = b(\mu) = \mu(K_1)b(\mu_1) + (1 - \mu(K_1))b(\mu_2),$$

where $b(\mu_1) \neq x$, a contradiction. Consequently, $\mu = \varepsilon_x$.

Conversely, if $x \in K \setminus \text{ex}K$, then there exist $x_1, x_2 \in K$ such that $x = \frac{1}{2}(x_1 + x_2)$ and $x_1, x_2 \neq x$. We then have

$$\varepsilon_x \sim \frac{1}{2}(\varepsilon_{x_1} + \varepsilon_{x_2}) \quad \text{and} \quad \varepsilon_x \neq \frac{1}{2}(\varepsilon_{x_1} + \varepsilon_{x_2}).$$

The proposition is proved.

PROPOSITION 2 (see [8], Proposition 3.1). For any $f \in C(K; \mathbb{R})$ and any $x \in K$ we have

$$\bar{f}(x) = \sup_K \left\{ \int_K f(x) d\mu(x); \mu \sim \varepsilon_x \right\}.$$

In particular, if $x \in \text{ex}K$, then $\bar{f}(x) = f(x)$.

Proof. Let us define $f'(x) = \sup_K \left\{ \int_K f(x) d\mu(x); \mu \sim \varepsilon_x \right\}$. Then we obviously have $f(x) \leq f'(x)$, $x \in K$, and f' is easily shown to be concave. In order to prove that it is upper semicontinuous, let $r \in \mathbb{R}$ and let $(x_\alpha)_{\alpha \in I}$ be a convergent net in K , $x = \lim_{\alpha \in I} x_\alpha$ such that $f'(x_\alpha) \geq r$, $\forall \alpha \in I$. For any $\varepsilon > 0$ and any $\alpha \in I$ we can choose a $\mu_\alpha \in \mathcal{M}_+^1(K)$ such that $\mu_\alpha \sim \varepsilon_{x_\alpha}$ and $r - \varepsilon \leq f'(x_\alpha) - \varepsilon < \mu_\alpha(f)$, $\forall \alpha \in I$. Since the set $\mathcal{M}_+^1(K)$ is compact for the vague topology, we can choose a convergent subnet $(\mu_{\alpha(\beta)})_{\beta \in J}$, $\lim_{\beta \in J} \mu_{\alpha(\beta)} = \mu$. We obviously have $b(\mu) = x$ and $r - \varepsilon \leq \mu(f) \leq f'(x)$, for any $\varepsilon > 0$; hence, $r \leq f'(x)$, and this shows that f' is upper semicontinuous. Consequently, the inequality $f \leq f'$ implies that $\bar{f} \leq f' = f$. On the other hand, we have (for any $\mu \in \mathcal{M}_+^1(K)$, $\mu \sim \varepsilon_x$),

$$\int_K f(x) d\mu(x) \leq \int_K \bar{f}(x) d\mu(x) \leq \bar{f}(x),$$

and this implies that $f'(x) \leq \bar{f}(x)$, $x \in K$.

COROLLARY 1. (a) For any $f \in C(K; \mathbb{R})$ we have $S_f \supset \text{ex}K$.

(b) If $H \subset A(K; \mathbb{R})$ is a total set, then

$$\bigcap_{h \in H} S_{h^2} = \text{ex}K.$$

(c) If K is metrizable, then there exists a sequence $(h_n)_{n \geq 0}$, $h_n \in A(K; \mathbb{R})$, $\|h_n\| \leq 1$, $\forall n \geq 0$, such that

$$\bigcap_{n=0}^{\infty} S_{h_n^2} = \text{ex}K.$$

Consequently, if K is metrizable, then $\text{ex}K$ is a G_δ -subset of K .

Proof. (a) is an immediate consequence of Proposition 2.

(b) Let us remark that if H is a total subset of $A(K; \mathbb{R})$, then H separates the points of K . Now let $x \in \bigcap_{h \in H} S_{h^2}$. If $x \notin \text{ex}K$, then there exist $x_1, x_2 \in K$, $x_1, x_2 \neq x$, such that $x = \frac{1}{2}(x_1 + x_2)$. Since the set H separates the points of K , there exists a $h_0 \in H$ such that $h_0(x_1) \neq h_0(x)$. Then, for any $h \in A(K; \mathbb{R})$, we have

$$h \geq h_0^2 \Rightarrow h(x) \geq |h_0(x_1) - h_0(x)|^2 + h_0^2(x).$$

Consequently, we have $\bar{h}_0^2(x) \geq |h_0(x_1) - h_0(x)|^2 + h_0^2(x) > h_0^2(x)$, and this implies that $x \notin S_{h_0^2}$.

(c) If K is metrizable, then $C(K; \mathbb{R})$ is separable. Consequently, $A(K; \mathbb{R})$ is separable and, therefore, in $\{h \in A(K; \mathbb{R}); \|h\| \leq 1\}$ there exists a countable, dense subset H , which, obviously, is total in $A(K; \mathbb{R})$. Since any S_f is a G_δ -subset in K , in this case $\text{ex}K$ is also a G_δ -subset.

COROLLARY 2. (a) For any (\blacktriangleleft) -maximal measure $\mu \in \mathcal{M}_+^1(K)$, and any $f \in S$ we have $\mu(S_f) = 1$; in particular, $\mu(S_{h^2}) = 1$, $\forall h \in A(K; \mathbb{R})$.

(b) If K is metrizable, and $\mu \in \mathcal{M}_+^1(K)$ is any (\blacktriangleleft) -maximal measure, then $\mu(\text{ex}K) = 1$.

Proof. Assertion (a) is an immediate consequence of Lemma 2 and of the definition of the set S_f , $f \in C(K; \mathbb{R})$. Assertion (b) is an immediate consequence of assertion (a) and of Corollary 1.

4

An important result in Convex Analysis is the following "Minimum Principle" of H. Bauer (see [2], Lemma 1).

THEOREM 1 (H. Bauer). Let $f_1, f_2: K \rightarrow \mathbb{R}$ be semicontinuous functions, with f_1 convex and f_2 concave. Then

$$f_1(x) \leq f_2(x), \forall x \in \text{ex}K \Rightarrow f_1(x) \leq f_2(x), \forall x \in K.$$

(Here any kind of semicontinuity is allowed for the two functions; moreover, one does not assume that they have the same kind of semicontinuity, i.e., any combination is allowed.)

We refer to [2] for the proof of this theorem. We shall need a slight, but partial, extension of this theorem, which will play the main role in our generalization of Choquet's and Bishop's de Leeuw's theorems.

THEOREM 2. *Let $f: K \rightarrow \mathbb{R}$ be a concave semicontinuous function. Then*

$$f(x) > 0, \forall x \in \text{ex}K \Rightarrow f(x) > 0, \forall x \in K.$$

Proof. (a) Let us assume that f is lower semicontinuous. Then $m = \inf f(x)$ is attained on K , i.e., there exists an $x_0 \in K$ such that $f(x_0) = m$. Let $K_0 = \{x \in K; f(x) = m\}$. Since f is lower semicontinuous, K_0 is a (non-empty) compact subset of K . Let $K_1 = \overline{\text{co}}(K_0)$. Milman's theorem implies that $\text{ex}K_1 \subset K_0$. Let $x_1 \in \text{ex}K_1$; then $f(x_1) = m$. If $x', x'' \in K$ are such that $x', x'' \neq x_1$, $x_1 = \frac{1}{2}(x' + x'')$, then, since f is concave, we have

$$m = f(x_1) \geq \frac{1}{2}(f(x') + f(x'')) \geq m,$$

and, therefore, $f(x') = f(x'') = m$. Hence $x', x'' \in K_0 \subset K_1$. Consequently, since $x_1 \in \text{ex}K_1$, we have $x' = x'' = x_1$. This shows that $x_1 \in \text{ex}K$, and $m = f(x_1) > 0$; therefore, we have $f(x) \geq m > 0$, for any $x \in K$.

(b') Let us now assume that f is upper semicontinuous, and let $x_0 \in K$. Then we have $f = \bar{f}$ and, therefore, we have

$$f(x_0) = \inf_{h \in A(K; \mathbb{R})} h(x_0).$$

For any $n \in \mathbb{N}^*$ we can choose a $h_n \in A(K; \mathbb{R})$ such that $h_n \geq f$ on K and

$$f(x_0) + 1/n > h_n(x_0), \quad n \geq 1.$$

Let $f_0 = \inf_{n \geq 1} h_n$. Then $f_0: K \rightarrow \mathbb{R}$ is a Baire measurable, concave, upper semicontinuous function such that

$$f_0(x) \geq f(x) > 0 \quad \text{for any } x \in \text{ex}K$$

and

$$f_0(x_0) = f(x_0).$$

(b'') For any $r \in \mathbb{R}$ let us define

$$K(r) = \{x \in K; f_0(x) < r\}.$$

For any $r \in \mathbb{R}$, the set $K(r)$ is Baire measurable and, therefore, there exists a countable subset $H_r \subset A(K; \mathbb{R})$ such that $K(r)$ belongs to the smallest σ -algebra Σ_r of subsets of K such that all functions in H_r are Σ_r -measurable.

Let $H = \bigcup_{r \in \mathbb{Q}} H_r$. Then H is a countable subset of $A(K; \mathbb{R})$. Let Σ be the smallest σ -algebra of subsets of K such that all functions in H are Σ -measurable. Then we have $\Sigma_r \subset \Sigma$, $\forall r \in \mathbb{Q}$, and, therefore, $K(r) \in \Sigma$, $\forall r \in \mathbb{Q}$.

For any $r \in \mathbb{R}$ and any sequence $(r_n)_{n \geq 0}$ such that $r_n \in \mathbb{Q}$, $r_n \uparrow r$, we have

$$K(r) = \bigcup_{n \geq 0} K(r_n),$$

and this implies that $K(r) \in \Sigma$ for any $r \in \mathbb{R}$.

Consequently, the function f_0 is Σ -measurable.

Let $(g_n)_{n \geq 0}$ be an enumeration of the functions in H .

(b'''). Let us now define a continuous affine mapping $g: K \rightarrow \mathbb{R}^N$ by the formula

$$g(x) = (g_n(x))_{n \geq 0}, \quad x \in K.$$

Then $g(K) \subset \mathbb{R}^N$ is a metrizable compact convex subset of \mathbb{R}^N . Let us now prove that if $x, y \in K$ and $g(x) = g(y)$, then $f_0(x) = f_0(y)$.

Indeed, let us define

$$L(x) = \{z \in K; f_0(z) = f_0(x)\}.$$

Then $L(x) \in \Sigma$. Since Σ is the smallest σ -algebra of subset of K such that all the functions g_n , $n \geq 0$, be measurable, we have

$$z_1 \in K, z_0 \in L(x), g_n(z_1) = g_n(z_0), \forall n \geq 0 \Rightarrow z_1 \in L(x).$$

Consequently, since $x \in L(x)$, we have $y \in L(x)$, i.e., $f_0(y) = f_0(x)$.

(b'') From (b''') we conclude that by the formula

$$k(g(x)) = f_0(x), \quad x \in K,$$

we correctly define a function $k: g(K) \rightarrow \mathbb{R}$.

It is obvious that this function is concave on $g(K)$.

Now let $y_1 \in \text{ex}g(K)$. Then there exists an $x_1 \in \text{ex}K$ such that $g(x_1) = y_1$ and, therefore, we have

$$k(y_1) = k(g(x_1)) = f_0(x_1) > 0, \quad \forall y_1 \in \text{ex}g(K).$$

From

$$\begin{aligned} \{y \in g(K); k(y) \geq r\} &= \{g(x); x \in K, f_0(x) \geq r\} \\ &= g(\{x; x \in K, f_0(x) \geq r\}), \quad \forall r \in \mathbb{R}, \end{aligned}$$

and from the fact that the mapping g is closed we infer that the function k is upper semicontinuous on $b(K)$.

(b') Now let $\mu_0 \in \mathcal{M}_+^1(g(K))$ be any ($<$)-maximal or (\leq)-maximal measure which represents the point $g(x_0)$. We then have (by taking into account Corollary 2 to Proposition 2, and also Lemma 1)

$$f_0(x_0) = k(g(x_0)) \geq \int_{g(K)} k(y) d\mu_0(y) = \int_{\text{ex}g(K)} k(y) d\mu_0(y) > 0.$$

The theorem is proved.

5

In this section we shall give our generalization of the Bishop-de Leeuw and Choquet theorems.

Let $D \subset K$ be a Baire measurable subset; it is easy to prove that there exists a sequence $(h_n)_{n \in \mathbb{N}}$, $h_n \in A(K; \mathbb{R})$, $n \in \mathbb{N}$, such that $\|h_n\| \leq 1$, $\forall n \in \mathbb{N}$, and $x \in D$, $y \in K$, $h_n(x) = h_n(y)$, $\forall n \in \mathbb{N} \Rightarrow y \in D$.

Let us define $D_0 = D \cap \left(\bigcap_{n=0}^{\infty} S_{h_n^2} \right)$.

PROPOSITION 3. For any $x_0 \in D_0 \setminus (\text{ex} K)$ there exist $x_1, x_2 \in D_0$ such that $x_1 \neq x_0 \neq x_2$ and $x_0 = \frac{1}{2}(x_1 + x_2)$.

Proof. Since $x_0 \notin \text{ex} K$, there exist $x_1, x_2 \in K$ such that $x_1 \neq x_0 \neq x_2$ and $x_0 = \frac{1}{2}(x_1 + x_2)$. Let us prove that $x_1, x_2 \in D_0$.

Indeed, we have

$$\bar{h}_n^2(x_0) = h_n^2(x_0), \quad \forall n \in N,$$

and

$$\bar{h}_n^2(\lambda x_1 + (1-\lambda)x_2) \geq h_n^2(\lambda x_1 + (1-\lambda)x_2)$$

for any $n \in N$, and any $\lambda \in [0, 1]$.

Since \bar{h}_n^2 is concave and h_n^2 is convex, we infer that

$$(*) \quad \bar{h}_n^2(\lambda x_1 + (1-\lambda)x_2) = h_n^2(\lambda x_1 + (1-\lambda)x_2)$$

for any $n \in N$ and any $\lambda \in [0, 1]$. We then infer that the mapping

$$\lambda \mapsto h_n^2(\lambda x_1 + (1-\lambda)x_2), \quad \lambda \in [0, 1],$$

is affine for any $n \in N$. Since it is the square of an affine function, we deduce that the function

$$\lambda \mapsto h_n(\lambda x_1 + (1-\lambda)x_2), \quad \lambda \in [0, 1],$$

is constant on $[0, 1]$ for any $n \in N$. Consequently, we have

$$h_n(x_1) = h_n(x_0) = h_n(x_2), \quad \forall n \in N,$$

and this implies that

$$x_1, x_2 \in D.$$

On the other hand, from (*) we infer that

$$x_1, x_2 \in \bigcap_{n=0}^{\infty} S_{h_n^2}.$$

The proposition is proved.

THEOREM 3. (a) Any (\blacktriangleleft)-maximal measure $\mu \in \mathcal{M}_+^1(K)$ is pseudoconcentrated on $\text{ex} K$. In particular, we have

(b) (Choquet) Any ($<$)-maximal measure $\mu \in \mathcal{M}_+^1(K)$ is pseudoconcentrated on $\text{ex} K$.

(c) (Bishop-de Leeuw) Any (\ll)-maximal measure $\mu \in \mathcal{M}_+^1(K)$ is pseudoconcentrated on $\text{ex} K$.

Proof. (a) It will be sufficient to prove that for any compact, Baire measurable set $D \subset K$ such that

$$D \cap (\text{ex} K) = \emptyset,$$

we have $\mu(D) = 0$. Indeed, let us consider the set D_0 from the preceding proposition. We shall prove that $D_0 = \emptyset$, i.e.,

$$D \subset \bigcup_{n=0}^{\infty} (CS_{h_n^2}).$$

Corollary 2 to Proposition 2 will then imply that $\mu(D) = 0$.

Let us assume that $D_0 \neq \emptyset$. From the equality

$$\overline{\text{co}}(D_0) = \overline{\text{co}}(\bar{D}_0),$$

and from the Milman theorem we then infer that

$$\text{ex}(\overline{\text{co}}(D_0)) \subset \bar{D}_0 \subset D,$$

because D is compact. Let us consider the function

$$\varphi = \sum_{n=0}^{\infty} \frac{1}{2^n} (\bar{h}_n^2 - h_n^2),$$

where h_n , $n \in N$, are the functions already used in Proposition 3. It is obvious that φ is concave, finite and upper semicontinuous on K , hence on $\overline{\text{co}}(D_0)$.

If $\varphi(x) > 0$ for any $x \in \text{ex}(\overline{\text{co}}(D_0))$, then, by Theorem 2, we would infer that $\varphi(x) > 0$ for any $x \in D_0$, a contradiction. Consequently, there exists an $x_0 \in \text{ex}(\overline{\text{co}}(D_0))$ such that $\varphi(x_0) = 0$; hence, we have

$$x_0 \in D_0 \subset D.$$

From $D \cap (\text{ex} K) = \emptyset$ we infer that

$$x_0 \notin \text{ex} K;$$

Proposition 3 now implies that there exist $x_1, x_2 \in D_0 \subset \overline{\text{co}}(D_0)$ such that

$$x_1 \neq x_0 \neq x_2 \quad \text{and} \quad x_0 = \frac{1}{2}(x_1 + x_2),$$

thus contradicting the extremality of x_0 in $\overline{\text{co}}(D_0)$.

The theorem is proved.

Added in proof. N. Boboc informed the author that Theorem 2 first appeared in N. Boboc and Gh. Bucur, *Conuri convexe de funcții continue pe spații compacte*, Ed. Academiei R.S.R., Bucarest 1976, whereas Theorem 3 first appeared in N. Boboc and A. Cornea, *Convex cones of lower semicontinuous functions on compact spaces*, Rev. Roumaine Math. pures et appl. 12 (1967), 471–525. The proofs given to them here are, however, new.

References

- [1] E. M. Alfsen, *Compact convex sets and boundary integrals*, Springer Verlag, Berlin-Heidelberg-New York 1971 (Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 57).
- [2] H. Bauer, *Kennzeichnung kompakter Simplexe mit abgeschlossener Extrempunktmenge*, Arkiv der Math. 14 (6) (1963), 415–421.
- [3] E. Bishop and K. de Leeuw, *The representations of linear functionals by measures on sets of extreme points*, Ann. Inst. Fourier 9 (1959), 305–331.
- [4] G. Choquet, *Existence et unicité des représentations intégrales au moyen des points extrémaux dans les cônes convexes*, Séminaire Bourbaki (Déc. 1956), 139, 15 pp.
- [5] —, *Le théorème de représentation intégrale dans les ensembles convexes compacts*, Ann. Inst. Fourier 10 (1960), 333–344.
- [6] J. L. Doob, *Stochastic process*, John Wiley and Sons, New York; Chapman and Hall, London 1953.

- [7] P. A. Meyer, *Probability and potentials*, Blaisdell Publishing Company, Waltham, Toronto, London 1966.
- [8] R. R. Phelps, *Lectures on Choquet's theorem*, D. Van Nostrand Co., Princeton, Toronto, New York, London 1966.
- [9] C. F. Skau, *Orthogonal measures on the state space of a C*-algebra*, in: *Algebras in Analysis*, W. Williamson (ed.), Academic Press, 1975 (p. 272-303).
- [10] S. Telemán, *On reduction theory*, Rev. Roum. Math. Pures et Appl. 21 (4) (1976), 465-486.
- [11] —, *On the irreducible disintegration of the representations of C*-algebras*, Preprint Series in Mathematics, No. 8/1977, INCREST.
- [12] W. Wils, *Désintégration centrale des formes positives sur les C*-algèbres*, C. R. Acad. Sci., Paris, 267 (1968), 810-812.

Presented to the semester
 Spectral Theory
 September 23-December 16, 1977

ALGEBRAS OF GENERALIZED ANALYTIC FUNCTIONS

TOMA V. TONEV

*Institute of Mathematics, Bulgarian Academy of Sciences,
 Sofia, Bulgaria*

1

Let Γ be an additive and dense subgroup of the real axis \mathbb{R} . We shall consider that Γ is equipped with the discrete topology. Let $G = \hat{\Gamma}$ be the compact group of characters of Γ , i.e. the group of multiplicative functions χ on Γ with $|\chi| = 1$. According to the Pontrjagin duality theorem, the character group of G equipped with the sup-norm topology is isomorphic and homeomorphic to Γ , and we shall always identify $\Gamma = \{a\}_{a \in \Gamma}$ with the group $\hat{G} = \{\chi_a\}_{a \in \Gamma}$.

By the *generalized complex plane* we mean in the following the space $C_G = G \times [0, \infty) / G \times \{0\}$ with factor-topology. By $*$ we denote the image of $G \times \{0\}$ in C_G . We extend the characters χ_a of G on C_G as $\tilde{\chi}_a(\lambda, g) = \lambda^a \chi_a(g)$, if $\lambda \neq 0$, $a \neq 0$; $\tilde{\chi}_a(*) = 0$, $a \neq 0$; and $\tilde{\chi}_0(\lambda, g) \equiv 1$ if $\chi_0 \equiv 1$. Let K be a compact set in C_G . By $P_G(K)$ [$R_G(K)$], we denote the closure in $C(K)$ (with sup-norm) of the [ratios of] finite linear combinations of the extensions $\tilde{\chi}_a$, $a \geq 0$, of the characters χ_a , and by $A_G(K)$ the closure in $C(K)$ of the ratios of functions from $P_G(K)$. Note that in the case when K is the set $\bar{A}_G = G \times [0, 1] / G \times \{0\} \subset C_G$, the algebra $P_G(K)$ coincides with the algebra A_G of so called generalized analytic functions on G , introduced by Arens and Singer [1]. We use the term *generalized analytic functions* to denote the elements of $A_G(K)$. It is easy to see, that if Γ consists of rational numbers, then $R_G(K) = A_G(K)$. In this paper we show that algebras $P_G(K)$, $R_G(K)$ and $A_G(K)$, when $K \subset C_G$, preserve most of the properties of classical algebras $P(K)$, $R(K)$ and $A(K)$, $K \subset \mathbb{C}^n$.

2

Let K be a bounded set in C_G . The (generalized) *polynomial hull* \hat{K} of K we call as usual the set of these $(\lambda, g) \in C_G$, for which

$$|P(\lambda, g)| \leq \sup_K |P(\lambda, g)|,$$

where P is any finite linear combination of functions $\tilde{\chi}_a$, $a \geq 0$. A bounded set $K \subset C_G$ is called *polynomially convex*, if $\hat{K} = K$.