

References

- [1] W. G. Bade, *On Boolean algebras of projections and algebras of operators*, Trans. Amer. Math. Soc. 80 (1955), p. 345-359.
- [2] — *A multiplicity theory for Boolean algebras of projections in Banach spaces*, ibidem 92 (1958), p. 508-530.
- [3] M. M. Day, *Normed linear spaces*, Berlin 1962.
- [4] N. Dunford and J. Schwartz, *Linear operators I*, New York 1958.
- [5] J. Lindenstrauss and A. Pełczyński, *Absolutely summing operators in \mathcal{L}_p -spaces and their applications*, Studia Math. 29 (1968), p. 275-326.
- [6] J. Lindenstrauss and M. Zippin, *Banach spaces with a unique unconditional basis* (to appear).
- [7] — *Banach spaces with sufficiently many Boolean algebras of projections* (to appear).
- [8] G. Mackey, *Commutative Banach algebras*, Notes de Matematica 4, Rio de Janeiro.
- [9] C. A. McCarthy and L. Tzafriri, *Projections in \mathcal{L}_1 and \mathcal{L}_∞ spaces*, Pacific J. Math. (to appear).
- [10] A. Pełczyński, *Projections in certain Banach spaces*, Studia Math. 19 (1960), p. 209-228.
- [11] J. Werner, *Commuting spectral measures on Hilbert space*, Pacific J. Math. 4 (1954), p. 255-261.
- [12] M. Zippin, *On perfectly homogeneous bases in Banach spaces*, Israel J. Math. 4 (1966), p. 265-272.

UNIVERSITY OF MINNESOTA

Reçu par la Rédaction le 6. 4. 1968

Note on the class $L \log L$

by

E. M. STEIN (Princeton)

1. The purpose of this note is to prove two theorems. Each of these incidentally characterizes the class $L \log L$ in terms of the converse of some well-known inequality.

In Theorem 1 the setting is \mathbb{R}^n , and for a given integrable function $f(x)$, we define the maximal function $Mf(x)$ by

$$(1) \quad (Mf)(x) = \sup_{r>0} \frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y)| dy,$$

where $B(x, r)$ denotes the ball of radius r centered at x and $m(B(x, r))$ is its Lebesgue measure.

THEOREM 1. Suppose that f is integrable and is supported on some finite ball B . Then $\int_B Mf dx < \infty$ if and only if

$$\int_B |f| \log^+ |f| dx < \infty.$$

One direction, that $f \in L \log L$ implies $Mf \in L$, is very well known; but the converse although not really deeper, seems to have been overlooked all these years.

We shall also obtain a consequence of this result dealing with the Hilbert transform and its n -dimensional generalization, the Riesz transforms. The most appropriate setting for this will be periodic functions, i.e. those that satisfy $f(x+m) = f(x)$, where $m = (m_1, m_2, \dots, m_n)$ is any vector with integral coordinates. We denote by Q the "fundamental cube" — $\frac{1}{2} < x_j \leq \frac{1}{2}$, $j = 1, \dots, n$. Let

$$(2) \quad f \sim \sum_m a_m e^{2\pi i m \cdot x}$$

be the Fourier series of a periodic function integrable over Q , and let its Riesz transforms be given by

$$(3) \quad R_k(f) \sim i \sum_m' \frac{m_k}{|m|} a_m e^{2\pi i m \cdot x}, \quad k = 1, \dots, n.$$

When $n = 1$, R_1 is the usual (periodic) Hilbert transform.

THEOREM 2. Suppose $f \geq 0$, and f is integrable. Then all the $R_k f$ are also integrable ⁽¹⁾ if and only if

$$(4) \quad \int_0^\infty f \log^+ f dx < \infty.$$

Our main interest here, as above, is in the direction $R_k f \in L^1$ implies $f \in L\log L$. The converse was known previously (see Calderón and Zygmund [3]). So was the other direction when $n = 1$, but this was by an argument which seems to be restricted to that case. The argument given below has the additional virtue of showing Theorem 1 and Theorem 2 are really closely linked together.

2. Proof of Theorem 1. Suppose $f \in L^1(\mathbb{R}^n)$. Then it is possible to estimate the distribution function of Mf as follows. For any $a > 0$

$$(5) \quad m\{x: Mf(x) > a\} \leq \frac{A}{a} \int_{|f| > a/2} |f| dx.$$

This known inequality may be found in Weiner [5].

Our basic observation is that the estimate (5) may also be reversed. In fact, if we apply to $|f|$ and a the lemma of Calderón and Zygmund ([2], p. 91), we obtain a sequence $\{Q_j\}$ of disjoint cubes so that $|f(x)| \leq a$, if $x \notin UQ_j$; also

$$a < \frac{1}{m(Q_j)} \int_{Q_j} |f| dx \leq 2^n a.$$

Because of this if $x \in Q_j$, $Mf(x) > c^{-1}a$, with $c = 2^{n/2} \times \text{volume of the unit ball}$. So

$$\{x: Mf(x) > c^{-1}a\} \supset U_j Q_j,$$

and therefore

$$m\{x: Mf(x) > c^{-1}a\} \geq \sum m(Q_j) \geq (2^{-n}/a) \int_{U_j Q_j} |f| dx \geq (2^{-n}/a) \int_{|f| > a} |f| dx,$$

since $x \in UQ_j$, if $|f(x)| > a$. Passing from a to ca then gives

$$(6) \quad m\{x: Mf(x) > a\} \geq \frac{2^{-n} c^{-1}}{a} \int_{|f| > ca} |f| dx.$$

3. Let now B be any fixed finite ball and B' the ball with the same center as B with twice the diameter. We observe that if f is supported in B and Mf is integrable over B , then Mf is also integrable over B' . In

fact, suppose for simplicity that $B = \{x: |x| \leq 1\}$, and $B' = \{x: |x| \leq 2\}$. Then if $x \in B' - B$ (i.e. $1 < |x| \leq 2$), we can easily verify that $Mf(x) \leq cMf(\bar{x})$, where $\bar{x} = x/|x|^2$, which proves that Mf is also integrable over B' .

However if we are outside B' , we are at a positive distance from B and there Mf is bounded, and in fact decreases to zero as $|x| \rightarrow \infty$ (again since f is supported in B). These observations can be summarized in the following lemma:

LEMMA 1. Suppose f is supported in a finite ball B and Mf is integrable over that ball. Then for any fixed $\alpha_0, \alpha > 0$, Mf is integrable over the set where $(Mf)(x) > \alpha_0$.

According to the lemma then

$$\int_{\alpha_0}^{\infty} m\{x: Mf > a\} da < \infty$$

and by (6) this shows that $\int_B |f| \log^+ |f| dx < \infty$ which proves the theorem.

4. Proof of Theorem 2. For the proof of Theorem 2 we shall need some facts from the n -dimensional theory of H^p -spaces as developed by G. Weiss and the author [4], but in the periodic setting. The periodic analogues can be given very similar proofs as in the non-periodic case treated in [4] and so we shall content ourselves with the bare statement of the needed results.

Suppose both (2) and (3) are Fourier series of integrable functions. We form the $n+1$ harmonic functions $u_j(x, t)$ given by

$$(7) \quad u_0(x, t) = \sum a_m e^{-2\pi|m|t} e^{2\pi i m \cdot x},$$

$$(8) \quad u_k(x, t) = i \sum' a_m \frac{m_k}{|m|} e^{-2\pi|m|t} e^{2\pi i m \cdot x}, \quad k = 1, \dots, n.$$

The $u_j(x, t)$, $j = 0, \dots, n$, satisfy the generalized Cauchy-Riemann equations

$$\sum_{j=0}^n \frac{\partial u_j(x, t)}{\partial x_j} = 0 \quad (x_0 = t),$$

$$\frac{\partial u_k(x, t)}{\partial x_j} = \frac{\partial u_j(x, t)}{\partial x_k}, \quad 0 \leq k, j \leq n.$$

Let

$$F(x, t) = (u_0(x, t), u_1(x, t), \dots, u_n(x, t)).$$

Then we say that $F \in H^1$ if

$$\sup_{t>0} \int_Q |F(x, t)| dx \leq M < \infty.$$

⁽¹⁾ More precisely, the series (3) are Fourier series of integrable functions.

The result we shall use is the following

LEMMA 2. Suppose $F \in H^1$. Then

$$\int_Q \sup_{t>0} |F(x, t)| dx \leq A \sup_{t>0} \int_Q |F(x, t)| dx.$$

(For the non-periodic analogue see [4], pp. 46 and 47).

5. We shall also need to make a few remarks about the Poisson kernel.

Let

$$P(x, t) = \frac{c_n}{(|x|^2 + t^2)^{(n+1)/2}}, \quad c_n = \frac{\Gamma((n+1)/2)}{\pi^{(n+1)/2}}$$

be the n -dimensional (non-periodic) Poisson kernel. Its periodic analogue $\mathcal{P}(x, t)$ is

$$\mathcal{P}(x, t) = \sum_m P(x+m, t).$$

It can be seen that

$$(9a) \quad \mathcal{P}(x, t) \geq 0, \quad \text{in fact } \mathcal{P}(x, t) \geq P(x+m, t);$$

$$(9b) \quad \int_Q \mathcal{P}(x, t) dx = 1;$$

$$(9c) \quad \mathcal{P}(x, t) = \sum_m e^{-2\pi|m|t} e^{2\pi im \cdot x}.$$

The only statement that really needs proof is (9c), but this is a consequence of the Poisson summation formula (see e.g. [1], p. 32).

6. We shall now prove that part of the theorem that concerns us. Suppose that f and the $R_k(f)$ given by (3) are in $L^1(Q)$. Since

$$u_0(x, t) = \int f(y) \mathcal{P}(x-y, t) dy$$

and

$$u_k(x, t) = \int (R_k f)(y) \mathcal{P}(x-y, t) dy,$$

we see that $F = (u_0, u_1, \dots, u_n) \in H^1$.

Next observe that if g is any non-negative function on \mathbf{R}^n which is integrable, then

$$(10) \quad \sup_{t>0} \int_{\mathbf{R}^n} g(y) P(x-y, t) dy \geq \sup_{t>0} \int_{|x-y|<t} g(y) P(x-y, t) dy \\ \geq \sup_{t>0} b t^{-n} \int_{|x-y|<t} g(y) dy = b' M(g)(x).$$

Now we have already noted the fact that $f, R_k(f) \in L'(Q)$ implies that $F \in H^1$. Thus by Lemma 2

$$\int_Q \sup_{t>0} |F(x, t)| dx < \infty$$

and so since $f \geq 0$ and $\mathcal{P}(x, t) \geq P(x, t)$, we get by (10)

$$\int_Q M g(x) dx < \infty,$$

where $g = f$ for $x \in Q$, and $g = 0$, $x \notin Q$. Similarly, since $\mathcal{P}(x, t) \geq P(x+m, t)$, we also have

$$\int_{Q+m} M g dx < \infty$$

for any vector m with integral components. Finally, suppose B is a fixed ball which contains Q . Then B is contained in the union of finitely many integral translates of Q . So

$$\int_B M g dx < \infty,$$

which by Theorem 1 implies that

$$\int_B g \log^+ g dx = \int_Q f \log^+ f dx < \infty.$$

7. Addendum. The following non-periodic analogue of theorem 2, is stated without proof. Suppose $f \in L^p(\mathbf{R}^n)$ for some p , $1 \leq p < \infty$. Then the non-periodic Riesz transform R_k is defined by

$$R_k(f)(x) = \lim_{\epsilon \rightarrow 0} \int_{|y| \geq \epsilon} f(x-y) \frac{y_k}{|y|^{n+1}} dy.$$

The limit exists almost everywhere, and if in addition $p > 1$, it also exists in L^p -norm.

Suppose now B_1 and B_2 are a pair of finite open balls so that $\bar{B}_1 \subset B_2$.

THEOREM 3. (a) Suppose $\int_{B_2} |f| \log^+ |f| dx < \infty$. Then

$$\int_{B_1} |R_k f| dx < \infty.$$

(b) Conversely, suppose $\int_{B_2} |R_k f| dx < \infty$, $k = 1, \dots, n$, and $f \geq 0$ on B_2 . Then

$$\int_{B_1} |f| \log^+ |f| dx < \infty.$$

Part (a) was known previously, see [3]. Part (b) is proved by an adaptation of the argument used for the proof of the periodic case, and may be left as an exercise to the interested reader.

The reason that this non-periodic version seems awkward in comparison to its periodic analogue is the fact that if $f \geq 0$ and $f \neq 0$, $f \in L^1(\mathbb{R}^n)$, then $R_k(f)$ never belongs to $L^1(\mathbb{R}^n)$.

The author profitted from several valuable conversations with A. Zygmund on the contents of this note.

References

- [1] S. Bochner, *Harmonic analysis and the theory of probability*, Berkeley 1955.
- [2] A. P. Calderón and A. Zygmund, *On the existence of certain singular integrals*, Acta Math. 88 (1952), p. 85-139.
- [3] — *Singular integrals and periodic functions*, Studia Math. 14 (1954), p. 249-271.
- [4] E. M. Stein and G. Weiss, *On the theory of harmonic functions of several variables I*, Acta Math. 103 (1960), p. 25-62.
- [5] N. Wiener, *The ergodic theorem*, Duke Math. J. 5 (1939), p. 1-18.

Reçu par la Rédaction le 27. 4. 1968
