

INVERSE PROBLEM ON THE STEINER WIENER INDEX

XUELIANG LI¹

Center for Combinatorics and LPMC-TJKLC
Nankai University, Tianjin 300071, China

e-mail: lxl@nankai.edu.cn

YAPING MAO²

Department of Mathematics
Qinghai Normal University, Qinghai 810008, China

e-mail: maoyaping@ymail.com

AND

IVAN GUTMAN

Faculty of Science P.O. Box 60
34000 Kragujevac, Serbia, and
State University of Novi Pazar, Novi Pazar, Serbia

e-mail: gutman@kg.ac.rs

Abstract

The Wiener index $W(G)$ of a connected graph G , introduced by Wiener in 1947, is defined as $W(G) = \sum_{u,v \in V(G)} d_G(u,v)$, where $d_G(u,v)$ is the distance (the length a shortest path) between the vertices u and v in G . For $S \subseteq V(G)$, the *Steiner distance* $d(S)$ of the vertices of S , introduced by Chartrand *et al.* in 1989, is the minimum size of a connected subgraph of G whose vertex set contains S . The k -th *Steiner Wiener index* $SW_k(G)$ of G is defined as $SW_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} d(S)$. We investigate the following problem: Fixed a positive integer k , for what kind of positive integer w does there exist a connected graph G (or a tree T) of order $n \geq k$ such that $SW_k(G) = w$ (or $SW_k(T) = w$)? In this paper, we give some solutions to this problem.

Keywords: distance, Steiner distance, Wiener index, Steiner Wiener index.

2010 Mathematics Subject Classification: 05C05, 05C12, 05C35.

¹Supported by NSFC No. 11371205.

²Supported by NSFC Nos. 11601254, 11661068, 11551001, 11161037 and SFQP Nos. 2016-ZJ-948Q, 2014-ZJ-907.

1. INTRODUCTION

All graphs in this paper are assumed to be undirected, finite and simple. We refer to [3] for graph theoretical notation and terminology not specified here. A distance is one of basic concepts of graph theory [4]. If G is a connected graph and $u, v \in V(G)$, then the *distance* $d(u, v) = d_G(u, v)$ between u and v is the length of a shortest path connecting u and v . For more details on this subject, see [13].

The *Wiener index* $W(G)$ of a connected graph G is defined by

$$W(G) = \sum_{u, v \in V(G)} d_G(u, v).$$

Mathematicians have studied this graph invariant since the 1970s in [11]; for details see the surveys [10, 33], the recent papers [2, 7, 14, 17, 15, 20] and the references cited therein. Information on chemical applications of the Wiener index can be found in [27, 28].

The Steiner distance of a graph, introduced by Chartrand *et al.* in [6] in 1989, is a natural and nice generalization of the concept of the classical graph distance. For a graph $G = (V, E)$ and a set $S \subseteq V$ of at least two vertices, an *S-Steiner tree* or a *Steiner tree connecting S* (or simply, an *S-tree*) is a subgraph $T = (V', E')$ of G that is a tree with $S \subseteq V'$. Let G be a connected graph of order at least 2 and let S be a nonempty set of vertices of G . Then the *Steiner distance* $d(S)$ among the vertices of S (or simply the distance of S) is the minimum size of a connected subgraph whose vertex set contains S . Note that if H is a connected subgraph of G such that $S \subseteq V(H)$ and $|E(H)| = d(S)$, then H is a tree. Clearly, $d(S) = \min\{|E(T)| : S \subseteq V(T)\}$, where T is a subtree of G . Furthermore, if $S = \{u, v\}$, then $d(S) = d(u, v)$ is nothing new, but the classical distance between u and v . Clearly, if $|S| = k$, then $d(S) \geq k - 1$. For more details on Steiner distance, we refer to [1, 5, 6, 8, 13, 26].

In [23], we proposed a generalization of the Wiener index concept, using Steiner distance. Thus, the *k-th Steiner Wiener index* $SW_k(G)$ of a connected graph G is defined by

$$SW_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} d(S).$$

For $k = 2$, the Steiner Wiener index coincides with the ordinary Wiener index. It is usual to consider SW_k for $2 \leq k \leq n - 1$, but the above definition implies $SW_1(G) = 0$ and $SW_n(G) = n - 1$ for a connected graph G of order n . For more details on Steiner Wiener index, we refer to [23, 24, 25].

A chemical application of SW_k was recently reported in [16].

It should be noted that in the 1990s, Dankelmann *et al.* in [8, 9] studied the *average Steiner distance*, which is related to our Steiner Wiener index via $SW_k(G)/\binom{n}{k}$.

The seemingly elementary question: “Which natural numbers are Wiener indices of graphs?” was much investigated in the past; see [12, 19, 21, 29, 31, 32]. We now consider the analogous question for Steiner Wiener indices.

Problem. Fixed a positive integer k , for what kind of positive integer w does there exist a connected graph G (or a tree T) of order $n \geq k$ such that $SW_k(G) = w$ (or $SW_k(T) = w$)?

For $k = 2$, the authors have nice results in [30, 32], completely solved a conjecture by Lepović and Gutman [22] for trees, which states that for all but 49 positive integers w one can find a tree with Wiener index w . This is different from our problem for trees, since here we consider graphs or trees with order n .

2. THE CASES $k = n$ AND $k = n - 1$

At first, let us consider the case $k = n$.

If G is a connected graph or a tree of order n , then for $k = n$, $SW_k(G) = n - 1$. Thus the following result is immediate.

Proposition 2.1. *For a positive integer w , there exists a connected graph G or a tree T of order n such that $SW_n(G) = w$ or $SW_n(T) = w$ if and only if $w = n - 1$.*

For the case $k = n - 1$, we need the following results in [23].

Lemma 2.2 [23]. *Let T be a tree of order n , possessing p pendant vertices. Then*

$$SW_{n-1}(T) = n(n - 1) - p$$

irrespective of any other structural detail of T .

Lemma 2.3 [23]. *Let K_n be the complete graph of order n , and let k be an integer such that $2 \leq k \leq n$. Then*

$$SW_k(K_n) = \binom{n}{k}(k - 1).$$

Lemma 2.4 [23]. *Let G be a connected graph of order n , and let k be an integer such that $2 \leq k \leq n$. Then*

$$\binom{n}{k}(k - 1) \leq SW_k(G) \leq (k - 1)\binom{n + 1}{k + 1}.$$

Moreover, the lower bound is sharp.

From the previous results, we can derive the following proposition.

Proposition 2.5. *For a positive integer w , there exists a connected graph G of order n such that $SW_{n-1}(G) = w$ if and only if $n^2 - 2n \leq w \leq n^2 - n - 2$.*

Proof. By Lemma 2.4, if G is a connected graph of order n , then

$$n(n-2) \leq SW_{n-1}(G) \leq (n+1)(n-2).$$

Therefore, $n^2 - 2n \leq w \leq n^2 - n - 2$.

By Lemma 2.3, $SW_{n-1}(K_n) = n^2 - 2n$.

Let T be a tree of order n with p pendant vertices with $2 \leq p \leq n-1$. By Lemma 2.2, $SW_{n-1}(T) = n^2 - n - p$, and thus for any integer w with $n^2 - n - (n-1) \leq w \leq n^2 - n - 2$, there exists a tree T of order n such that $SW_{n-1}(T) = w$. ■

From the proof of Proposition 2.5 the next result immediately follows.

Proposition 2.6. *For a positive integer w , there exists a tree T of order n such that $SW_{n-1}(T) = w$ if and only if $n^2 - 2n + 1 \leq w \leq n^2 - n - 2$.*

3. THE CASE $k = n - 2$

Similarly to Lemma 2.2, we can derive the following result.

Lemma 3.1. *Let T be a tree of order n , possessing p pendant vertices. Let q be the number of vertices of degree 2 in T that are adjacent to a pendant vertex. Then*

$$(1) \quad SW_{n-2}(T) = \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q).$$

Proof. For any $S \subseteq V(T)$ and $|S| = n-2$, let $\bar{S} = \{u, v\}$. If $d_T(u) = d_T(v) = 1$, then $d_T(S) = n-3$, and this case contributes to SW_{n-2} by

$$\sum_{\substack{u, v \in \bar{S} \\ d_T(u)=d_T(v)=1}} d_T(S) = \binom{p}{2}(n-3).$$

If $d_T(u) \geq 2$ and $d_T(v) \geq 2$, then $d_T(S) = n-1$, and this case contributes to SW_{n-2} by

$$\sum_{\substack{u, v \in \bar{S} \\ d_T(u) \geq 2, d_T(v) \geq 2}} d_T(S) = \binom{n-p}{2}(n-1).$$

Suppose that $d_T(u) = 1$ and $d_T(v) \geq 2$. If $d_T(u) = 1$, $d_T(v) = 2$ and $uv \in E(G)$, then $d_T(S) = n-3$. If $d_T(u) = 1$, $d_T(v) \geq 3$ and $uv \in E(T)$, then

$d_T(S) = n - 2$. If $d_T(u) = 1$, $d_T(v) \geq 2$ and $uv \notin E(T)$, then $d_T(S) = n - 2$. Therefore, this case contributes to SW_{n-2} by

$$\begin{aligned} \sum_{\substack{u,v \in \bar{S} \\ d_T(u)=1, d_T(v) \geq 2}} d_T(S) &= \sum_{\substack{u,v \in \bar{S}, uv \in E(T) \\ d_T(u)=1, d_T(v)=2}} d_T(S) + \sum_{\substack{u,v \in \bar{S}, uv \in E(T) \\ d_T(u)=1, d_T(v) \geq 3}} d_T(S) + \sum_{\substack{u,v \in \bar{S}, uv \notin E(T) \\ d_T(u)=1, d_T(v) \geq 2}} d_T(S) \\ &= q(n-3) + (p-q)(n-2) + p(n-p-1)(n-2). \end{aligned}$$

From the above argument, we have

$$\begin{aligned} SW_{n-2}(T) &= \binom{p}{2}(n-3) + \binom{n-p}{2}(n-1) + q(n-3) \\ &\quad + (p-q)(n-2) + p(n-p-1)(n-2) \\ &= \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q). \quad \blacksquare \end{aligned}$$

Li *et al.* obtained the following sharp lower and upper bounds of $SW_k(T)$ for a tree T .

Lemma 3.2 [23]. *Let T be a tree of order n , and let k be an integer such that $2 \leq k \leq n$. Then*

$$\binom{n-1}{k-1}(n-1) \leq SW_k(T) \leq (k-1) \binom{n+1}{k+1}.$$

Moreover, among all trees of order n , the star S_n minimizes the Steiner Wiener k -index, whereas the path P_n maximizes the Steiner Wiener k -index.

For trees, we have the following result.

Theorem 3.3. *For a positive integer w , there exists a tree T of order n ($n \geq 5$), possessing p pendant vertices, such that $SW_{n-2}(T) = w$ if and only if $w = \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q)$, where q is the number of vertices of degree 2 in T that are adjacent to a pendant vertex, and one of the following holds:*

- (1) $2 \leq q \leq \lfloor \frac{n-1}{2} \rfloor$ and $q \leq p \leq n - q - 1$;
- (2) $q = 1$ and $3 \leq p \leq n - 2$;
- (3) $q = 0$ and $4 \leq p \leq n - 1$.

Proof. Suppose that $w = \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q)$, where $0 \leq q \leq \lfloor \frac{n-1}{2} \rfloor$, $q \leq p \leq n - q - 1$. Let $K_{1,p-1}$ be a star of order p , and let v be the center of $K_{1,p-1}$. Then $K_{1,p-1}^*$ is a graph obtained from $K_{1,p-1}$ by picking up $q-1$ edges and then replacing each of them by a path of length 2. Note that $K_{1,p-1}^*$ is a

subdivision of $K_{1,p-1}$. Let G be a graph obtained by $K_{1,p-1}^*$ and a path $P_{n-p-q+2}$ by identifying v and one endvertex of the path. Clearly, G is a tree of order n with p pendant vertices, and there are exactly q vertices of degree 2 in T such that each of them is adjacent to a pendant vertex. From Lemma 3.1, we have $SW_{n-2}(T) = \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q) = w$, as desired.

Conversely, for any tree T of order n ($n \geq 5$) with p pendant vertices, from Lemma 3.1, $SW_{n-2}(T) = \frac{1}{2}(n^3 - 2n^2 + n - 2np + 2p - 2q)$. We now show that p, q satisfy one of (1), (2), (3). Clearly, $p \geq 2$, $0 \leq q \leq \lfloor \frac{n-1}{2} \rfloor$ and $q \leq p$.

Claim 1. $p + q \leq n - 1$.

Proof. Assume, to the contrary, that $p + q = n$. Then T is a path of order n . Since $n \geq 5$, it follows that there exists a vertex of degree 2 having no adjacent pendant vertex, which contradicts to $p + q = n$. \square

If $q \geq 2$, then it follows from Claim 1 and $q \leq p$ that $q \leq p \leq n - q - 1$. If $q = 1$, then it follows from Claim 1 that $2 \leq p \leq n - 2$. Furthermore, if $p = 2$, then T is a path of order n . Since $n \geq 5$, it follows that $q = 2$, a contradiction. If $q = 0$, then it follows from Claim 1 that $2 \leq p \leq n - 1$. Furthermore, if $p = 2$, then T is a path of order n . Since $n \geq 5$, it follows that $q = 2$, a contradiction. If $p = 3$, then T is a tree of order n . Since $n \geq 5$, it follows that $q \geq 1$, a contradiction. \blacksquare

4. THE CASE FOR GENERAL k

For trees, we have the following result.

Theorem 4.1. *Let T be a graph obtained from a path P_t and a star S_{n-t+1} by identifying a pendant vertex of P_t and the center v of S_{n-t+1} , where $1 \leq t \leq n - 1$ and $k \leq n$. Then*

$$SW_k(T) = t \binom{n-1}{k} - \binom{t}{k+1} - \binom{n}{k+1} + \binom{n-t+1}{k+1} + (k-1) \binom{n}{k}.$$

Proof. For any $S \subseteq V(T)$ and $|S| = k$, if $S \subseteq V(S_{n-t+1}) - v$, then $d_G(S) = k$. There are $\binom{n-t}{k}$ such subsets, contributing to SW_k by $k \binom{n-t}{k}$. If $S \subseteq V(P_t)$, then it contributes to SW_k by $(k-1) \binom{t+1}{k+1}$ from Lemma 3.2. Suppose that $S \cap V(P_t) \neq \emptyset$ and $S \cap (V(S_{n-t+1}) - v) \neq \emptyset$. Let $|S \cap V(S_{n-t+1} - v)| = i$, $|S \cap V(P_t)| = k - i$ and $P_t = u_1 u_2 \cdots u_t$, where $v = u_1$. Without loss of generality, let $S \cap V(P_t) = \{u_{j_1}, u_{j_2}, \dots, u_{j_{k-i}}\}$, where $1 \leq j_1 < j_2 < \cdots < j_{k-i} \leq t$. Then $k - i \leq j_{k-i} \leq t$. Let $j_{k-i} = j$. Then $d_G(S) = i + j - 1$, and $k - i \leq j \leq t$. Once the vertex u_j is chosen, we have $\binom{j-2}{k-i-1}$ ways to choose $u_{j_1}, u_{j_2}, \dots, u_{j_{k-i-1}}$. In

this case, we contribute to SW_k by

$$X = \sum_{i=1}^{k-1} \binom{n-t}{i} \left[\sum_{j=k-i}^t \binom{j-1}{k-i-1} (j+i-1) \right].$$

Since

$$\begin{aligned} \binom{j-1}{k-i-1} (j+i-1) &= \binom{j-1}{k-i-1} j + \binom{j-1}{k-i-1} (i-1) \\ &= (k-i) \binom{j}{k-i} + (i-1) \binom{j-1}{k-i-1}, \end{aligned}$$

it follows that

$$\begin{aligned} \sum_{j=k-i}^t \binom{j-1}{k-i-1} (j+i-1) &= (k-i) \sum_{j=k-i}^t \binom{j}{k-i} + (i-1) \sum_{j=k-i}^t \binom{j-1}{k-i-1} \\ &= (k-i) \binom{t+1}{k-i+1} + (i-1) \binom{t}{k-i}, \end{aligned}$$

and hence

$$\begin{aligned} X &= \sum_{i=1}^{k-1} \binom{n-t}{i} \left[\sum_{j=k-i}^t \binom{j-1}{k-i-1} (j+i-1) \right] \\ &= \sum_{i=1}^{k-1} \binom{n-t}{i} \left[(k-i) \binom{t+1}{k-i+1} + (i-1) \binom{t}{k-i} \right] \\ &= \sum_{i=1}^{k-1} \binom{n-t}{i} (k-i) \binom{t+1}{k-i+1} + \sum_{i=1}^{k-1} \binom{n-t}{i} (i-1) \binom{t}{k-i} \\ &= \sum_{i=1}^{k-1} (k-i) \binom{t}{k-i+1} \binom{n-t}{i} + \sum_{i=1}^{k-1} (k-i) \binom{t}{k-i} \binom{n-t}{i} \\ &\quad + \sum_{i=1}^{k-1} (i-1) \binom{t}{k-i} \binom{n-t}{i} \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^{k-1} (k-i) \binom{t}{k-i+1} \binom{n-t}{i} + (k-1) \sum_{i=1}^{k-1} \binom{t}{k-i} \binom{n-t}{i} \\
&= \sum_{i=1}^{k-1} (k-i) \binom{t}{k-i+1} \binom{n-t}{i} + (k-1) \left[\binom{n}{k} - \binom{t}{k} - \binom{n-t}{k} \right].
\end{aligned}$$

Let

$$Y = \sum_{i=1}^{k-1} (k-i) \binom{t}{k-i+1} \binom{n-t}{i}.$$

Then

$$\begin{aligned}
Y &= \sum_{i=1}^{k-1} (k-i+1) \binom{t}{k-i+1} \binom{n-t}{i} - \sum_{i=1}^{k-1} \binom{t}{k-i+1} \binom{n-t}{i} \\
&= t \sum_{i=1}^{k-1} \binom{t-1}{k-i} \binom{n-t}{i} - \sum_{i=1}^{k-1} \binom{t}{k+1-i} \binom{n-t}{i} \\
&= t \left[\binom{n-1}{k} - \binom{t-1}{k} - \binom{n-t}{k} \right] \\
&\quad - \left[\binom{n}{k+1} - \binom{t}{k+1} - t \binom{n-t}{k} - \binom{n-t}{k+1} \right],
\end{aligned}$$

and hence

$$\begin{aligned}
SW_k(T) &= (k-1) \binom{t+1}{k+1} + k \binom{n-t}{k} + X \\
&= (k-1) \binom{t+1}{k+1} + k \binom{n-t}{k} + Y + (k-1) \left[\binom{n}{k} - \binom{t}{k} - \binom{n-t}{k} \right] \\
&= (k-1) \binom{t+1}{k+1} + k \binom{n-t}{k} + t \left[\binom{n-1}{k} - \binom{t-1}{k} - \binom{n-t}{k} \right] \\
&\quad - \left[\binom{n}{k+1} - \binom{t}{k+1} - t \binom{n-t}{k} - \binom{n-t}{k+1} \right] \\
&\quad + (k-1) \left[\binom{n}{k} - \binom{t}{k} - \binom{n-t}{k} \right]
\end{aligned}$$

$$\begin{aligned}
 &= (k-1) \binom{t}{k+1} + (k-1) \binom{t}{k} + k \binom{n-t}{k} + t \binom{n-1}{k} - t \binom{t-1}{k} \\
 &\quad - t \binom{n-t}{k} - \binom{n}{k+1} + \binom{t}{k+1} + t \binom{n-t}{k} + \binom{n-t}{k+1} \\
 &\quad + (k-1) \binom{n}{k} - (k-1) \binom{t}{k} - (k-1) \binom{n-t}{k} \\
 &= (k-1) \binom{t}{k+1} + k \binom{n-t}{k} + t \binom{n-1}{k} - t \binom{t-1}{k} \\
 &\quad - \binom{n}{k+1} + \binom{t}{k+1} + \binom{n-t}{k+1} + (k-1) \binom{n}{k} - (k-1) \binom{n-t}{k} \\
 &= k \binom{t}{k+1} + \binom{n-t}{k} + t \binom{n-1}{k} - t \binom{t-1}{k} - \binom{n}{k+1} + \binom{n-t}{k+1} \\
 &\quad + (k-1) \binom{n}{k} \\
 &= k \binom{t}{k+1} + t \binom{n-1}{k} - t \binom{t-1}{k} - \binom{n}{k+1} + \binom{n-t+1}{k+1} + (k-1) \binom{n}{k} \\
 &= t \binom{n-1}{k} - \binom{t}{k+1} - \binom{n}{k+1} + \binom{n-t+1}{k+1} + (k-1) \binom{n}{k}. \quad \blacksquare
 \end{aligned}$$

The following corollary is immediate from Theorem 4.1.

Corollary 4.2. *For a positive integer w , there exists a tree T of order n such that $SW_k(T) = w$ if*

$$w = t \binom{n-1}{k} - \binom{t}{k+1} - \binom{n}{k+1} + \binom{n-t+1}{k+1} + (k-1) \binom{n}{k},$$

where $1 \leq t \leq n-1$ and $k \leq n$.

For general graphs, we have the following.

Theorem 4.3. *Let G be a graph obtained from a clique K_{n-r} and a star S_{r+1} by identifying a vertex of K_{n-r} and the center v of S_{r+1} . For $k \leq r \leq n-1-k$,*

$$SW_k(G) = (n-1) \binom{n-1}{k-1} - \binom{n-r-1}{k}.$$

Proof. For any $S \subseteq V(G)$ and $|S| = k$, if $S \subseteq V(K_{n-r})$, then $d_G(S) = k - 1$. There are $\binom{n-r}{k}$ such subsets, contributing to SW_k by $(k-1)\binom{n-r}{k}$. If $S \subseteq V(S_{r+1}) - v$, then $d_G(S) = k$. There are $\binom{r}{k}$ such subsets, contributing to SW_k by $k\binom{r}{k}$. Suppose that $S \cap V(K_{n-r}) \neq \emptyset$ and $S \cap (V(S_{r+1}) - v) \neq \emptyset$. If $v \in S$, then $d_G(S) = k - 1$. There are $\binom{n-r-1}{k-x-1}\binom{r}{x}$ such subsets, contributing to SW_k by $(k-1)\sum_{x=1}^{k-1}\binom{n-r-1}{k-x-1}\binom{r}{x}$. If $v \notin S$, then $d_G(S) = k$. There are $\binom{n-r-1}{k-x}\binom{r}{x}$ such subsets, contributing to SW_k by $k\sum_{x=1}^{k-1}\binom{n-r-1}{k-x}\binom{r}{x}$. Then

$$\begin{aligned}
& SW_k(G) \\
&= (k-1)\binom{n-r}{k} + k\binom{r}{k} + (k-1)\sum_{x=1}^{k-1}\binom{n-r-1}{k-x-1}\binom{r}{x} \\
&\quad + k\sum_{x=1}^{k-1}\binom{n-r-1}{k-x}\binom{r}{x} \\
&= (k-1)\binom{n-r}{k} + k\binom{r}{k} + (k-1)\left[\binom{n-1}{k-1} - \binom{n-1-r}{k-1}\right] \\
&\quad + k\left[\binom{n-1}{k} - \binom{n-1-r}{k} - \binom{r}{k}\right] \\
&= (k-1)\binom{n-r}{k} + (k-1)\left[\binom{n-1}{k-1} - \binom{n-1-r}{k-1}\right] \\
&\quad + k\left[\binom{n-1}{k} - \binom{n-1-r}{k}\right] \\
&= (k-1)\binom{n-r}{k} + (n-1)\binom{n-1}{k-1} - (k-1)\binom{n-1-r}{k-1} - k\binom{n-1-r}{k} \\
&= (n-1)\binom{n-1}{k-1} + (k-1)\binom{n-r-1}{k} - k\binom{n-1-r}{k} \\
&= (n-1)\binom{n-1}{k-1} - \binom{n-1-r}{k},
\end{aligned}$$

as desired. ■

The following corollary is immediate from Theorems 4.1 and 4.3.

Corollary 4.4. *For a positive integer w , there exists a connected graph G of order n such that $SW_k(G) = w$ if w satisfies one of the following conditions.*

- (1) $w = t \binom{n-1}{k} - \binom{t}{k+1} - \binom{n}{k+1} + \binom{n-t+1}{k+1} + (k-1) \binom{n}{k}$, where $1 \leq t \leq n-1$ and $k \leq n$.
- (2) $w = (n-1) \binom{n-1}{k-1} - \binom{n-r-1}{k}$, where $k \leq r \leq n-1-k$ and $k \leq n$.

Acknowledgement

The authors are very grateful to the referees for their valuable comments and suggestions.

REFERENCES

- [1] P. Ali, P. Dankelmann and S. Mukwembi, *Upper bounds on the Steiner diameter of a graph*, Discrete Appl. Math. **160** (2012) 1845–1850.
doi:10.1016/j.dam.2012.03.031
- [2] E.O.D. Andriantiana, S. Wagner and H. Wang, *Maximum Wiener index of trees with given segment sequence*, MATCH Commun. Math. Comput. Chem. **75** (2016) 91–104.
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory* (Springer, New York, 2008).
- [4] F. Buckley and F. Harary, *Distance in Graphs* (Addison-Wesley, Redwood, 1990).
- [5] J. Cáceres, A. Márquez and M.L. Puertas, *Steiner distance and convexity in graphs*, European J. Combin. **29** (2008) 726–736.
doi:10.1016/j.ejc.2007.03.007
- [6] G. Chartrand, O.R. Oellermann, S.L. Tian and H.B. Zou, *Steiner distance in graphs*, Časopis Pest. Mat. **114** (1989) 399–410.
- [7] L. Chen, X. Li and M. Liu, *The (revised) Szeged index and the Wiener index of a nonbipartite graph*, European J. Combin. **36** (2014) 237–246.
doi:10.1016/j.ejc.2013.07.019
- [8] P. Dankelmann, O.R. Oellermann and H.C. Swart, *The average Steiner distance of a graph*, J. Graph Theory **22** (1996) 15–22.
doi:10.1002/(SICI)1097-0118(199605)22:1<15::AID-JGT3>3.0.CO;2-O
- [9] P. Dankelmann, O.R. Oellermann and H.C. Swart, *On the average Steiner distance of graphs with prescribed properties*, Discrete Appl. Math. **79** (1997) 91–103.
doi:10.1016/S0166-218X(97)00035-8
- [10] A. Dobrynin, R. Entringer and I. Gutman, *Wiener index of trees: theory and applications*, Acta Appl. Math. **66** (2001) 211–249.
doi:10.1023/A:1010767517079
- [11] R.C. Entringer, D.E. Jackson and D.A. Snyder, *Distance in graphs*, Czechoslovak Math. J. **26** (1976) 283–296.
- [12] J. Fink, B. Lužar and R. Škrekovski, *Some remarks on inverse Wiener index problem*, Discrete Appl. Math. **160** (2012) 1851–1858.
doi:10.1016/j.dam.2012.02.028

- [13] W. Goddard and O.R. Oellermann, *Distance in graphs*, in: M. Dehmer (Ed.), *Structural Analysis of Complex Networks* (Birkhäuser, Dordrecht, 2011) 49–72.
doi:10.1007/978-0-8176-4789-6_3
- [14] M. Goubko, *Minimizing Wiener index for vertex-weighted trees with given weight and degree sequences*, *MATCH Commun. Math. Comput. Chem.* **75** (2016) 3–27.
- [15] I. Gutman, R. Cruz and J. Rada, *Wiener index of Eulerian graphs*, *Discrete Appl. Math.* **162** (2014) 247–250.
doi:10.1016/j.dam.2013.08.024
- [16] I. Gutman, B. Furtula and X. Li, *Multicenter Wiener indices and their applications*, *J. Serb. Chem. Soc.* **80** (2015) 1009–1017.
doi:10.2298/JSC150126015G
- [17] I. Gutman, K. Xu and M. Liu, *A congruence relation for Wiener and Szeged indices*, *Filomat* **29** (2015) 1081–1083.
doi:10.2298/FIL1505081G
- [18] I. Gutman and Y.N. Yeh, *The sum of all distances in bipartite graphs*, *Math. Slovaca* **45** (1995) 327–334.
- [19] I. Gutman, Y.N. Yeh and J.C. Chen, *On the sum of all distances in graphs*, *Tamkang J. Math.* **25** (1994) 83–86.
- [20] M. Knor and R. Škrekovski, *Wiener index of generalized 4-stars and of their quadratic line graphs*, *Australas. J. Combin.* **58** (2014) 119–126.
- [21] M. Krnc and R. Škrekovski, *On Wiener inverse interval problem*, *MATCH Commun. Math. Comput. Chem.* **75** (2016) 71–80.
- [22] M. Lepović and I. Gutman, *A collective property of trees and chemical trees*, *J. Chem. Inf. Comput. Sci.* **38** (1998) 823–826.
doi:10.1021/ci980004b
- [23] X. Li, Y. Mao and I. Gutman, *The Steiner Wiener index of a graph*, *Discuss. Math. Graph Theory* **36** (2016) 455–465.
doi:10.7151/dmgt.1868
- [24] Y. Mao, Z. Wang and I. Gutman, *Steiner Wiener index of graph products*, *Trans. Comb.* **5** (2016) 39–50.
- [25] Y. Mao, Z. Wang, Y. Xiao and C. Ye, *Steiner Wiener index and connectivity of graphs*, *Util. Math.*, in press.
- [26] O.R. Oellermann and S. Tian, *Steiner centers in graphs*, *J. Graph Theory* **14** (1990) 585–597.
doi:10.1002/jgt.3190140510
- [27] D.H. Rouvray, *Harry in the limelight: The life and times of Harry Wiener*, in: D.H. Rouvray, R.B. King (Eds.), *Topology in Chemistry: Discrete Mathematics of Molecules* (Horwood, Chichester, 2002) 1–15.

- [28] D.H. Rouvray, *The rich legacy of half a century of the Wiener index*, in: D.H. Rouvray, R.B. King (Eds.), *Topology in Chemistry: Discrete Mathematics of Molecules* (Horwood, Chichester, 2002) 16–37.
- [29] S.G. Wagner, *A note on the inverse problem for the Wiener index*, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 639–646.
- [30] S.G. Wagner, *A class of trees and its Wiener index*, *Acta Appl. Math.* **91** (2006) 119–132.
doi:10.1007/s10440-006-9026-5
- [31] S.G. Wagner, H. Wang and G. Yu, *Molecular graphs and the inverse Wiener index problem*, *Discrete Appl. Math.* **157** (2009) 1544–1554.
doi:10.1016/j.dam.2008.06.008
- [32] H. Wang and G. Yu, *All but 49 numbers are Wiener indices of trees*, *Acta Appl. Math.* **92** (2006) 15–20.
doi:10.1007/s10440-006-9037-2
- [33] K. Xu, M. Liu, K.C. Das, I. Gutman and B. Furtula, *A survey on graphs extremal with respect to distance-based topological indices*, *MATCH Commun. Math. Comput. Chem.* **71** (2014) 461–508.

Received 16 September 2015

Revised 5 October 2016

Accepted 5 October 2016