2019 Volume 14 Issue 1

DOI: 10.20858/tp.2019.14.1.1

Keywords: drink; driving error; driving performance; food; intake

Iftikar Zahedi SUTALAKSANA, Gradiyan Budi PRATAMA*, Putra Alif Ramdhani YAMIN, Herman Rahadian SOETISNA

Department of Industrial Engineering, Faculty of Industrial Technology Institut Teknologi Bandung Jl. Ganesa 10, Bandung, Indonesia 40132 *Corresponding author. E-mail: gradiyan@gmail.com

THE EFFECTS OF FOOD AND DRINK INTAKE TO DRIVING PERFORMANCE: A SYSTEMATIC REVIEW

Summary. Human-related factors are considered to be the main cause of traffic incidents or accidents, causing 69.70% of the incidents. Several studies have been conducted to identify the relationship between drowsiness or fatigue and driving performance. Furthermore, a number of other studies not only discussed the symptoms causing drowsiness but also tried to investigate related factors that cause sleepiness or fatigue while driving. On the other hand, some discussed the quantity and quality of sleep as well as food and drink intake before and while driving. This systematic review, which is based on the PRISMA method, aims to map previous studies that investigated the effect of different food/drink consumption, either taken prior to driving or while driving, on the on-road driving characteristics of drivers. Furthermore, this article is expected to serve as a reference for further research that could potentially contribute to minimizing driving errors that lead to incident or accident. From 1871 articles screened, 7 studies related to food/drink intake and driving performance were reviewed. On the basis of the existing studies, no real evidence showing the presence of the association between food intake and the monotony of the road to decrease the driving performance has been found; therefore, further research is needed.

1. INTRODUCTION

The growth rate of traffic accidents in Indonesia is still high. During 2007-2016, in Indonesia, road accidents and traffic collisions were the most common type of accidents (65.6%). Among various types of traffic accidents, in 2014, a total number of 18,147 car accidents occurred. Although there is a decrease in the number of passenger car accidents by 8.8% per year, the number of accidents is still high, and thus resulting in large losses. There was an average increase of 35.8% casualties due to traffic accidents in the period 2007-2016 [1]. As per the data from the National Transportation Safety Committee, the main factors causing the accident were human factors (69.70%), compared to facilities factor (21.21%) and infrastructure (9.09%).

One of the major factors is driving errors due to drowsiness [2] and fatigue [3]. Several studies have been conducted to identify the relationship between drowsiness or fatigue and driving performance. In addition to fatigue and drowsiness, the driver's cognitive aspect, especially the level of driver's awareness of various road conditions, is also an interesting characteristic to consider. A number of other studies not only discussed the symptoms causing drowsiness but also tried to investigate related factors that cause sleepiness or fatigue while driving. On the other hand, some discussed the quantity and quality of sleep as well as food and drink intake before and while driving.

Some research studies have investigated the effect of food/drink consumption on the sleepiness and human driving performance. Orr et al. [4] studied the meal characteristics and their effect on postprandial sleepiness. Another research by Karl, et al. [5] examined how the composition of protein and carbohydrate within meal influences mood, cognition, and sleep. Jacobson et al. [6] discussed the relationship of a meal consumption style, which causes obesity, with the performance of driving. However, studies indicating the relationship between various types of food/drink intake and driving performance are still scarce. This study aims to review some of the literature that reported the results of previous studies on driving performance and food/beverage intake. This literature review is expected to serve as a reference for further research that could potentially contribute to minimizing driving errors.

2. METHODS

This literature review is based on the Prefered Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) approach [7]. PRISMA method was applied because it provides a clear procedure and milestone for the review of specific area of study. First, the author defined the research question, the scope of the study, and inclusion/exclusion criteria, complying with the PRISMA statement. Afterward, a systematic screening of available literature was performed to select a set of eligible studies. The review steps were conducted by two independent groups (I.Z.S-G.B.P. & P.A.R.Y-H.R.S) and required a consensus before moving to the next step. Ultimately, the selected articles were reviewed and summarized.

The literature search was done using a combination of key words: (driver OR driving) AND (simulator OR vehicle OR road) AND (food OR drink OR diet OR meal OR intake OR consumption) AND (performance OR sleepiness OR response OR reaction OR alertness OR vigilance OR attention). Some inclusion and exclusion criteria for paper selection were also considered, as listed in Table 1. Related literature was collected from search results using three main reference search engines including ScienceDirect, Transportation Research Information Documentation (TRID), and SCOPUS (Table 2).

Table 1 Inclusion and exclusion criteria for paper selection

Study Element	Inclusion Criteria	Exclusion Criteria
Article type	Research paper published in international journal	Book chapter, reports, conference proceedings were excluded
Subjects	Healthy, non-professional driver, free from sleep disorders	-
Exposure (Independent variable)	Food and drink consumption were the main factors	-
Outcome measures (Dependent variable)	Driving performance: driving error, speed, number of incident/accident Sleepiness Driver reaction	-
Publication date	Studies published between 2008 and 2018	-
Publication language	Articles published in English	Articles published in other language were excluded

Search characteristics from each database

Table 2

No	Database	Limiters	Identified	Selected
			articles	articles
1	ScienceDirect	Publication type: only journals,	481	8
		all sources, all sciences		
		Publication year: 2008-2018		
2	Transportation	Publication type: only article and	465	8
	Research	papers		
	Information	Language: English		
	Documentation	Publication year: 2008-2018		
	(TRID)			
3	Scopus	Limit document type: article,	925	10
		article in press		
		Limit language: English		
		Publication year: 2008-2018		

First of all, the title of the study became the basic parameter for filtering the literature in order to produce a fairly relevant list of papers. After the literature was filtered into smaller lists, a review of paper abstracts was conducted to collect the most focused studies on driving errors as well as food/drink intake. The final step was to review each paper to be analyzed so that the researcher can map the related studies that have been done so far in this area. The more detailed procedure of article search based on PRISMA flow is described in Fig. 1.




Fig. 1. Flow diagram of a systematic review (PRISMA)

3. RESULTS

Based on the procedure of the literature search for a systematic review, there were 7 studies reviewed in this paper. The specific title, study element, and the study findings are presented in Table 3.

4. DISCUSSION

Fatigue and sleepiness are a major contributing factor to transportation accidents [3], and food or drink intake is one of the potential causes of it. As regards to this fact, several studies have evaluated different possible types of intake that might be taken by the driver and their effect on driving behavior. According to the PRISMA-based systematic search in this review, there is only limited available evidence related to the effects of food and drink intake on driving performance. From the 7 studies reviewed in this paper, there is a heterogeneity of multiple study parameters and outcomes, which limits the generalizability of the findings. These study parameters are divided into driving-/driver-related measures, intake-related measures, and the road characteristics.

The road characteristics in the studies reviewed were mainly simulated long monotonous roads. Prolonged driving on monotonous roads is ideal for obtaining sleepiness and fatigue data on driving. Five out of the seven studies measured the sleepiness of the driver. A mixture of using objective and subjective measures has proven to be reliable in this area of research, especially when using Electroencephalograph (EEG) and Electrooculograph (EOG) [8, 9]. The subjective measures used included the well-proven Karolinska Sleepiness Scale [9, 10] or the Stanford Sleepiness Scale [11]. Other sleepiness parameters were eye blink, eye closure, eyes rolling upwards, and vacant staring.

Driving performance was measured in a number of different ways. Most studies measured the Standard Deviation of Lane Position (SDLP) and speed deviation [8, 10-14]. Other measures include braking reaction time [12] and the number of crashes or hits [8, 14]. It is arguable that driving measures that are more sensitive to change, such as lane and speed deviation, give better overall data to process, whereas measures like crashes, which are less likely to occur, are ineffective to measure driving performance related to food intake.

Intake-related measures are considered to be the most heterogeneous parameter and research related to them is scarce. Braag et al. [11] studied caffeine consumption and meal glycemic load. Ronen et al. [13] and Mets et al. [10] examined the effects of energy drinks. Watson et al. [8] limited the intake of drinking water, whereas Li et al (2016) studied the effects of alcohol. Only two studies evaluated the effects of food and drink intake on driving performance. The participants of the study conducted by Young et al. [14] ate simple snacks and consumed a bottle of water while driving. In contrast, Reyner et al. [9] suggested the consumption of light (305 calories) and heavy (922 calories) meal before driving in the driving simulator.

There are several studies that investigated the effects of alcohol and caffeine on driving performance, as reviewed in two separate articles [15-17]. However, currently, there is not much literature that discussed the influence of food or drink (besides alcohol and coffee) intake before/while driving on driving performance.

This literature review contributes to knowledge of the consequences that arise when driving on monotonous roads and its connection with food or drink consumed. This systematic review found, in general, no real evidence showing the presence of the association between food intake and the monotony of the road to affect driving performance. However, one study did find that heavy meals contribute to higher drowsiness levels and caffeinated drinks could increase driving performance and reduce the level of driver's sleepiness. There are several points that can be used as a reference for further research on driving behavior. Setting a straight and long path with a low-level stimulant proved to contribute to driver fatigue. There has been no real evidence showing the presence of the association of food intake and the monotony of the road to decrease the driving performance; hence, we recommend further study on this subject. Furthermore, there is still no study that compares different type of specific ingredients consumed by the driver.

1			Parameters		
Author (Year)/ Title	Participant	Driving-/Driver-related measures	Intake-related measures	Road Characteristics	Findings
Braag et al. (2017) Effect of meal glycemic load and caffeine consumption on prolonged monotonous driving performance	M: 7, F: 3 51±7 years old 97.4±19,9kg BMI 31.3±5.1 kg.m ⁻²	Vigilance: Visual Analogue Scale Sleepiness: Stanford Sleepiness Scale Driving error: Standard Deviation of Lane Position / SDLP, Average Lane Position / AVLP, Lane Crossing / LC Driving speed: Average Speed / AVSP, Standard Deviation of Speed / SDSP	• Caffeine (blood sample) • Glycemic load / GL (Blood Glucose Level)	Straight and long way (monotonic), 90 minutes driving (SCANeR Studio simulation engine v1.2r109)	• Significant decrease in SDLP (0.36 ± 0.20 m vs. 0.41 ± 0.19 m, p = 0.004) and LC (34.4 ± 31.4 vs. 56.7 ± 31.5, p = 0.018) on low GL conditions • There is no difference in the effect of glycemic consumption on driving performance
Li et al. (2016) A simulation study of the effects of alcohol on driving performance in a Chinese	M: 34, F: 18 21-61y.o. (avg 38.2 y.o.) Min. 1-year driver's license	Driving behavior (braking reaction time/BRT, standard deviation of lateral pos./SD-LANE, standard deviation of speed (SD-SPEED)	• Alcohol (Breath alcohol concentration/ BrAC	Free driving, urban area, 4 hours (XP-300 Driving Simulator)	□ Driving performance (BRT and SDLANE) → is highly correlated with the level of alcohol consumption □ 1 unit increase Brac → BRT decrease (0.3%) and SDLANE (0.2%)

walson of al.	M: 11	 Vigilance: Visual Analogue 	Hydration level	Long and straight road, 2	 Mild dehydration leads to a
(2015)	22±4 years old	Scale	(blood and urine	hours driving (immobile	significant increase in
	Height 1.75±0.06m	 Sleepiness: excessive 	sample)	car, with full size,	driving errors
Mild hypo-	Body mass	blinking, eye closure, eyes	 Hydrated trial 	interactive, computer-	 It is unclear if the difference
hydration	77.4±10.0kg	rolling upwards, vacant	(200 mL of	generated road projection)	was caused due to the fluid
increases the		staring ahead, EEG, EOG	fluid every	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	restriction, different fluid
frequency of		 Driving error: Lane drifting, 	hour)	1	intake during driving task,
driver errors		late breaking, car wheel	 Fluid restricted 		or combination of both
during a		touching lane line, car hits a	trial (25 mL		factors.
prolonged,		barrier or another car	every hour)		
monotonous		 Driving speed: Average 			
driving task		Speed/AVSP, Standard			
		Deviation of speed/SDSP	3		
Reyner et al.	M: 12	Sleepiness (Karolinska	Meal size	Straight and long	Heavy meal → Significant
(2012)	19-25 years old	Sleepiness Scale, EEG, EOG)	 Heavy meal 	(monotone), 120 minutes	increase in incident and EEG
	(avg: 21years old)	201	(922 cal.)	driving (immobile car,	Power
Post-lunch	BMI 20-27 kg.m ⁻²		• Light meal (305	interactive, computer	Heavy meal \rightarrow higher
sleepiness during	(avg: 23)	i gran	cal.)	generated road projection)	subjective drowsiness trends
prolonged,					
monotonous				ķz	
driving – Effects		8		(30)	
of meal size					

Ronen et al. (2014) The combination of short rest and	M: 15 24-29 years old Minimum 5 years of truck driving experience	 Driving performance: lane position, longitudinal speed, steering wheel deviations Subjective questionnaires: SOFI-20, NASA TLX 	• Energy drink • Energy drink + resting after 100 minutes of driving	Two-way monotonous and straight road, 40 minutes (STI-SIM fixed-based driving simulator, System Technology, Inc.)	Consumption of energy drink helped maintaining driving performance during the first part of drive
energy drink consumption as	•	 Physiological measures: ECG signals (R wave, R-R 	0		
fatigue		intervals, HRV)			
during a				600	
prolonged drive		C.			
of professional truck drivers			J		
Mets et al. (2010)		• Driving performance:	Energy drink	Monotonous and	An energy drink significantly
	volunteers	standard deviation of lateral	(caffeine, taurine,	prolonged road, 4 hours	improves driving performance
Positive effects of		position (SDLP), standard	glucuronolactone,	driving (STI-SIM driving	during prolonged driving
an energy drink		deviation of speed	vitamin B,	simulator)	
on driving		 Sleepiness: Karolinska 	inositol)		
performance		Sleepiness Scale (KSS)			
during prolonged					
Volung et al	M· 16· F· 10	• Driving performance: speed	Eating and	Urban environment	There is no significant
(2008)	37.5 ± 12.9 years old	lane position, number of	drinking while	representative: two-way	influence of eating and
		crashes, time-to-contact	driving (simple	road (1.75 km), mixed	drinking while driving on
Crash dieting:		(TTC)	snack: a packet of	curved and straight	driving performance
The effects of		 Mental workload (NASA 	sweets and a	sections (11 curves); 553	
eating and		Task Load Index)	bottle of water)	seconds (average) of	
drinking on				driving	
driving					
performance					

References

- 1. Saputra, A.D. Study of Traffic Accident Rate in Indonesia Base on KNKT (Komite Nasional Keselamatan Transportasi) Database from 2007-2016. *Warta Penelitian Perhubungan*. 2017. Vol. 29. P. 179-190.
- 2. Dinges, D.F. An overview of sleepiness and accidents. *Journal of Sleep Research*. 1995. Vol. 4. No. s2. P. 4-14.
- 3. Lal, S.K.L. & Craig, A. A critical review of the psychophysiology of driver fatigue. *Biological Psychology*. 2001. Vol. 55. No. 3. P. 173-194.
- 4. Orr, W.C. & Shadid, G. & Harnish, M.J. & Elsenbruch, S. Meal composition and its effect on postprandial sleepiness. *Physiology & behavior*. 1997. Vol. 62(4). P. 709-712.
- 5. Karl, J.P. & Thompson, L.A. & Niro, P.J. & Margolis, L.M. & McClung, J.P. & Cao, J.J. & Whigham, L.D. & Combs, G.F. & Young, A.J. & Lieberman, H.R. & Pasiakos, S.M. Transient decrements in mood during energy deficit are independent of dietary protein-to-carbohydrate ratio. *Physiology & behavior*. 2015. Vol. 139. P. 524-531.
- 6. Jacobson, S.H. & King, D.M. & Yuan, R. A note on the relationship between obesity and driving. *Transport Policy*. 2011 Sep 1. Vol. 18(5). P. 772-776.
- 7. Moher, D. & Liberati, A. & Tetzlaff, J. & Altman, D.G. Preferred reporting items for systematic reviews and metaanalyses: the PRISMA statement. *Annals of internal medicine*. 2009. Vol. 151(4). P. 264-269.
- 8. Watson, P. & Whale, A. & Mears, S.A. et al. Mild hypohydration increases the frequency of driver errors during a prolonged, monotonous driving task. *Physiology and Behavior*. 2015. Vol. 147. P. 313-318.
- 9. Reyner, L.A. & Wells, S.J. & Mortlock, V. & Horne, J.A. 'Post-lunch' sleepiness during prolonged, monotonous driving Effects of meal size. *Physiology and Behavior*. 2012. Vol. 105. No. 4. P. 1088-1091.
- 10. Mets, M.A.J. & Ketzer, S. & Blom, C. & Olivier, B. & Verster, J.C. Positive effects of an Energy Drink on driving performance during prolonged driving. *Psychopharmacology*. 2011. Vol. 214. No. 3. P. 737-745.
- 11. Bragg, C. & Desbrow, B. & Hall, S. & Irwin, C. Effect of meal glycemic load and caffeine consumption on prolonged monotonous driving performance. *Physiology and Behavior*. 2017. Vol. 181. P. 110-116.
- 12. Li, Y.C. & Sze, N.N. & Wong, S.C. & Yan, W. & Tsui, K.L. & So, F.L. A simulation study of the effects of alcohol on driving performance in a Chinese population. *Accident Analysis & Prevention*. 2016 Oct. 1. Vol. 95. P. 334-342.
- 13. Ronen, A. & Oron-Gilad, T. & Gershon, P. The combination of short rest and energy drink consumption as fatigue countermeasures during a prolonged drive of professional truck drivers. *Journal of Safety Research*. 2014. Vol. 49. P. 39-43.
- 14. Young, M.S. et al. Crash dieting: The effects of eating and drinking on driving performance. *Accident Analysis & Prevention.* 2008. Vol. 40. P. 142-148.
- 15. Heatherley, S.V. Caffeine withdrawal, sleepiness, and driving performance: What does the research really tell us? *Nutritional Neuroscience*. 2011. Vol. 14. No. 3. P. 89-95.
- 16. Irwin, C. et al. Effects of acute alcohol consumption on measures of simulated driving: A systematic review and meta-analysis. *Accident Analysis & Prevention*. 2017. Vol. 102. P. 248-266.
- 17. Soleimanloo, S.S. & Melanie J.W. & Garcia-Hansen, V. & Smith, S.S. The effects of sleep loss on young drivers' performance: A systematic review. *PLOS One.* 2017. Vol. 12(8). DOI: 10.1371/journal.pone.0184002