

ON THE PARTIAL FINITE ALTERNATING SUMS OF  
RECIPROCAL OF BALANCING AND  
LUCAS-BALANCING NUMBERS

UTKAL KESHARI DUTTA

AND

PRASANTA KUMAR RAY

*Sambalpur University, Jyoti Vihar, Burla, India*

**e-mail:** utkaldutta@gmail.com  
prasantamath@suniv.ac.in

**Abstract**

In this note, the finite alternating sums of reciprocals of balancing and Lucas-balancing numbers are considered and several identities involving these sums are deduced.

**Keywords:** balancing numbers, Lucas-balancing numbers, partial sums, reciprocal, floor function.

**2010 Mathematics Subject Classification:** 11B37, 11B39.

1. INTRODUCTION

A natural number  $n$  is said to be a balancing number if it is the solution of a simple Diophantine equation  $1+2+\cdots+(n-1) = (n+1)+(n+2)+\cdots+(n+l)$ , where  $l$  is a balancer corresponding to  $n$  [1]. Let  $\{B_n\}_{n \geq 0}$  be the balancing sequence and is recursively defined as  $B_0 = 0$ ,  $B_1 = 1$  and  $B_n = 6B_{n-1} - B_{n-2}$  for  $n \geq 2$ . For any balancing number  $B_n$ , the positive square roots of  $8B_n^2 + 1$  generate a sequence called as Lucas-balancing sequence  $\{C_n\}_{n \geq 0}$ . Lucas-balancing sequence satisfies the same recurrence as that of balancing sequence but with different initials, that is,  $C_n = 6C_{n-1} - C_{n-2}$  for  $n \geq 2$  with  $C_0 = 1$  and  $C_1 = 3$  [8].

Many researchers studied the partial infinite sums of reciprocal Fibonacci and other related numbers. Ohtsuka and Nakamura [7] studied the partial infinite

sums of reciprocal Fibonacci numbers and derived the following results, where  $[\cdot]$  denotes the floor function. For all  $n \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{\infty} \frac{1}{F_k} \right)^{-1} \right] = \begin{cases} F_{n-2}, & \text{if } n \text{ is even;} \\ F_{n-2} - 1, & \text{if } n \text{ is odd} \end{cases}$$

and for all  $n \geq 1$ ,

$$\left[ \left( \sum_{k=n}^{\infty} \frac{1}{F_k^2} \right)^{-1} \right] = \begin{cases} F_n F_{n-1} - 1, & \text{if } n \text{ is even;} \\ F_n F_{n-1}, & \text{if } n \text{ is odd.} \end{cases}$$

Later, Holiday and Komatsu [3] established several identities for generalized Fibonacci numbers  $G_n$  defined by  $G_n = aG_{n-1} + G_{n-2}$  for  $n \geq 2$  with  $(G_0, G_1) = (0, 1)$ . Recently, Wang and Wen [9] strengthened the above results to the finite case and deduced the following identities. For all  $m \geq 3$  and  $n \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_k} \right)^{-1} \right] = \begin{cases} F_{n-2}, & \text{if } n \text{ is even;} \\ F_{n-2} - 1, & \text{if } n \text{ is odd} \end{cases}$$

and for all  $m \geq 2$  and  $n \geq 1$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_k^2} \right)^{-1} \right] = \begin{cases} F_n F_{n-1} - 1, & \text{if } n \text{ is even;} \\ F_n F_{n-1}, & \text{if } n \text{ is odd.} \end{cases}$$

Several authors studied the bounds for partial infinite and finite reciprocal sums involving terms from Fibonacci sequence, Pell sequence (e.g., see [2, 4, 6, 10–13]). More recently, Komatsu and Panda [5] studied the partial infinite alternating sums of reciprocal of balancing numbers and derived some identities involving these sums. Among other results they have deduced the following result. For  $n \geq 1$ ,

$$\left[ \left( \sum_{k=n}^{\infty} \frac{(-1)^k}{B_k} \right)^{-1} \right] = \begin{cases} B_n + B_{n-1}, & \text{if } n \text{ is even;} \\ -(B_n + B_{n-1} + 1), & \text{if } n \text{ is odd.} \end{cases}$$

In the present study, we consider the partial finite alternating sums of reciprocals of balancing numbers, squared balancing numbers, even-indexed balancing numbers, odd-indexed balancing numbers, product of consecutive balancing numbers etc. We derive some identities relating to these sums that enhances the results of Komatsu and Panda [5].

2. AUXILIARY RESULTS

In this section, we discuss some well known results which are used to prove our main theorems.

The following results are found in [8].

**Lemma 1.** For every positive integer  $n \geq 1$ ,  $B_n^2 - B_{n-1}B_{n+1} = 1$ .

**Lemma 2.** For every positive integers  $m$  and  $n$ ,  $B_{m+n} = B_mB_{n+1} - B_{m-1}B_n$ .

Using the above results, we deduce the following lemmas.

**Lemma 3.** For any even positive integer  $n \geq 2$ ,  $f_1(n) + f_1(n + 1) + f_1(2n) + \frac{1}{B_{2n+1}+B_{2n}-1} > 0$ , where  $f_1(n) = \frac{1}{B_n+B_{n-1}-1} - \frac{(-1)^n}{B_n} - \frac{1}{B_{n+1}+B_n-1}$ .

*Proof.* Let

$$f_1(n) = \frac{1}{B_n + B_{n-1} - 1} - \frac{(-1)^n}{B_n} - \frac{1}{B_{n+1} + B_n - 1}.$$

For even  $n$ ,  $f_1(n)$  is negative. Now,

$$\begin{aligned} & f_1(n) + f_1(n + 1) + f_1(2n) + \frac{1}{B_{2n+1} + B_{2n} - 1} \\ &= \frac{1}{B_n + B_{n-1} - 1} - \frac{1}{B_n} + \frac{1}{B_{n+1}} - \frac{1}{B_{n+2} + B_{n+1} - 1} + \frac{1}{B_{2n} + B_{2n-1} - 1} - \frac{1}{B_{2n}} \\ &> \frac{1}{B_n + B_{n-1} - 1} - \frac{1}{B_n} + \frac{1}{B_{n+1}} - \frac{1}{B_{n+2} + B_{n+1} - 1} - \frac{1}{B_{2n}} \\ &= \frac{(B_{n+2} + B_{n+1}B_{n+2} + B_n) - (B_{n+1} + B_nB_{n-1} + B_{n-1})}{(B_n^2 + B_nB_{n-1} - B_n)(B_{n+1}^2 + B_{n+1}B_{n+2} - B_{n+1})} - \frac{1}{B_{2n}}. \end{aligned}$$

By virtue of Lemmas 1 and 2, it can be easily checked that

$$\begin{aligned} & B_{2n}((B_{n+2} + B_{n+1}B_{n+2} + B_n) - (B_{n+1} + B_nB_{n-1} + B_{n-1})) \\ &> (B_n^2 + B_nB_{n-1} - B_n)(B_{n+1}^2 + B_{n+1}B_{n+2} - B_{n+1}). \end{aligned}$$

Therefore,

$$f_1(n) + f_1(n + 1) + f_1(2n) + \frac{1}{B_{2n+1} + B_{2n} - 1} > 0.$$

This completes the proof. ■

**Lemma 4.** For any integer  $m \geq 2$  and odd positive integer  $n$ ,  $f_1(n) + f_1(n + 1) - \frac{1}{B_{mn+1}+B_{mn}+1} > 0$ , where  $f_1(n) = \frac{-1}{B_n+B_{n-1}+1} - \frac{(-1)^n}{B_n} + \frac{1}{B_{n+1}+B_n+1}$ .

**Proof.** Let

$$f_1(n) = \frac{-1}{B_n + B_{n-1} + 1} - \frac{(-1)^n}{B_n} + \frac{1}{B_{n+1} + B_n + 1}.$$

For odd  $n$ ,  $f_1(n)$  is positive and hence  $f_1(n) + f_1(n+1)$  is positive. In order to show the result, it suffices to prove that

$$f_1(n) + f_1(n+1) > \frac{1}{B_{2n+1} + B_{2n} + 1}.$$

Now,

$$\begin{aligned} & f_1(n) + f_1(n+1) - \frac{1}{B_{2n+1} + B_{2n} + 1} \\ &= \frac{-1}{B_n + B_{n-1} + 1} + \frac{1}{B_n} + \frac{1}{B_{n+2} + B_{n+1} + 1} - \frac{1}{B_{n+1}} - \frac{1}{B_{2n+1} + B_{2n} + 1} \\ &= \frac{B_{n+1}B_{n+2} + B_{n+1} + B_{n-1} - (B_{n+2} + B_n B_{n-1} + B_n)}{B_n B_{n+1} (B_n + B_{n-1} + 1) (B_{n+2} + B_{n+1} + 1)} - \frac{1}{B_{2n+1} + B_{2n} + 1}. \end{aligned}$$

Using Lemmas 1 and 2, the above identity simplifies  $f_1(n) + f_1(n+1) - \frac{1}{B_{2n+1} + B_{2n} + 1} > 0$ . This ends the proof.  $\blacksquare$

**Lemma 5.** For any integer  $m \geq 2$  and odd positive integer  $n$ ,  $f_2(n) + f_2(n+1) + f_2(mn) < 0$ , where  $f_2(n) = \frac{-1}{B_n + B_{n-1}} - \frac{(-1)^n}{B_n} + \frac{1}{B_{n+1} + B_n}$ .

**Proof.** Let

$$f_2(n) = \frac{-1}{B_n + B_{n-1}} - \frac{(-1)^n}{B_n} + \frac{1}{B_{n+1} + B_n}.$$

For odd  $n$ ,  $f_2(n)$  is positive and hence

$$f_2(n) + f_2(n+1) = \frac{B_{n-1} - B_{n+2}}{B_n B_{n+1} (B_n + B_{n-1}) (B_{n+2} + B_{n+1})},$$

which is negative. For  $m \geq 2$  and odd  $n$ , two cases arise. For  $mn$  is even,  $f_2(mn)$  negative. Thus, it is clear that  $f_2(n) + f_2(n+1) + f_2(mn) < 0$  for even  $mn$ . If  $mn$  is odd,  $m$  must be odd and greater than 2 and therefore

$$f_2(mn) = \frac{-1}{B_{mn} + B_{mn-1}} + \frac{1}{B_{mn}} + \frac{1}{B_{mn+1} + B_{mn}} < \frac{1}{B_{3n}}.$$

By virtue of Lemmas 1 and 2, it can be easily checked that

$$B_{3n}(B_{n+2} - B_{n-1}) > B_n B_{n+1} (B_n + B_{n-1}) (B_{n+2} + B_{n+1}).$$

Therefore,

$$\begin{aligned} & f_2(n) + f_2(n + 1) + f_2(mn) \\ & < \frac{B_{n-1} - B_{n+2}}{B_n B_{n+1} (B_n + B_{n-1})(B_{n+2} + B_{n+1})} + \frac{1}{B_{3n}} \\ & = \frac{B_{3n}(B_{n-1} - B_{n+2}) + B_n B_{n+1} (B_n + B_{n-1})(B_{n+2} + B_{n+1})}{B_n B_{n+1} B_{3n} (B_n + B_{n-1})(B_{n+2} + B_{n+1})} < 0. \end{aligned}$$

This finishes the proof. ■

### 3. MAIN RESULTS

Now, we are in a position to derive our main results.

**Theorem 6.** *If any integer  $n \geq 2$  is even, then  $\left[ \left( \sum_{k=n}^{2n} \frac{(-1)^k}{B_k} \right)^{-1} \right] = B_n + B_{n-1} - 1$ .*

*Proof.* For any positive integer  $k$ , consider

$$(3.1) \quad f_1(k) = \frac{1}{B_k + B_{k-1} - 1} - \frac{(-1)^k}{B_k} - \frac{1}{B_{k+1} + B_k - 1}.$$

For even  $k$ , it is clear that  $f_1(k)$  is negative and therefore

$$\begin{aligned} f_1(k) + f_1(k + 1) &= \left( \frac{1}{B_k + B_{k-1} - 1} - \frac{1}{B_k} \right) + \left( \frac{1}{B_{k+1}} - \frac{1}{B_{k+2} + B_{k+1} - 1} \right) \\ &= \frac{1 - B_{k-1}}{B_k(B_k - (1 - B_{k-1}))} - \frac{1 - B_{k+2}}{B_{k+1}(B_{k+1} - (1 - B_{k+2}))} \\ &= \frac{1}{B_k \left( \frac{B_k}{1 - B_{k-1}} - 1 \right)} - \frac{1}{B_{k+1} \left( \frac{B_{k+1}}{1 - B_{k+2}} - 1 \right)} \\ &= \frac{1}{\left( \frac{B_{k+1}B_{k-1} + 1}{1 - B_{k-1}} - B_k \right)} - \frac{1}{\left( \frac{B_{k+2}B_{k+1}}{1 - B_{k+2}} - B_{k+1} \right)} \\ &= \frac{-1}{B_{k+1} + B_k + \left( \frac{B_{k+1} + 1}{B_{k-1} - 1} \right)} + \frac{1}{B_{k+1} + B_k + \left( \frac{B_{k+1}}{B_{k+2} - 1} \right)} \\ &> 0. \end{aligned}$$

Taking summation over  $k$  from  $n$  to  $2n$  in (3.1), we get

$$\begin{aligned} \sum_{k=n}^{2n} \frac{(-1)^k}{B_k} &= \sum_{k=n}^{2n} \left( \frac{1}{B_k + B_{k-1} - 1} - \frac{1}{B_{k+1} + B_k - 1} \right) - \sum_{k=n}^{2n} f_1(k) \\ &= \frac{1}{B_n + B_{n-1} - 1} - \left[ \frac{1}{B_{2n+1} + B_{2n} - 1} + f_1(n) + f_1(n+1) + f_1(2n) \right] \\ &\quad - \sum_{k=n+2}^{2n-1} f_1(k). \end{aligned}$$

Since  $\sum_{k=n+2}^{2n-1} f_1(k) > 0$  and from Lemma 3, we have,

$$\sum_{k=n}^{2n} \frac{(-1)^k}{B_k} < \frac{1}{B_n + B_{n-1} - 1}.$$

On the other hand, consider  $f_2(k) = \frac{1}{B_k + B_{k-1}} - \frac{(-1)^k}{B_k} - \frac{1}{B_{k+1} + B_k}$ . For even  $k$ ,  $f_2(k)$  is negative. One can observe that  $f_2(k) + f_2(k+1) < 0$ . Hence

$$\begin{aligned} \sum_{k=n}^{2n} \frac{(-1)^k}{B_k} &= \frac{1}{B_n + B_{n-1}} - \frac{1}{B_{2n+1} + B_{2n}} - \sum_{k=n}^{2n} f_2(k) \\ &= \frac{1}{B_n + B_{n-1}} - \left[ \frac{1}{B_{2n+1} + B_{2n}} + f_2(2n) \right] - \sum_{k=n}^{2n-1} f_2(k) > \frac{1}{B_n + B_{n-1}}, \end{aligned}$$

the result follows. ■

**Theorem 7.** For any odd positive integer  $n$  and any integer  $m \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} \right)^{-1} \right] = -(B_n + B_{n-1} + 1).$$

*Proof.* In order to prove the theorem, it suffices to show that  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{-1}{B_n + B_{n-1} + 1}$  and  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} > \frac{-1}{B_n + B_{n-1}}$ . Consider

$$(3.2) \quad f_1(k) = \frac{-1}{B_k + B_{k-1} + 1} - \frac{(-1)^k}{B_k} + \frac{1}{B_{k+1} + B_k + 1},$$

and

$$(3.3) \quad f_2(k) = \frac{-1}{B_k + B_{k-1}} - \frac{(-1)^k}{B_k} + \frac{1}{B_{k+1} + B_k}.$$

For odd  $k$ , both  $f_1(k)$  and  $f_2(k)$  are positive. It is checked that  $f_1(k) + f_1(k+1)$  is positive for any odd positive integer  $k$ . Similarly, one can check that  $f_2(k) + f_2(k+1)$

1) is negative. Therefore, from the above results, we conclude  $\sum_{k=n}^{mn(\text{even})} f_1(k) > 0$  and  $\sum_{k=n}^{mn(\text{even})} f_2(k) < 0$ . Summing (3.2) over  $k$  from  $n$  to  $mn$ ,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \sum_{k=n}^{mn} \left( \frac{-1}{B_k + B_{k-1} + 1} + \frac{1}{B_{k+1} + B_k + 1} \right) - \sum_{k=n}^{mn} f_1(k) \\ &= \frac{-1}{B_n + B_{n-1} + 1} + \frac{1}{B_{mn+1} + B_{mn} + 1} - \sum_{k=n}^{mn} f_1(k). \end{aligned}$$

The following cases arise. When  $mn$  is odd,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \frac{-1}{B_n + B_{n-1} + 1} + \frac{1}{B_{mn+1} + B_{mn} + 1} - f_1(mn) - \sum_{k=n}^{mn-1} f_1(k) \\ &= \frac{-1}{B_n + B_{n-1} + 1} - \frac{B_{mn-1} + 1}{B_{mn}(B_{mn} + B_{mn-1} + 1)} - \sum_{k=n}^{mn-1} f_1(k). \end{aligned}$$

Since  $\sum_{k=n}^{mn-1} f_1(k) > 0$ , then  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{-1}{B_n + B_{n-1} + 1}$ . Now, for even  $mn$ ,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \frac{-1}{B_n + B_{n-1} + 1} + \frac{1}{B_{mn+1} + B_{mn} + 1} - f_1(n) - f_1(n+1) - \sum_{k=n+2}^{mn} f_1(k) \\ &= \frac{-1}{B_n + B_{n-1} + 1} - \sum_{k=n+2}^{mn} f_1(k) - \left[ f_1(n) + f_1(n+1) - \frac{1}{B_{mn+1} + B_{mn} + 1} \right]. \end{aligned}$$

Since  $\sum_{k=n+2}^{mn} f_1(k) > 0$  and using Lemma 4, we conclude

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{-1}{B_n + B_{n-1} + 1}.$$

On the other hand, taking summation over  $k$  from  $n$  to  $mn$  in (3.3), we obtain

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \sum_{k=n}^{mn} \left( \frac{-1}{B_k + B_{k-1}} + \frac{1}{B_{k+1} + B_k} \right) - \sum_{k=n}^{mn} f_2(k) \\ &= \frac{-1}{B_n + B_{n-1}} + \frac{1}{B_{mn+1} + B_{mn}} - \sum_{k=n}^{mn} f_2(k). \end{aligned}$$

If  $mn$  is even, then  $\sum_{k=n}^{mn(\text{even})} f_2(k) < 0$  and therefore

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \frac{-1}{B_n + B_{n-1}} + \frac{1}{B_{mn+1} + B_{mn}} - \sum_{k=n}^{mn} f_2(k) \\ &> \frac{-1}{B_n + B_{n-1}}. \end{aligned}$$

As  $\sum_{k=n+2}^{mn-1} f_2(k) < 0$  and from Lemma 5,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \frac{-1}{B_n + B_{n-1}} + \frac{1}{B_{mn+1} + B_{mn}} - \sum_{k=n}^{mn} f_2(k) \\ &= \frac{-1}{B_n + B_{n-1}} + \frac{1}{B_{mn+1} + B_{mn}} - [f_2(n) + f_2(n+1) + f_2(mn)] \\ &\quad - \sum_{k=n+2}^{mn-1} f_2(k) > \frac{-1}{B_n + B_{n-1}}. \end{aligned}$$

This completes the proof of the theorem. ■

**Theorem 8.** For any even positive integer  $n$  and any integer  $m \geq 3$ ,  
 $\left\lfloor \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} \right)^{-1} \right\rfloor = B_n + B_{n-1}$ .

**Proof.** In order to show the above the result, it is sufficient to prove that  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{1}{B_n + B_{n-1}}$  and  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} > \frac{1}{B_n + B_{n-1} + 1}$ . Consider

$$(3.4) \quad g_1(k) = \frac{1}{B_k + B_{k-1}} - \frac{(-1)^k}{B_k} - \frac{1}{B_{k+1} + B_k}.$$

For even  $k$ ,  $g_1(k) < 0$  and  $g_1(k) + g_1(k+1)$  is positive which can be easily checked. Taking summation over  $k$  from  $n$  to  $mn$  in (3.4), we have

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \sum_{k=n}^{mn} \left( \frac{1}{B_k + B_{k-1}} - \frac{1}{B_{k+1} + B_k} \right) - \sum_{k=n}^{mn} g_1(k) \\ &= \frac{1}{B_n + B_{n-1}} - \frac{1}{B_{mn+1} + B_{mn}} - \sum_{k=n}^{mn} g_1(k). \end{aligned}$$

If  $mn$  is odd, then  $\sum_{k=n}^{mn} g_1(k) > 0$  and therefore  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{1}{B_n + B_{n-1}}$ . For even  $mn$ ,  $\sum_{k=n}^{mn-1} g_1(k) > 0$  and hence



$$\begin{aligned} & \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} \\ &= \frac{1}{B_n + B_{n-1}} - \frac{1}{B_{mn+1} + B_{mn}} - \sum_{k=n+2}^{mn-1} g_1(k) - (g_1(n) + g_1(n+1) + g_1(mn)) \\ &= \frac{1}{B_n + B_{n-1}} - \sum_{k=n+2}^{mn-1} g_1(k) - \left( g_1(n) + g_1(n+1) + g_1(mn) + \frac{1}{B_{mn+1} + B_{mn}} \right). \end{aligned}$$

One can observe that  $g_1(n) + g_1(n+1) + g_1(mn) + \frac{1}{B_{mn+1} + B_{mn}} > 0$  and therefore,  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} < \frac{1}{B_n + B_{n-1}}$ . Let

$$(3.5) \quad g_2(k) = \frac{1}{B_k + B_{k-1} + 1} - \frac{(-1)^k}{B_k} - \frac{1}{B_{k+1} + B_k + 1}.$$

For even  $k$ ,  $g_2(k)$  and  $g_2(k) + g_2(k+1)$  are negative. Summing (3.5) over  $k$  from  $n$  to  $mn$ , we obtain

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k} &= \sum_{k=n}^{mn} \left( \frac{1}{B_k + B_{k-1} + 1} - \frac{1}{B_{k+1} + B_k + 1} \right) - \sum_{k=n}^{mn} g_2(k) \\ &= \frac{1}{B_n + B_{n-1} + 1} - \left( \frac{1}{B_{mn+1} + B_{mn} + 1} + g_2(mn) \right) - \sum_{k=n}^{mn-1} g_2(k) \\ &= \frac{1}{B_n + B_{n-1} + 1} - \left( \frac{1}{B_{mn} + B_{mn-1} + 1} - \frac{1}{B_{mn}} \right) - \sum_{k=n}^{mn-1} g_2(k). \end{aligned}$$

Since  $\sum_{k=n}^{mn-1} g_2(k) < 0$ , it follows that  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k} > \frac{1}{B_n + B_{n-1} + 1}$ . This ends the proof of the theorem. ■

**Theorem 9.** For any even positive integer  $n$  and any integer  $m \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} \right)^{-1} \right] = B_n^2 + B_{n-1}^2.$$

*Proof.* Consider  $g_1(k) = \frac{1}{B_k^2 + B_{k-1}^2} - \frac{(-1)^k}{B_k^2} - \frac{1}{B_{k+1}^2 + B_k^2}$ . For even  $k$ ,  $g_1(k) < 0$  and it can be observed that  $g_1(k) + g_1(k+1) > 0$ . With the help of (3.4),

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \sum_{k=n}^{mn} \left( \frac{1}{B_k^2 + B_{k-1}^2} - \frac{1}{B_{k+1}^2 + B_k^2} \right) - \sum_{k=n}^{mn} g_1(k) = \frac{1}{B_n^2 + B_{n-1}^2} \\ &\quad - \sum_{k=n+2}^{mn-1} g_1(k) - \left[ g_1(n) + g_1(n+1) + g_1(mn) + \frac{1}{B_{mn+1}^2 + B_{mn}^2} \right]. \end{aligned}$$

It is observed that  $g_1(n) + g_1(n+1) + g_1(mn) + \frac{1}{B_{mn+1}^2 + B_{mn}^2} > 0$  and  $\sum_{k=n+2}^{mn-1} g_1(k) > 0$ . Therefore,

$$(3.6) \quad \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} < \frac{1}{B_n^2 + B_{n-1}^2}.$$

On the other hand, consider  $g_2(k) = \frac{1}{B_k^2 + B_{k-1}^2 + 1} - \frac{(-1)^k}{B_k^2} - \frac{1}{B_{k+1}^2 + B_k^2 + 1}$ . For even  $k$ , both  $g_2(k)$  and  $g_2(k) + g_2(k+1)$  are negative. Therefore,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \sum_{k=n}^{mn} \left( \frac{1}{B_k^2 + B_{k-1}^2 + 1} - \frac{1}{B_{k+1}^2 + B_k^2 + 1} \right) - \sum_{k=n}^{mn} g_2(k) \\ &= \frac{1}{B_n^2 + B_{n-1}^2 + 1} - \sum_{k=n}^{mn-1} g_2(k) - \left( g_2(mn) + \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} \right). \end{aligned}$$

As  $\sum_{k=n}^{mn-1} g_2(k) < 0$  and  $g_2(mn) + \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} < 0$ ,

$$(3.7) \quad \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} > \frac{1}{B_n^2 + B_{n-1}^2 + 1}.$$

The result follows from (3.6) and (3.7). ■

**Theorem 10.** For any positive odd integer  $n$  and any integer  $m \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} \right)^{-1} \right] = -(B_n^2 + B_{n-1}^2 + 1).$$

*Proof.* In order to prove the result, it is sufficient to show that  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} < \frac{-1}{B_n^2 + B_{n-1}^2 + 1}$  and  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} > \frac{-1}{B_n^2 + B_{n-1}^2}$ . Consider

$$(3.8) \quad s_1(k) = \frac{-1}{B_k^2 + B_{k-1}^2 + 1} - \frac{(-1)^k}{B_k^2} + \frac{1}{B_{k+1}^2 + B_k^2 + 1},$$

and

$$(3.9) \quad s_2(k) = \frac{-1}{B_k^2 + B_{k-1}^2} - \frac{(-1)^k}{B_k^2} + \frac{1}{B_{k+1}^2 + B_k^2}.$$

For any odd positive integer  $k$ ,  $s_1(k)$  and  $s_2(k)$  are positive. It can be easily checked that  $s_1(k) + s_1(k+1) > 0$  and  $s_2(k) + s_2(k+1) < 0$  for any odd positive integer  $k$ . Therefore,

$$\sum_{k=n}^{mn(\text{even})} s_1(k) > 0 \quad \text{and} \quad \sum_{k=n}^{mn(\text{even})} s_2(k) < 0.$$

Taking summation over  $k$  from  $n$  to  $mn$  in (3.8),

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \sum_{k=n}^{mn} \left( \frac{-1}{B_k^2 + B_{k-1}^2 + 1} + \frac{1}{B_{k+1}^2 + B_k^2 + 1} \right) - \sum_{k=n}^{mn} s_1(k) \\ &= \frac{-1}{B_n^2 + B_{n-1}^2 + 1} + \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} - \sum_{k=n}^{mn} s_1(k). \end{aligned}$$

For odd  $mn$ , we write

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \frac{-1}{B_n^2 + B_{n-1}^2 + 1} + \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} - s_1(mn) - \sum_{k=n}^{mn-1} s_1(k) \\ &= \frac{-1}{B_n^2 + B_{n-1}^2 + 1} - \frac{B_{mn-1}^2 + 1}{B_{mn}^2(B_{mn}^2 + B_{mn-1}^2 + 1)} - \sum_{k=n}^{mn-1} s_1(k). \end{aligned}$$

Since  $\sum_{k=n}^{mn-1} s_1(k) > 0$ , from the above identity, it follows that  $\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} < \frac{-1}{B_n^2 + B_{n-1}^2 + 1}$ . When  $mn$  is even, we can write

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \frac{-1}{B_n^2 + B_{n-1}^2 + 1} + \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} - s_1(n) - s_1(n+1) - \sum_{k=n+2}^{mn} s_1(k) \\ &= \frac{-1}{B_n^2 + B_{n-1}^2 + 1} - \sum_{k=n+2}^{mn} s_1(k) \\ &\quad - \left[ s_1(n) + s_1(n+1) - \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} \right]. \end{aligned}$$

It can be easily checked that  $s_1(n) + s_1(n+1) - \frac{1}{B_{mn+1}^2 + B_{mn}^2 + 1} > 0$  and  $\sum_{k=n+2}^{mn} s_1(k) > 0$ . Therefore,

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} < \frac{-1}{B_n^2 + B_{n-1}^2 + 1},$$

which completes the first part of the theorem. On the other hand, taking summation over  $k$  from  $n$  to  $mn$  in (3.9), we get

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \sum_{k=n}^{mn} \left( \frac{-1}{B_k^2 + B_{k-1}^2} + \frac{1}{B_{k+1}^2 + B_k^2} \right) - \sum_{k=n}^{mn} s_2(k) \\ &= \frac{-1}{B_n^2 + B_{n-1}^2} + \frac{1}{B_{mn+1}^2 + B_{mn}^2} - \sum_{k=n}^{mn} s_2(k). \end{aligned}$$

Since  $\sum_{k=n}^{mn} s_2(k) < 0$  for even  $mn$  and therefore

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} = \frac{-1}{B_n^2 + B_{n-1}^2} + \frac{1}{B_{mn+1}^2 + B_{mn}^2} - \sum_{k=n}^{mn} s_2(k) > \frac{-1}{B_n^2 + B_{n-1}^2}.$$

For odd  $mn$ , we proceed as follows.

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} &= \frac{-1}{B_n^2 + B_{n-1}^2} + \frac{1}{B_{mn+1}^2 + B_{mn}^2} - \sum_{k=n}^{mn} s_2(k) = \frac{-1}{B_n^2 + B_{n-1}^2} \\ &+ \frac{1}{B_{mn+1}^2 + B_{mn}^2} - \left[ s_2(n) + s_2(n+1) + s_2(mn) \right] - \sum_{k=n+2}^{mn-1} s_2(k). \end{aligned}$$

It can be easily checked that  $s_2(n) + s_2(n+1) + s_2(mn) < 0$  and  $\sum_{k=n+2}^{mn-1} s_2(k) < 0$ . Therefore,

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k^2} > \frac{-1}{B_n^2 + B_{n-1}^2}.$$

This finishes the proof. ■

The following results deal with the finite alternating sums of reciprocals of even and odd-indexed balancing numbers. The proofs are analogous to Theorems 7 and 8.

**Theorem 11.** *For any positive integer  $m \geq 2$  and any even integer  $n \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k}} \right)^{-1} \right] = B_{2n} + B_{2n-2}.$$

**Theorem 12.** *For any odd positive integer  $n$  and any integer  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k}} \right)^{-1} \right] = -(B_{2n} + B_{2n-2} + 1).$$

**Theorem 13.** *For any even positive integer  $n$  and any integer  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k+1}} \right)^{-1} \right] = \begin{cases} B_{2n+1} + B_{2n-1} - 1, & \text{if } m = 2; \\ B_{2n+1} + B_{2n-1}, & \text{if } m \geq 3. \end{cases}$$

**Theorem 14.** *For any odd positive integer  $n$  and any integer  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k+1}} \right)^{-1} \right] = -(B_{2n+1} + B_{2n-1} + 1).$$

The following result concerns with the finite alternating sums of reciprocals of product of two consecutive balancing numbers.

**Theorem 15.** *For any positive integers  $n \geq 1$  and  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_k B_{k+1}} \right)^{-1} \right] = \begin{cases} B_{n-1}B_n + B_nB_{n+1}, & \text{if } n \text{ is even;} \\ -(B_{n-1}B_n + B_nB_{n+1} + 1), & \text{if } n \text{ is odd.} \end{cases}$$

*Proof.* Consider

$$(3.10) \quad S_1(k) = \frac{1}{B_{k-1}B_k + B_kB_{k+1}} - \frac{(-1)^k}{B_kB_{k+1}} - \frac{1}{B_kB_{k+1} + B_{k+1}B_{k+2}}$$

and

$$(3.11) \quad S_2(k) = \frac{1}{B_{k-1}B_k + B_kB_{k+1} + 1} - \frac{(-1)^k}{B_kB_{k+1}} - \frac{1}{B_kB_{k+1} + B_{k+1}B_{k+2} + 1}.$$

For even  $k$ , both  $S_1(k)$  and  $S_2(k)$  are negative. Now,

$$\begin{aligned} & S_1(k) + S_1(k+1) \\ &= \frac{1}{B_{k-1}B_k + B_kB_{k+1}} - \frac{1}{B_kB_{k+1}} + \frac{1}{B_{k+1}B_{k+2}} - \frac{1}{B_{k+1}B_{k+2} + B_{k+2}B_{k+3}} \\ &= \frac{1}{B_{k+1}B_{k+2} \left(1 + \frac{B_{k+1}}{B_{k+3}}\right)} - \frac{1}{B_kB_{k+1} \left(1 + \frac{B_{k+1}}{B_{k-1}}\right)} \\ &= \frac{1}{B_{k+1}B_{k+2} + (1 + B_kB_{k+2})\frac{B_{k+2}}{B_{k+3}}} - \frac{1}{B_kB_{k+1} + (1 + B_kB_{k+2})\frac{B_k}{B_{k-1}}} \\ &= \frac{1}{B_{k+1}B_{k+2} + B_kB_{k+1} + \frac{B_k+B_{k+2}}{B_{k+3}}} - \frac{1}{B_{k+1}B_{k+2} + B_kB_{k+1} + \frac{B_kB_{k+2}}{B_{k-1}}} > 0, \end{aligned}$$

as

$$\frac{B_k + B_{k+2}}{B_{k+3}} < \frac{B_kB_{k+2}}{B_{k-1}}.$$

In a similar manner, we can check that  $S_2(k) + S_2(k+1) < 0$  for any even integer  $k \geq 2$ . Taking summation over  $k$  from  $n$  to  $mn$  in (3.10), we have

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{B_k B_{k+1}} &= \sum_{k=n}^{mn} \left[ \frac{1}{B_{k-1}B_k + B_kB_{k+1}} - \frac{1}{B_kB_{k+1} + B_{k+1}B_{k+2}} \right] - \sum_{k=n}^{mn} S_1(k) \\ &= \frac{1}{B_{n-1}B_n + B_nB_{n+1}} - \frac{1}{B_{mn}B_{mn+1} + B_{mn+1}B_{mn+2}} \\ &\quad - \left[ S_1(n) + S_1(n+1) + S_1(mn) \right] - \sum_{k=n+2}^{mn-1} S_1(k). \end{aligned}$$

It can be easily checked that  $S_1(n)+S_1(n+1)+S_1(mn) > 0$  and  $\sum_{k=n+2}^{mn-1} S_1(k) > 0$ . Therefore,

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k B_{k+1}} < \frac{1}{B_{n-1} B_n + B_n B_{n+1}}.$$

Similarly, with the help of (3.11), we can prove that

$$\sum_{k=n}^{mn} \frac{(-1)^k}{B_k B_{k+1}} > \frac{1}{B_{n-1} B_n + B_n B_{n+1} + 1},$$

which completes the theorem for even  $n$ . Considering

$$S_3(k) = \frac{-1}{B_{k-1} B_k + B_k B_{k+1} + 1} - \frac{(-1)^k}{B_k B_{k+1}} + \frac{1}{B_k B_{k+1} + B_{k+1} B_{k+2} + 1}$$

and

$$S_4(k) = \frac{-1}{B_{k-1} B_k + B_k B_{k+1}} - \frac{(-1)^k}{B_k B_{k+1}} + \frac{1}{B_k B_{k+1} + B_{k+1} B_{k+2}},$$

we can prove that

$$\frac{-1}{B_{n-1} B_n + B_n B_{n+1}} < \sum_{k=n}^{mn} \frac{(-1)^k}{B_k B_{k+1}} < \frac{-1}{B_{n-1} B_n + B_n B_{n+1} + 1}.$$

This completes the proof of the theorem. ■

Similarly, the following results can be proved.

**Theorem 16.** For any positive integers  $n \geq 1$  and  $m \geq 2$ ,

- (i)  $\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k}^2} \right)^{-1} \right] = \begin{cases} B_{2n}^2 + B_{2n-2}^2, & \text{if } n \text{ is even;} \\ -(B_{2n}^2 + B_{2n-2}^2 + 1), & \text{if } n \text{ is odd.} \end{cases}$
- (ii)  $\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k-1}^2} \right)^{-1} \right] = \begin{cases} B_{2n-1}^2 + B_{2n-3}^2, & \text{if } n \text{ is even;} \\ -(B_{2n-1}^2 + B_{2n-3}^2 + 1), & \text{if } n \text{ is odd.} \end{cases}$
- (iii)  $\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k-1} B_{2k+1}} \right)^{-1} \right] = \begin{cases} B_{2n}^2 + B_{2n-2}^2 - 1, & \text{if } n \text{ is even;} \\ -(B_{2n}^2 + B_{2n-2}^2), & \text{if } n \text{ is odd.} \end{cases}$
- (iv)  $\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{B_{2k} B_{2k+2}} \right)^{-1} \right] = \begin{cases} B_{2n+1}^2 + B_{2n-1}^2 - 1, & \text{if } n \text{ is even;} \\ -(B_{2n+1}^2 + B_{2n-1}^2), & \text{if } n \text{ is odd.} \end{cases}$

The following are the corresponding results for Lucas-balancing numbers  $C_n$  which can be analogously shown.

**Theorem 17.** For any positive integers  $n \geq 1$  and  $m \geq 2$ ,

$$\begin{aligned}
 \text{(i)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_k} \right)^{-1} \right] = \begin{cases} C_n + C_{n-1} - 1, & \text{if } n \text{ is even;} \\ -(C_n + C_{n-1}), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(ii)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k}} \right)^{-1} \right] = \begin{cases} C_{2n} + C_{2n-2} - 1, & \text{if } n \text{ is even;} \\ -(C_{2n} + C_{2n-2}), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(iii)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k+1}} \right)^{-1} \right] = \begin{cases} C_{2n+1} + C_{2n-1} - 1, & \text{if } n \text{ is even;} \\ -(C_{2n+1} + C_{2n-1}), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(iv)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_k^2} \right)^{-1} \right] = \begin{cases} C_n^2 + C_{n-1}^2 - 1, & \text{if } n \text{ is even;} \\ -(C_n^2 + C_{n-1}^2), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(v)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_k C_{k+1}} \right)^{-1} \right] = \begin{cases} C_{n-1} C_n + C_n C_{n+1} - 1, & \text{if } n \text{ is even;} \\ -(C_{n-1} C_n + C_n C_{n+1}), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(vi)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k}^2} \right)^{-1} \right] = \begin{cases} C_{2n}^2 + C_{2n-2}^2 - 1, & \text{if } n \text{ is even;} \\ -(C_{2n}^2 + C_{2n-2}^2), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(vii)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k-1}^2} \right)^{-1} \right] = \begin{cases} C_{2n-1}^2 + C_{2n-3}^2 - 1, & \text{if } n \text{ is even;} \\ -(C_{2n-1}^2 + C_{2n-3}^2), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(viii)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k-1} C_{2k+1}} \right)^{-1} \right] = \begin{cases} C_{2n}^2 + C_{2n-2}^2 - 1, & \text{if } n \text{ is even;} \\ -(C_{2n}^2 + C_{2n-2}^2), & \text{if } n \text{ is odd.} \end{cases} \\
 \text{(ix)} \quad & \left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{C_{2k} C_{2k+2}} \right)^{-1} \right] = \begin{cases} C_{2n+1}^2 + C_{2n-1}^2 - 1, & \text{if } n \text{ is even;} \\ -(C_{2n+1}^2 + C_{2n-1}^2), & \text{if } n \text{ is odd.} \end{cases}
 \end{aligned}$$

REFERENCES

[1] A. Behera and G. K. Panda, *On the square roots of triangular numbers*, Fibonacci Quart. **37** (1999) 98–105.

[2] Y. Choo, *On the reciprocal of sums of products of Pell numbers*, Int. J. Math. Anal. **12** (2018) 595–602.  
doi:10.12988/ijma.2018.81074

[3] S.H. Holliday and T. Komatsu, *On the sum of reciprocal generalized Fibonacci numbers*, Integers **11** (2011) 441–455.  
doi:10.1515/integ.2011.031

[4] E. Kilic and T. Arıkan, *More on the infinite sums of reciprocal Fibonacci, Pell and higher order recurrences*, Appl. Math. Comput. **219** (2013) 7783–7788.  
doi:10.1016/j.amc.2013.02.003

[5] T. Komatsu and G.K. Panda, *On several kinds of sums of balancing numbers*, Ars Combin. (to appear), arXiv:1608.05918.

- [6] R. Liu and A.Y. Wang, *Sums of products of two reciprocal Fibonacci numbers*, Adv. Differ. Equ. **2016** (2016) Article ID 136.  
doi:10.1186/s13662-016-0860-0
- [7] H. Ohtsuka and S. Nakamura, *On the sum of reciprocal Fibonacci numbers*, Fibonacci Quart. **46/47** (2008/2009) 153–159.
- [8] G.K. Panda, *Some fascinating properties of balancing numbers*, Congr. Numer. **194** (2009) 185–189.
- [9] A.Y. Wang and P. Wen, *On the partial finite sums of the reciprocals of the Fibonacci numbers*, J. Inequal. Appl. **2015** (2015) Article ID 73.  
doi:10.1186/s13660-015-0595-6
- [10] A.Y. Wang and T. Yuan, *Alternating sums of the reciprocal Fibonacci numbers*, J. Integer Seq. **20** (2017) Article ID 17.1.4.
- [11] A.Y. Wang and W. Zhang, *The reciprocal sums of even and odd terms in the Fibonacci sequence*, J. Inequal. Appl. **2015** (2015) Article ID 376.  
doi:10.1186/s13660-015-0902-2
- [12] A.Y. Wang and F. Zhang, *The reciprocal sums of the Fibonacci 3-subsequences*, Adv. Differ. Equ. **2016** (2016) Article ID 27.  
doi:10.1186/s13662-016-0761-2
- [13] W. Zhang and T. Wang, *The infinite sum of reciprocal Pell numbers*, Appl. Math. Comput. **218** (2012) 6164–6167.  
doi:10.1016/j.amc.2011.11.090

Received 8 November 2019

Revised 21 January 2020

Accepted 15 June 2020