

## ON eGE-ALGEBRAS

R.K. BANDARU<sup>1</sup>

*Department of Mathematics*  
*GITAM (Deemed to be University)*  
*Hyderabad Campus, India-502 329*

**e-mail:** ravimaths83@gmail.com

N. RAFI

*Department of Mathematics*  
*Bapatla Engineering College*  
*Bapatla, Andhra Pradesh, India-522 101*

**e-mail:** rafimaths@gmail.com

AND

A. REZAEI

*Department of Mathematics*  
*Payame Noor University*  
*P.O. Box 19395-3697, Tehran, Iran*

**e-mail:** rezaei@pnu.ac.ir

### Abstract

A new algebraic structure was introduced, called an eGE-algebra, which is a generalisation of a GE-algebra and investigated its properties. We explore the definition of filters and the quotient algebra associated with such filters.

**Keywords:** BE-algebra, GE-algebra, eGE-algebra, transitive, filter.

**2010 Mathematics Subject Classification:** 06D20, 06F35, 03G25.

---

<sup>1</sup>Corresponding author.

## 1. INTRODUCTION AND PRELIMINARIES

L. Henkin and T. Skolem developed the idea of Hilbert algebra in the early 50-ties for some investigations of implication in intuitionistic and other nonclassical logics. In 60-ties, these algebras were particularly studied by Horn and Diego [6] from algebraic point of view. Hilbert algebras are a valuable tool for some algebraic logic investigations as they can be regarded as fragments of any propositional logic that contains a logical connective implication ( $\rightarrow$ ) and the constant 1 that is assumed to be the logical meaning “true”. Many researchers have done a significant amount of work on Hilbert algebras [3–5, 7–9, 13–16]. As a generalization of Hilbert algebras, Bandaru *et al.* [1] introduce the notion of GE-algebras. They studied the various properties and filter theory of GE-algebras. BCK-algebras and BCI-algebras were introduced by Imai and Iseki [10, 11]. H.S. Kim and Y.H. Kim [12] developed the concept of BE-algebra as a generalization of dual BCK-algebra. Many researchers developed theory of BE-algebras [2, 17–19]. Rezaei [20] has introduced the notion of eBE-algebra as a generalization of BE-algebra and has studied some of its properties. It is important to make clear the corresponding algebraic structures for the creation of many-valued logical system. As a generalization of GE-algebra, we are inspired to concentrate on a new algebraic structure, called eGE-algebra, and thus to investigate some properties.

**Definition 1.1** [1]. Let  $X$  be a non-empty set with a constant 1 and  $*$  a binary operation on  $X$ . Then an algebraic structure  $(X, *, 1)$  of type  $(2, 0)$  is said to be a GE-algebra if it satisfies the following axioms:

$$(GE1) \quad u * u = 1,$$

$$(GE2) \quad 1 * u = u,$$

$$(GE3) \quad u * (v * w) = u * (v * (u * w))$$

for all  $u, v, w \in X$ .

In a GE-algebra  $X$ , a binary relation “ $\leq$ ” is defined by

$$(1) \quad (\forall u, v \in X) (u \leq v \Leftrightarrow u * v = 1).$$

**Proposition 1.2** [1]. Every GE-algebra  $X$  satisfies the following items.

$$(2) \quad (\forall u \in X) (u * 1 = 1).$$

$$(3) \quad (\forall u, v \in X) (u * (u * v) = u * v).$$

$$(4) \quad (\forall u, v \in X) (u \leq v * u).$$

**Definition 1.3** [1]. A GE-algebra  $X$  is said to be transitive, if it satisfies:

$$(5) \quad (\forall x, y, z \in X) (x * y \leq (z * x) * (z * y)).$$

**Proposition 1.4** [1]. Every transitive GE-algebra  $X$  satisfies the following assertions.

- (6)  $(\forall x, y, z \in X) (x * y \leq (y * z) * (x * z)).$   
 (7)  $(\forall x, y, z \in X) (x \leq y \Rightarrow z * x \leq z * y, y * z \leq x * z).$

**Definition 1.5** [1]. A subset  $F$  of a GE-algebra  $X$  is called a *filter* of  $X$  if it satisfies:

- (8)  $1 \in F,$   
 (9)  $(\forall x, y \in X)(x * y \in F, x \in F \Rightarrow y \in F).$

**Lemma 1.6** [1]. In a GE-algebra  $X$ , every filter  $F$  of  $X$  satisfies:

- (10)  $(\forall x, y \in X) (x \leq y, x \in F \Rightarrow y \in F).$

**Definition 1.7** [20]. Let  $X$  be a non-empty set. By an eBE-algebra we shall mean an algebra  $(X, *, A)$  such that “ $*$ ” is a binary operation on  $X$  and  $A$  is a non-empty subset of  $X$  satisfying the following axioms:

- (eBE1)  $x * x \in A,$   
 (eBE2)  $x * A \subseteq A,$   
 (eBE3)  $A * x = \{x\},$   
 (eBE4)  $x * (y * z) = y * (x * z)$

for all  $x, y, z \in X$ .

**Definition 1.8** [20]. An eBE-algebra  $X$  is said to be self distributive if it satisfies:

- (11)  $(\forall x, y, z \in X) x * (y * z) = (x * y) * (x * z).$

## 2. ON eGE-ALGEBRAS

In this section, we present the notion of eGE-algebra as a generalization of GE-algebra and study its properties.

**Definition 2.1.** An algebraic structure  $(X, *, E)$ , where  $*$  is a binary operation on a non-empty set  $X$  and  $E$  is a non-empty subset of  $X$ , is said to be an extended GE-algebra (eGE-algebra for short) if it satisfies the following axioms:

- (eGE1)  $u * u \in E,$   
 (eGE2)  $u * E \subseteq E,$   
 (eGE3)  $E * u = \{u\},$   
 (eGE4)  $u * (v * w) = u * (v * (u * w))$

for all  $u, v, w \in X$ .

Throughout the paper,  $E * u = \{e * u \mid e \in E\}$  and  $u * E = \{u * e \mid e \in E\}$ . If  $a, b \in E$  then, by (eGE3), we have  $a * b = b \in E$  and  $b * a = a \in E$ . Hence  $E$  is a closed subset of  $X$ .

We introduce a relation  $\leq$  on  $X$  by  $u \leq v$  if and only if  $u * v \in E$ . By (eGE1) the relation  $\leq$  is reflexive.

**Theorem 2.2.** *Every GE-algebra is an eGE-algebra.*

*Proof.* Put  $E = \{1\}$ . Then  $(X, *, E)$  is an eGE-algebra. ■

Every eGE-algebra need not be a GE-algebra which is shown in the following example.

**Example 2.3.** Let  $X = \{a, b, c, d, e\}$  be a set and  $*$  a binary operation given in the table:

*	a	b	c	d	e
a	c	b	c	c	b
b	d	d	d	d	d
c	a	b	c	d	e
d	a	b	c	d	e
e	a	c	c	d	c

Then  $(X, *, E)$ , where  $E = \{c, d\}$ , is an eGE-algebra, but not a GE-algebra. Since  $b * b = d$  and  $c * c = c$  and there is no  $1 \in X$ , such that  $u * u = 1$ , for all  $u \in X$ .

Note that the relation  $\leq$  need not be transitive in an eGE-algebra. From Example 2.3, we can observe that  $e * b = c \in E, b * a = d \in E$ , but  $e * a = a \notin E$ .

In the following example, we show that the axioms (eGE1) to (eGE4) are independent.

**Example 2.4.** (i) Let  $X = \{a, b, c, d\}$  be a set and  $*$  a binary operation on  $X$  given in the following table:

*	a	b	c	d
a	b	b	d	d
b	a	b	c	c
c	a	b	b	b
d	a	b	b	b

Then  $(X, *, E)$ , where  $E = \{a, b\}$ , satisfies (eGE2), (eGE3) and (eGE4), but it does not satisfy (eGE1), since  $E * u \neq \{u\}$  i.e.,  $a * c \neq c$  and  $a \in E$ .

(ii) Let  $X = \{a, b, c, d\}$  be a set and  $*$  a binary operation on  $X$  in the following table:

*	a	b	c	d
a	b	b	c	c
b	a	b	c	d
c	a	b	c	d
d	a	a	c	c

Then  $(X, *, E)$ , where  $E = \{b, c\}$ , satisfies (eGE1), (eGE3) and (eGE4), but it does not satisfy (eGE2), since  $u * E \not\subseteq E$ , i.e.,  $d * b = a \notin E$  and  $b \in E$ .

(iii) Let  $X = \{a, b, c, d\}$  be a set and  $*$  a binary operation on  $X$  given in the following table:

*	a	b	c	d
a	b	b	b	b
b	a	b	c	d
c	a	b	c	d
d	a	c	c	a

Then  $(X, *, E)$ , where  $E = \{b, c\}$  satisfies (eGE1), (eGE2) and (eGE4), but it does not satisfy (eGE3), since  $d * d = a \notin E$ .

(iv) Let  $X = \{a, b, c, d\}$  be a set and  $*$  a binary operation on  $X$  given in the following table:

*	a	b	c	d
a	c	b	c	c
b	b	c	c	c
c	a	b	c	d
d	a	b	c	d

Then  $(X, *, E)$ , where  $E = \{c, d\}$ , satisfies (eGE1), (eGE2) and (eGE3), but it does not satisfy (eGE4), since

$$a * (b * a) = a * b = b \neq c = a * c = a * (b * c) = a * (b * (a * a)).$$

**Theorem 2.5.** *Let  $(X, *, E)$  be an eGE-algebra. If  $E$  is a singleton set, then  $(X, *, E)$  is a GE-algebra.*

**Proof.** Let  $E = \{a\}$  be a singleton set. If we put  $1 = a$ , then  $(X, *, 1)$  is a GE-algebra. ■

**Theorem 2.6.** *Let  $(X, *, E_i)$ , for  $i = 1, 2$ , be two eGE-algebras. Then  $(X, *, E_1 \cap E_2)$  is also an eGE-algebra.*

**Proof.** Let  $u \in X$ . Since  $u * u \in E_1$  and  $u * u \in E_2$ , we have  $u * u \in E_1 \cap E_2$ , and so (eGE1) holds. Let  $a \in u * (E_1 \cap E_2)$ . Then, we can find  $b \in E_1 \cap E_2$  such that  $a = u * b$ . Since  $b \in E_1$ ,  $u * b \in E_1$  and  $b \in E_2$ ,  $u * b \in E_2$ , we have  $a = u * b \in E_1 \cap E_2$  and so  $u * (E_1 \cap E_2) \subseteq E_1 \cap E_2$ . Hence (eGE2) holds. Let  $a \in (E_1 \cap E_2) * u$ . Then, we can find  $b \in E_1 \cap E_2$  such that  $a = b * u$ . Since  $b * u = u$ , we have  $a = u$ , and so  $(E_1 \cap E_2) * u = \{u\}$ . Hence (eGE3) holds. (eGE4) is obvious. Thus  $(X, *, E_1 \cap E_2)$  is also an eGE-algebra. ■

**Corollary 2.7.** *If  $(X, *, E_i)$ , for  $i \in \Lambda$ , is a family of eGE-algebras, then  $(X, *, \bigcap_{i \in \Lambda} E_i)$  is an eGE-algebra.*

**Theorem 2.8.** *Let  $(X, *, E_i)$ , for  $i = 1, 2$ , be two eGE-algebras. Then  $(X, *, E_1 \cup E_2)$  is also an eGE-algebra.*

**Proof.** Let  $u \in X$ . Since  $u * u \in E_1$  and  $u * u \in E_2$ , we have  $u * u \in E_1 \cup E_2$  and so (eGE1) holds. For (eGE2), let  $a \in u * (E_1 \cup E_2)$ . Then, we can find  $b \in E_1 \cup E_2$  such that  $a = u * b$ . If  $b \in E_1$ , then  $a \in E_1$ . Also, if  $b \in E_2$ , then  $a \in E_2$ . Thus  $a \in E_1 \cup E_2$  and so  $u * (E_1 \cup E_2) \subseteq E_1 \cup E_2$ . Let  $a \in (E_1 \cup E_2) * u$ . Then, we can find  $b \in E_1 \cup E_2$  such that  $a = b * u$ . Since  $b * u = u$ , we have  $a = u$  and so  $(E_1 \cup E_2) * u = \{u\}$ . Therefore (eGE3) holds. (eGE4) is obvious. Thus  $(X, *, E_1 \cup E_2)$  is an eGE-algebra. ■

**Corollary 2.9.** *If  $(X, *, E_i)$ , for  $i \in \Lambda$ , is a family of eGE-algebras, then  $(X, *, \bigcup_{i \in \Lambda} E_i)$  is also an eGE-algebra.*

**Lemma 2.10.** *Let  $(X, *, E)$  be an eGE-algebra and  $u, v \in X$ . Then  $u * (u * v) = u * v$ .*

**Proof.** Let  $u, v \in X$ . Using (eGE1), (eGE3) and (eGE4), we get

$$u * (u * v) = u * ((u * u) * (u * v)) = u * ((u * u) * v) = u * v. \quad \blacksquare$$

**Theorem 2.11.** *Every self-distributive eBE-algebra is an eGE-algebra.*

**Proof.** Let  $(X, *, E)$  be a self-distributive eBE-algebra and  $u, v, w \in X$ . Then, by (eBE1), (eBE3), (eBE4), and self-distributivity,

$$u * (v * w) = (u * u) * (u * (v * w)) = u * (u * (v * w)) = u * (v * (u * w)).$$

Hence  $X$  is an eGE-algebra. ■

The converse of the Theorem 2.11 does not have to be true. From Example 2.3, we can observe that  $X$  is an eGE-algebra, but not a self-distributive eBE-algebra.

**Theorem 2.12.** *Let  $(X, *, E)$  be an eBE-algebra having the property  $u*(u*v) = u*v$ , for all  $u, v \in X$ . Then  $X$  is an eGE-algebra.*

**Proof.** Let  $u, v, w \in X$  and  $u*(u*v) = u*v$ . Then  $u*(v*w) = u*(u*(v*w)) = u*(v*(u*w))$ . Hence  $X$  is an eGE-algebra. ■

**Proposition 2.13.** *Let  $(X, *, E)$  be an eGE-algebra. Then*

- (i)  $(X; *, X \setminus E)$  is not an eGE-algebra,
- (ii)  $v*w \in E$  implies  $u*(v*w) \in E$ ,
- (iii)  $u*(v*u) \in E$ ,
- (iv)  $u \leq v*w$  implies  $v \leq u*w$ ,
- (v)  $u \leq (u*v)*u$ ,
- (vi)  $u*(v*w) \in E$  implies  $v*(u*w) \in E$  and  $v*(u*(v*w)) \in E$ ,
- (vii)  $u*(v*w) \leq v*(u*w)$ ,
- (viii)  $u*(v*w) \notin E$  implies  $u*w \notin E$  for all  $u, v, w \in X$ .

**Proof.** (i) (eGE2) does not hold, since  $u*E \not\subseteq X \setminus E$  and  $u*E \subseteq E$ .

(ii) By (eGE2), (ii) is obvious.

(iii) Using (eGE4), (eGE1) and (eGE2), we have

$$u*(v*u) = u*(v*(u*u)) \in u*E \subseteq E.$$

(iv) Let  $u \leq v*w$ . Hence  $u*(v*w) \in E$ . Then, by (eGE4) and (eGE2), we have  $v*(u*w) = v*(u*(v*w)) \in v*E \subseteq E$ . Therefore  $v \leq u*w$ .

(v) From (eGE4), (eGE1) and (eGE2) we have

$$u*((u*v)*u) = u*((u*v)*(u*u)) \in u*E \subseteq E.$$

Therefore  $u \leq (u*v)*u$ .

(vi) Applying (iv) and (eGE4), we can prove (vi).

(vii) By routine calculation we can see that

$$\begin{aligned} & (u*(v*w))*(v*(u*w)) \\ &= (u*(v*w))*(v*((u*(v*w)))) \\ &= (u*(v*w))*(v*((u*(v*w))*(u*(v*w)))) \in E. \end{aligned}$$

Thus  $u*(v*w) \leq v*(u*w)$ .

(viii) It is obvious by (ii). ■

**Theorem 2.14.** *Let  $(X, *, E)$  be an eGE-algebra. The following are equivalent.*

- (i)  $u*v \leq (w*u)*(w*v)$ ,
- (ii)  $u*v \leq (v*w)*(u*w)$

for all  $u, v, w \in X$ .

**Proof.** (i) $\Rightarrow$ (ii) Let  $u, v, w \in X$  and assume (i). Then

$$(u * v) * ((w * u) * (w * v)) \in E.$$

Hence, by (eGE4) and (eGE2), we get

$$(u * v) * ((v * w) * (u * w)) = (u * v) * ((v * w) * ((u * v) * (u * w))) \in E.$$

Therefore  $u * v \leq (v * w) * (u * w)$ .

(ii) $\Rightarrow$ (i) Let  $u, v, w \in X$  and assume (ii). Then  $(u * v) * ((v * w) * (u * w)) \in E$ . Hence, by (eGE4) and (eGE2), we get

$$(u * v) * ((w * u) * (w * v)) = (u * v) * ((w * u) * ((u * v) * (w * v))) \in E.$$

Therefore  $u * v \leq (w * u) * (w * v)$ . ■

**Definition 2.15.** An eGE-algebra  $(X, *, E)$  is said to be transitive if it satisfies:

$$(12) \quad (\forall u, v, w \in X) (u * v \leq (w * u) * (w * v)).$$

**Example 2.16.** Let  $X = \{a, b, c, d\}$  be a set and  $*$  a binary operation given in the following table:

$*$	a	b	c	d
a	d	d	d	d
b	a	c	c	c
c	a	b	c	d
d	a	b	c	d

Then  $(X, *, E)$ , where  $E = \{c, d\}$ , is a transitive eGE-algebra but not an eBE-algebra, since  $a * (b * c) = a * c = d \neq c = b * d = b * (a * c)$ .

The following theorem can be proved easily.

**Theorem 2.17.** Let  $(X, *, E)$  be a transitive eGE-algebra. The following hold:

- (1)  $u \leq v$  implies  $w * u \leq w * v$ ,
- (2)  $u * v \leq (v * w) * (u * w)$ ,
- (3)  $u \leq v$  implies  $v * w \leq u * w$ ,
- (4)  $((u * v) * v) * w \leq u * w$ ,
- (5)  $u \leq v$  and  $v \leq w$  imply  $u \leq w$ ,
- (6)  $u * (v * w) \leq (u * v) * (u * w)$

for all  $u, v, w \in X$ .



**Theorem 2.18.** *Let  $(X, *, E)$  be an eGE-algebra. Consider  $Y := (X \setminus E) \cup \{1\}$  and define the operation  $\triangleright$  on  $Y$  as follows:*

$$u \triangleright v = \begin{cases} u * v & \text{if } u, v \neq 1 \text{ and } u * v \notin E, \\ 1 & \text{if } u, v \neq 1 \text{ and } u * v \in E, \\ v & \text{if } u = 1, \\ 1 & \text{if } v = 1. \end{cases}$$

*Then  $(Y, \triangleright, 1)$  is a GE-algebra.*

**Proof.** By (eGE1),  $u * u \in E$ , for all  $u \in X$ . Thus  $u \triangleright u = 1$ , for all  $u \in Y$ , and so (GE1) holds. By definition of  $\triangleright$ , (GE2) hold. To prove  $(Y; \triangleright, 1)$  is a GE-algebra it is sufficient to prove that  $u \triangleright (v \triangleright w) = u \triangleright (v \triangleright (u \triangleright w))$ , for all  $u, v, w \in Y$ . If  $u = 1$  or  $v = 1$  or  $w = 1$ , then we have  $u \triangleright (v \triangleright w) = u \triangleright (v \triangleright (u \triangleright w))$ . Now, let  $u, v, w \neq 1$ . If  $v * w \in E$ , then  $v \triangleright w = 1$ , and so  $u \triangleright (v \triangleright w) = 1$ . On the other hand, if  $u * w \in E$ , then  $u \triangleright w = 1$  and  $u \triangleright (v \triangleright (u \triangleright w)) = u \triangleright (v \triangleright 1) = u \triangleright 1 = 1 = u \triangleright (v \triangleright w)$ . If  $u * w \notin E$ , then  $u * w = u \triangleright w$ . By Proposition 2.13(ii), and  $v * w \in E$ , we have  $v * (u * w) \in E$ . Hence  $u \triangleright (v \triangleright (u \triangleright w)) = u \triangleright 1 = 1 = u \triangleright (v \triangleright w)$ . If  $v * w \notin E$ , then  $v \triangleright w = v * w$ . We have two cases:  $u * (v * w) \in E$  or  $u * (v * w) \notin E$ . If  $u * (v \triangleright w) = u * (v * w) \in E$ , then  $u \triangleright (v \triangleright w) = 1$ . By Proposition 2.13(vi),  $v * (u * w) \in E$ , and so  $u \triangleright (v \triangleright (u \triangleright w)) = u \triangleright (v \triangleright (u * w)) = u \triangleright 1 = 1$ . Thus  $u \triangleright (v \triangleright w) = u \triangleright (v \triangleright (u \triangleright w))$ , in this case. If  $u * (v * w) \notin E$  then, by Proposition 2.13(ii & viii),  $u * w \notin E, v * w \notin E$  and  $v * (u * w) \notin E$ . So that  $u \triangleright w = u * w, v \triangleright w = v * w$  and  $v \triangleright (u * w) = v * (u * w)$ . Hence  $u \triangleright (v \triangleright w) = u * (v * w)$ . Also, by (eGE4),  $u * (v * (u * w)) = u * (v * w) \notin E$ . Hence  $u \triangleright (v \triangleright (u \triangleright w)) = u * (v \triangleright (u \triangleright w)) = u * (v * (u * w)) = u * (v * w) = u \triangleright (v \triangleright w)$ . Thus  $u \triangleright (v \triangleright w) = u \triangleright (v \triangleright (u \triangleright w))$ . Therefore  $(Y, \triangleright, 1)$  is a GE-algebra. ■

**Example 2.19.** Let  $X = \{a, b, c, d\}$  and  $E = \{c, d\}$ . A binary operation  $*$  on  $X$  is given in the following table:

*	a	b	c	d
a	c	b	c	c
b	d	d	d	d
c	a	b	c	d
d	a	b	c	d

Then  $(X, *, E)$  is an eGE-algebra which is not an eBE-algebra.

Now,  $Y = (X \setminus E) \cup \{1\} = \{1, a, b\}$ . Define  $\triangleright$  on  $Y$  with the following table:

$\triangleright$	1	a	b
1	1	a	b
a	1	1	b
b	1	1	1

Then  $(Y, \triangleright, 1)$  is a GE-algebra.

We conclude this section with the following theorem whose proof is straightforward.

**Theorem 2.20.** *Let  $(X, *, 1)$  be a GE-algebra and  $E_0$  be a set such that  $E_0 \cap X = \emptyset$ . If we define  $Y = X \cup E_0$ ,  $E = E_0 \cup \{1\}$  and define the operation  $\triangleleft$  on  $Y$  as follows:*

$$x \triangleleft y = \begin{cases} x * y & \text{if } x, y \notin A_0, \\ y & \text{otherwise.} \end{cases}$$

*Then  $(Y, \triangleleft, E)$  is an eGE-algebra.*

### 3. QUOTIENT EGE-ALGEBRAS

In this section, we introduce the notion of a filter in an eGE-algebra and study its properties. We construct a quotient eGE-algebra via a filter of an eGE-algebra. Throughout this section,  $X$  means  $(X, *, E)$  is an eGE-algebra, unless specified otherwise.

**Definition 3.1.** A subset  $F$  of  $X$  is called a filter of  $X$  if it satisfies:

- (eGEF1)  $E \subseteq F$ ,
- (eGEF2)  $u \in F$  and  $u * v \in F$  imply  $v \in F$ .

The set of all filters of  $X$  will be denoted by  $\mathcal{F}(X)$ . Clearly,  $\mathcal{F}(X) \neq \emptyset$ , since  $X \in \mathcal{F}(X)$ .

**Example 3.2.** Let  $X = \{a, b, c, d, e\}$  be a set and  $*$  a binary operation on  $X$  given in the following table:

*	a	b	c	d	e
a	c	c	c	c	c
b	a	d	d	d	e
c	a	b	c	d	e
d	a	b	c	d	e
e	a	b	c	d	d

Then  $(X, *, E)$ , where  $E = \{c, d\}$ , is an eGE-algebra. Let  $F = \{c, d, e\}$ . Then  $F \in \mathcal{F}(X)$ .

**Proposition 3.3.** *Let  $F \in \mathcal{F}(X)$ . If  $u \in F$  and  $u \leq v$ , then  $v \in F$ .*

**Proof.** Let  $u \in F$  and  $u \leq v$ . Then  $u * v \in E$  and  $E \subseteq F$ . So that  $u * v \in F$ . Hence  $v \in F$ , since  $u \in F$  and  $F$  is a filter of  $X$ . ■

**Theorem 3.4.** *In  $X$ ,  $E \in \mathcal{F}(X)$ .*

*Proof.* Clearly (eGEF1) holds, since  $E \subseteq E$ . Now we prove (eGEF2). Let  $u, u * v \in E$ . Now, by (eGE3), we have  $v = u * v \in E$ . Therefore  $E \in \mathcal{F}(X)$ . ■

**Proposition 3.5.** *If  $F_i \in \mathcal{F}(X)$ , for  $i \in \Lambda$ , then  $\bigcap_{i \in \Lambda} F_i \in \mathcal{F}(X)$ .*

**Theorem 3.6.** *Let  $F \in \mathcal{F}(X)$ . Then  $F_1 = (F \setminus E) \cup \{1\}$  is a filter of  $(Y, \triangleright, 1)$ , which is defined in Theorem 2.18.*

*Proof.* Clearly  $1 \in F_1$ . Let  $u \in F_1$  and  $u \triangleright v \in F_1$ . If  $u = 1$ , then  $v = 1 \triangleright v \in F_1$ . Let  $u \neq 1$ . If  $v = 1$ , then  $v \in F_1$ . If  $v \neq 1$ . Then  $u \in F \setminus E$  and  $v \in X \setminus E$ . If  $u \triangleright v = 1$  by definition of  $\triangleright$  we get  $u * v \in E$ . Then  $u * v \in F$ , since  $F$  is a filter of  $X$ . Hence  $v \in F$ . Thus  $v \in F_1$ . If  $u \triangleright v \neq 1$ , then by definition of  $\triangleright$ ,  $u * v \notin E$  and  $u \triangleright v = u * v \in F_1$ . Thus  $u * v \in F$ . Since  $F \in \mathcal{F}(X)$ , we have  $v \in F$ . Hence  $v \in F_1$ . Therefore  $F_1 \in \mathcal{F}(Y)$ . ■

**Example 3.7.** From Theorem 2.18 and Example 2.19, we get  $Y = \{1, a, b\}$  with the following table:

$\triangleright$	1	a	b
1	1	a	b
a	1	1	b
b	1	1	1

which is a GE-algebra. We can observe that  $F = \{a, c, d\}$  is a filter of  $(X, *, E)$  and  $F_1 = (F \setminus E) \cup \{1\} = \{1, a\}$  is a filter of  $(Y, \triangleright, 1)$ .

The following theorem can be proved easily.

**Theorem 3.8.** *Let  $(X, *, 1)$  be a GE-algebra,  $F \in \mathcal{F}(X)$  and  $E_0$  be a set such that  $X \cap E_0 = \emptyset$ . Then  $F_0 = F \cup E_0$  is a filter of an eGE-algebra  $(Y, \triangleleft, E)$ , which is defined in Theorem 2.20.*

The following example describes the above theorem

**Example 3.9.** Let  $X = \{1, a, b\}$  and  $E_0 = \{c, d\}$ . According to Example 2.19,  $(X, \triangleright, 1)$  is a GE-algebra. We can observe that  $F = \{1, a\}$  is a filter of  $X$ . By Theorem 2.20, we get  $Y = \{1, a, b, c, d\}$ ,  $E = \{1, c, d\}$  and  $(Y, \triangleleft, E)$  is an eGE-algebra with the following table:

$\triangleleft$	1	a	b	c	d
1	1	a	b	c	d
a	1	1	b	c	d
b	1	1	1	c	d
c	1	a	b	c	d
d	1	a	b	c	d

We can observe that  $F_0 = F \cup E_0 = \{1, a, c, d\}$  is a filter of  $Y$ .

**Proposition 3.10.** *A non-empty subset  $F$  of an eGE-algebra  $X$  is a filter of  $X$  if and only if it satisfies:*

- (i)  $E \subseteq F$ ,
- (ii)  $u * (v * w) \in F, v \in F$  implies  $u * w \in F$

for all  $u, v, w \in X$ .

**Proof.** Suppose  $F \in \mathcal{F}(X)$ . Then  $E \subseteq F$ . Let  $u, v, w \in X$  be such that  $u * (v * w) \in F$  and  $v \in F$ . Then, by Theorem 2.13(vii) and Proposition 3.3, we have  $v * (u * w) \in F$ . Then  $u * w \in F$ . Conversely, assume that the conditions hold. It is sufficient to prove (eGEF2). Let  $x \in F$  and  $x * y \in F$ . Then  $x * x \in E \subseteq F$  and  $(x * x) * (x * y) = x * y \in F$ . Hence  $(x * x) * y = y \in F$ . Thus  $F \in \mathcal{F}(X)$ . ■

**Theorem 3.11.** *Let  $F$  be a subset of  $X$  satisfying the following conditions:*

- (eGEF1)  $E \subseteq F$ ,
- (eGEF3)  $u \in X$  and  $r \in F$  imply  $u * r \in F$ ,
- (eGEF4)  $u \in X, r, s \in F$  imply  $(r * (s * u)) * u \in F$ .

Then  $F \in \mathcal{F}(X)$ .

**Proof.** It is sufficient to prove (eGEF2). Let  $u \in F$  and  $u * v \in F$ . Then, by (eGE1), (eGE3) and (eGEF4),  $v = [(u * v) * (u * v)] * v \in F$  and hence (eGEF2) holds. Therefore  $F \in \mathcal{F}(X)$ . ■

**Theorem 3.12.** *If  $X$  is an eGE-algebra and  $F$  is a filter of  $X$ , then  $F$  satisfies (eGEF1), (eGEF3) and (eGEF4).*

**Proof.** It is sufficient to prove (eGEF3) and (eGEF4). Let  $F \in \mathcal{F}(X)$  and  $r \in F, u \in X$ . Then  $r * (u * r) \in E \subseteq F$  and hence, by (GEF2),  $u * r \in F$ . Let  $r, s \in F$ . Since  $r * ((r * (s * u)) * (s * u)) \in E \subseteq F$  and  $r \in F$ , we have  $(r * (s * u)) * (s * u) \in F$ . Hence, by (eGE4) and (eGEF3),  $s * ((r * (s * u)) * u) = s * ((r * (s * u)) * (s * u)) \in F$ . Thus, by (eGEF2),  $(r * (s * u)) * u \in F$ . ■

**Theorem 3.13.** *Let  $F \in \mathcal{F}(X)$ . Then  $(r * u) * u \in F$  for all  $r \in F$  and  $u \in X$ .*

For a non-empty subset  $I$  of  $X$ , we define the binary relation  $\sim_I$  in the following way:

$$u \sim_I v \text{ if and only if } u * v \in I \text{ and } v * u \in I.$$

The set  $\{s \mid r \sim_I s\}$  will be denoted by  $[r]_I$ .

**Lemma 3.14.** *In the above relation  $\sim_I$ , if  $E \subseteq I$  and  $r \in E$ , then  $[r]_I = I$ .*

**Proof.** Let  $u \in I$  and  $r \in E$ . By (eGE3), we have  $r * u \in E * u = \{u\} \subseteq I$  and so  $r * u \in I$ . From (eGE2), we have  $u * r \in u * E \subseteq E \subseteq I$ , then  $u * r \in I$ . Hence  $r \sim_I u$ . Therefore  $I \subseteq [r]_I$ . Conversely, let  $r \in E$  and  $u \in [r]_I$ . Then  $u \sim_I r$  and so  $u * r \in I$  and  $r * u = u \in I$ . Hence  $[r]_I \subseteq I$ . Therefore  $[r]_I = I$ . ■

**Theorem 3.15.** *Let  $(X, *, E)$  be a transitive eGE-algebra and  $F \in \mathcal{F}(X)$ . Then  $\sim_F$  is a congruence relation on  $X$ .*

**Proof.** Since  $u * u \in E \subseteq F$ , we have  $u * u \in F$ , and so  $u \sim_F u$ . If  $u \sim_F v$ , then clearly  $v \sim_F u$ . Now, let  $u \sim_F v$  and  $v \sim_F w$ . Then  $u * v, v * u \in F$  and  $v * w, w * v \in F$ . By Proposition 2.17(1), we have  $v * w \leq (u * v) * (u * w)$ , and so by Proposition 3.3, we have  $(u * v) * (u * w) \in F$ . Since  $F$  is a filter and  $u * v \in F$ , we have  $u * w \in F$ . Similarly, we can prove that  $w * u \in F$ . Thus  $u \sim_F w$ . Therefore  $\sim_F$  is an equivalent relation on  $X$ . If  $r \sim_F s$  and  $u \sim_F v$ , then  $r * s, s * r \in F$  and  $u * v, v * u \in F$ . By Proposition 2.17(1), we have  $u * v \leq (r * u) * (r * v)$  and  $v * u \leq (r * v) * (r * u)$ , and so by Proposition 3.3, we have  $(r * u) * (r * v) \in F$  and  $(r * v) * (r * u) \in F$ . Thus  $r * u \sim_F r * v$ . Similarly, we can prove that  $r * v \sim_F s * v$ . Since the relation  $\sim_F$  is transitive, we have  $r * u \sim_F s * v$  which proves that  $\sim_F$  is a congruence relation on  $X$ . ■

**Proposition 3.16.** *Let  $\sim_G$  be a congruence relation on  $X$ ,  $E \subseteq G$  and  $r \in E$ . Then  $[r]_G \in \mathcal{F}(X)$ .*

**Proof.** By Lemma 3.14, we have  $[r]_G = G$ . Let  $u, u * v \in [r]_G$ . Thus  $u \sim_G r$  and  $u * v \sim_G r$ . Since  $v \sim_G v$  and  $\sim_G$  is a congruence relation, we can observe that  $r \sim_G u * v \sim_G r * v = v$  (by (eGE3)). Thus  $v \in [r]_G$ . Therefore  $[r]_G \in \mathcal{F}(X)$ . ■

Denote  $\frac{X}{\sim_G} = \{[u]_G \mid u \in X\}$ . Define a binary operation  $\bullet$  on  $\frac{X}{\sim_G}$  by  $[u]_G \bullet [v]_G := [u * v]_G$ . Then by above theorem,  $\bullet$  is well defined. The following theorem shows that for a transitive eGE-algebra  $(X, *, E), r \in E$  and  $F \in \mathcal{F}(X)$ , the quotient algebra  $\left(\frac{X}{\sim_{[r]_F}}, \bullet, [r]_F\right)$  is a GE-algebra.

**Theorem 3.17.** *Let  $(X, *, E)$  be a transitive eGE-algebra,  $F \in \mathcal{F}(X)$  and  $r \in E$ . Then  $\left(\frac{X}{\sim_{[r]_F}}, \bullet, [r]_F\right)$  is a GE-algebra.*

**Proof.** Since  $E \subseteq F$ , we can observe that  $E \subseteq [r]_F$ , for all  $r \in E$ . Hence  $[r]_F$  is a filter by Proposition 3.16 and so  $\sim_{[r]_F}$  is a congruence relation on  $X$  by Theorem 3.15. Now, we have

- (GE1)  $[u]_F \bullet [u]_F = [u * u]_F = [r]_F$ , since  $u * u \in E \subseteq [r]_F$ ,
  - (GE2)  $[r]_F \bullet [u]_F = [r * u]_F = [u]_F$ , since  $E * u = \{u\}$  and so  $r * u = u$ ,
  - (GE3)  $[u]_F \bullet ([v]_F \bullet [w]_F) = [u]_F \bullet [v * w]_F = [u * (v * w)]_F = [u * (v * (u * w))]_F = [u]_F \bullet [(v * (u * w))]_F = [u]_F \bullet ([v]_F \bullet [u * v]_F) = [u]_F \bullet ([v]_F \bullet ([u]_F \bullet [w]_F))$ .
- Thus  $\left(\frac{X}{\sim_{[r]_F}}, \bullet, [r]_F\right)$  is a GE-algebra. ■

## REFERENCES

- [1] R.K. Bandaru, A. Borumand Saeid and Y.B. Jun, *On GE-algebras*, Bulletin of the Section of Logic, (2020), in press.  
<https://doi.org/10.18778/0138-0680.2020.20>
- [2] A. Borumand Saeid, A. Rezaei and R.A. Borzooei, *Some types of filters in BE-algebras*, Math. Comput. Sci. **7** (2013) 341–352.  
<https://doi.org/10.1007/s11786-013-0157-6>
- [3] S. Celani, *A note on homomorphisms of Hilbert algebras*, Internat. J. Math. and Math. Sci. **29** (2002) 55–61.  
<https://doi.org/10.1155/S0161171202011134>
- [4] I. Chajda and R. Halas, *Congruences and ideals in Hilbert algebras*, Kyungpook Math. J. **39** (1999) 429–432.
- [5] I. Chajda, R. Halas and Y.B. Jun, *Annihilators and deductive systems in commutative Hilbert algebras*, Comment. Math. Univ. Carolinae **43** (2002) 407–417.
- [6] A. Diego, *Sur les algebres de Hilbert*, Collection de Logique Mathematique, Edition Hermann, Ser. A, XXI, 1966.
- [7] Y.B. Jun, *Commutative Hilbert algebras*, Soochow J. Math. **22** (1996) 477–484.
- [8] Y.B. Jun and K.H. Kim, *H-filters of Hilbert algebras*, Sci. Math. Jpn. e-2005, 231–236.
- [9] S.M. Hong and Y.B. Jun, *On deductive systems of Hilbert algebras*, Comm. Korean Math. Soc. **11** (1996) 595–600.
- [10] I. Iséki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Jpn. **23** (1978) 1–26.
- [11] K. Iseki, *On BCI-algebras*, Mathematics Seminar Notes (Kobe University) **8** (1980) 125–130.
- [12] H.S. Kim and Y.H. Kim, *On BE-algebras*, Sci. Math. Jpn. **66** (2007) 113–116.
- [13] A.S. Nasab and A. Borumand Saeid, *Semi maximal filter in Hilbert algebras*, J. Intelligent & Fuzzy Systems **30** (2016) 7–15.  
<https://doi.org/10.3233/IFS-151706>
- [14] A.S. Nasab and A. Boruman Saeid, *Stonean Hilbert algebra*, J. Intelligent & Fuzzy Systems **30** (2016) 485–492.  
<https://doi.org/10.3233/IFS-151773>
- [15] A.S. Nasab and A. Borumand Saeid, *Study of Hilbert algebras in point of filters*, An St. Univ. Ovidius Constanta **24** (2016) 221–251.  
<https://doi.org/10.1515/auom-2016-0039>
- [16] A. Rezaei, A. Borumand Saeid and R.A. Borzooei, *Relation between Hilbert algebras and BE-algebras*, Appl. Math. **8** (2013) 573–584.

- [17] A. Rezaei and A. Borumand Saeid, *Relation between Dual S-algebras and BE-algebras*, *Le Matematiche* **70** (2015) 71–79.  
<https://doi.org/10.4418/2015.70.1.5>
- [18] A. Rezaei and A. Borumand Saeid, *Relation between BE-algebras and g-Hilbert algebras*, *Discuss. Math. Gen. Alg. Appl.* **38** (2018) 33–45.  
<https://doi.org/10.7151/dmgaa.1285>
- [19] A. Rezaei and A. Borumand Saeid, *Some results in BE-algebras*, *An. Univ. Oradea, Fasc. Math.* **19** (2012) 33–44.
- [20] A. Rezaei, A. Borumand Saeid and A. Radfar, *On eBE-algebra*, *TWMS J. Pure Appl. Math.* **7** (2016) 200–210.

Received 15 September 2020

Revised 24 October 2020

Accepted 24 October 2020