

ON THE SIMPLICITY AND SUBDIRECT IRREDUCIBILITY
OF BOOLEAN ULTRAPOWERS

BY

STANLEY BURRIS AND ERROL JEFFERS (WATERLOO, ONTARIO)

In [2], Frayne et al. gave an example of a simple group with ultrapowers which are not simple. In this paper* we will obtain necessary and sufficient conditions for a Boolean ultrapower to be simple, or subdirectly irreducible, provided the language is countable.

Let $\mathfrak{A} = \langle A, \mathcal{F} \rangle$ be an algebra, and $\mathfrak{B} = \langle B, \vee, \wedge, ', 0, 1 \rangle$ a Boolean algebra. Assume that \mathfrak{B} is complete if \mathfrak{A} is infinite. The *Boolean power* $\mathfrak{A}[\mathfrak{B}]$ has as its universe (written $|\mathfrak{A}[\mathfrak{B}]|$) the set of all mappings α of A into B such that

- (i) if $a, b \in A$, $a \neq b$, then $\alpha(a) \wedge \alpha(b) = 0$;
- (ii) $\bigvee_{a \in A} \alpha(a) = 1$.

The fundamental operations are defined by

- (iii) $f(\alpha_0, \dots, \alpha_{n-1})(a) = \bigvee \{ \alpha_0(a_0) \wedge \dots \wedge \alpha_{n-1}(a_{n-1}) : f(a_0, \dots, a_{n-1}) = a \}$.

Let \mathcal{U} be an ultrafilter on a Boolean algebra \mathfrak{B} . Define the relation $\theta_{\mathcal{U}}(\mathfrak{A})$ on $\mathfrak{A}[\mathfrak{B}]$ by

$$\theta_{\mathcal{U}}(\mathfrak{A}) = \{ \langle \alpha, \beta \rangle \in |\mathfrak{A}[\mathfrak{B}]|^2 : \bigvee_{a \in A} \alpha(a) \wedge \beta(a) \in \mathcal{U} \}.$$

It can easily be shown that $\theta_{\mathcal{U}}(\mathfrak{A})$ is a congruence on $\mathfrak{A}[\mathfrak{B}]$. We denote the quotient algebra $\mathfrak{A}[\mathfrak{B}]/\theta_{\mathcal{U}}(\mathfrak{A})$ by $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$, and call it a *Boolean ultrapower* of \mathfrak{A} . For $\xi \in |\mathfrak{A}[\mathfrak{B}]|$ let $[\xi]_{\mathcal{U}}$ denote the image in $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$.

Remark 1. If $\mathfrak{B} \cong \mathfrak{2}^I$ for some I (where $\mathfrak{2}$ is the two-element Boolean algebra), then $\mathfrak{A}[\mathfrak{B}] \cong \mathfrak{A}[\mathfrak{2}^I] \cong \mathfrak{A}^I$. Therefore, $\mathfrak{A}[\mathfrak{B}]/\mathcal{U} \cong \mathfrak{A}^I/\mathcal{U}$, and the Boolean ultrapower in this case is just the familiar ultrapower.

An algebra \mathfrak{A} is *simple* if $|A| > 1$ and the only congruence relations on \mathfrak{A} are Δ_A and ∇_A , where $\Delta_A = \{ \langle a, a \rangle : a \in A \}$, and $\nabla_A = A \times A$. An algebra \mathfrak{A} is said to be *(a, b)-irreducible* if $a \neq b$ and every non-trivial

* This research was supported by NRC Grant No. A7256.

congruence on \mathfrak{A} identifies a and b . An algebra \mathfrak{A} is said to be *subdirectly irreducible* if there are $a, b \in A$ such that \mathfrak{A} is (a, b) -irreducible. A *simplicity sentence* is a first-order sentence all models of which are simple. Similarly we define a *subdirect irreducibility sentence*.

An ultrafilter \mathcal{U} on a Boolean algebra is said to be ω -complete if, whenever $\{x_n: n < \omega\} \subseteq \mathcal{U}$,

$$\bigwedge_{n < \omega} x_n \in \mathcal{U}.$$

\mathcal{U} is said to be ω -incomplete if it is not ω -complete.

Remark 2. Principal ultrafilters on Boolean algebras are always ω -complete. Therefore, \mathfrak{B} must be infinite in order that ω -incomplete ultrafilters may exist.

An algebra \mathfrak{A} is α -saturated if every set of formulae $\{\sigma_i(x_0): i \in I\}$ in the language of \mathfrak{A} , with fewer than α parameters from $|\mathfrak{A}|$, which is finitely satisfiable in \mathfrak{A} is also satisfiable in \mathfrak{A} .

From now on we assume that the language of \mathfrak{A} is countable.

LEMMA 1. *An ω -saturated algebra \mathfrak{A} satisfies a simplicity (subdirect irreducibility) sentence iff \mathfrak{A} is simple (subdirectly irreducible).*

Proof. For the non-trivial direction, assume that \mathfrak{A} does not satisfy a simplicity sentence. Taylor has shown in [4] that, for any $a, b, c, d \in A$, $(c, d) \in \theta(a, b)$ iff there exists an existential positive formula $\varphi(x, y, u, v)$ which satisfies certain conditions, and $\mathfrak{A} \models \varphi(a, b, c, d)$. Let $\{\varphi_i(x, y, u, v)\}_{i < \omega}$ be all such formulae in our language. From [4] one can conclude the following:

(1) \mathfrak{A} is not simple iff, for some $a, b, c, d \in A$,

$$\mathfrak{A} \models \neg \varphi_i(a, b, c, d) \& a \neq b \quad \text{for all } i < \omega;$$

(2) for every choice of $\varphi_{i_0}(x, y, u, v), \dots, \varphi_{i_n}(x, y, u, v)$,

$$\forall xyuv (x \neq y \rightarrow \varphi_{i_0} \vee \dots \vee \varphi_{i_n})$$

is a simplicity sentence.

Therefore, suppose that \mathfrak{A} does not satisfy a simplicity sentence. Then, for any $i_0, \dots, i_n < \omega$,

(3) $\mathfrak{A} \models \neg \forall xyuv (x \neq y \rightarrow \varphi_{i_0} \vee \dots \vee \varphi_{i_n})$.

Let Γ be the set of all formulae of the form

$$(x \neq y) \& (\neg \varphi_i(x, y, u, v)).$$

From (3) we see that Γ is finitely satisfiable in \mathfrak{A} . Since \mathfrak{A} is ω -saturated, and the members of Γ contain no parameters from $|\mathfrak{A}|$, Γ is satisfiable in \mathfrak{A} . (1) now implies that \mathfrak{A} is not simple. A similar proof holds for the subdirect irreducibility case.

Now we need two results in [3].

LEMMA 2. *If \mathcal{U} is an ω -incomplete ultrafilter on a Boolean algebra \mathfrak{B} , then $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is ω_1 -saturated.*

LEMMA 3. *Let $\xi_0, \dots, \xi_n \in |\mathfrak{A}[\mathfrak{B}]|$ and suppose that $\sigma([\xi_0]_{\mathcal{U}}, \dots, [\xi_n]_{\mathcal{U}})$ is a sentence. Then*

$$\mathfrak{A}[\mathfrak{B}]/\mathcal{U} \models \sigma([\xi_0]_{\mathcal{U}}, \dots, [\xi_n]_{\mathcal{U}})$$

iff

$$\bigvee \{ \xi_0(a_0) \wedge \dots \wedge \xi_n(a_n) : \mathfrak{A} \models \sigma(a_0, \dots, a_n) \} \in \mathcal{U}.$$

THEOREM 1. *Let \mathfrak{B} be a Boolean algebra, and \mathcal{U} an ultrafilter on \mathfrak{B} . Then $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is simple (subdirectly irreducible) iff either \mathcal{U} is ω -complete and \mathfrak{A} is simple (subdirectly irreducible) or \mathfrak{A} satisfies a simplicity (subdirect irreducibility) sentence.*

Proof. We will consider the case of simplicity — the subdirect irreducibility case has a similar treatment. If \mathfrak{A} satisfies a simplicity sentence, then, since \mathfrak{A} can elementarily be embedded in $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ (see [3]), $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is simple. So assume that \mathfrak{A} is simple and \mathcal{U} is ω -complete. Let the formulae φ mentioned in the proof of Lemma 1 be enumerated as follows: $\varphi_0, \varphi_1, \dots, \varphi_n, \dots$ ($n < \omega$). Let

$$S_i = \{ \langle a_0, a_1, c_0, c_1 \rangle : \mathfrak{A} \models \varphi_i(a_0, a_1, c_0, c_1) \}.$$

Let ξ, η, α, β be arbitrary elements of $|\mathfrak{A}[\mathfrak{B}]|$ such that

$$\bigvee_{a_0 \neq a_1} \xi(a_0) \wedge \eta(a_1) \in \mathcal{U}$$

(i.e. $[\xi]_{\mathcal{U}} \neq [\eta]_{\mathcal{U}}$ in $|\mathfrak{A}[\mathfrak{B}]/\mathcal{U}|$). Then, since \mathfrak{A} is simple,

$$\bigvee_{i < \omega} \bigvee \{ \xi(a_0) \wedge \eta(a_1) \wedge \alpha(c_0) \wedge \beta(c_1) : \langle a_0, a_1, c_0, c_1 \rangle \in S_i \} \in \mathcal{U}.$$

Hence (by ω -completeness) for some $i < \omega$ we have

$$\{ \xi(a_0) \wedge \eta(a_1) \wedge \alpha(c_0) \wedge \beta(c_1) : \langle a_0, a_1, c_0, c_1 \rangle \in S_i \} \in \mathcal{U},$$

i.e.

$$\mathfrak{A}[\mathfrak{B}]/\mathcal{U} \models \varphi_i([\xi]_{\mathcal{U}}, [\eta]_{\mathcal{U}}, [\alpha]_{\mathcal{U}}, [\beta]_{\mathcal{U}}),$$

so $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is simple.

Now, assume that $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is simple and \mathcal{U} is ω -incomplete. By Lemma 2, $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is ω_1 -saturated, hence ω -saturated. By Lemma 1, together with the fact that \mathfrak{A} is isomorphic to an elementary substructure of $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$, the proof is complete.

COROLLARY. *For a given algebra \mathfrak{A} and for a given infinite Boolean algebra \mathfrak{B} , $\mathfrak{A}[\mathfrak{B}]/\mathcal{U}$ is simple (subdirectly irreducible) for all \mathcal{U} iff \mathfrak{A} satisfies a simplicity sentence (subdirect irreducibility sentence).*

In view of Remark 1 the above applies to a special case of ultrapowers. Indeed, a similar result can be stated for ultraproducts.

THEOREM 2. *Let \mathcal{U} be an ultrafilter on a given infinite set I , and assume that the language of our algebras \mathfrak{A}_i is countable. Then $\prod_{i \in I} \mathfrak{A}_i / \mathcal{U}$ is simple (subdirectly irreducible) iff either \mathcal{U} is ω -complete and*

$$\{i \in I: \mathfrak{A}_i \text{ is simple (subdirectly irreducible)}\} \in \mathcal{U}$$

or, for some simplicity sentence (subdirect irreducibility sentence) σ ,

$$\{i \in I: \mathfrak{A}_i \models \sigma\} \in \mathcal{U}.$$

REFERENCES

- [1] J. L. Bell and A. B. Slomson, *Models and ultraproducts*, Amsterdam 1971.
- [2] T. Frayne, A. Morel and D. Scott, *Reduced direct products*, *Fundamenta Mathematicae* 51 (1962), p. 195-228.
- [3] R. Mansfield, *The theory of Boolean ultrapowers*, *Annals of Mathematical Logic* 2 (1971), p. 297-323.
- [4] W. Taylor, *Residually small varieties*, *Algebra Universalis* 2 (1972), p. 33-53.

Reçu par la Rédaction le 14. 7. 1975