

## PRESOLID VARIETIES OF $n$ -SEMIGROUPS

AVAPA CHANTASARTRASSMEE

*The University of the Thai Chamber of Commerce*  
*Department of Mathematics*  
*126/1 Vibhavadee-Rangsit Road*  
*Din Daeng Bangkok 10400, Thailand*  
**e-mail:** avapa\_a@hotmail.com

AND

JÖRG KOPPITZ

*University of Potsdam, Institute of Mathematics*  
*Am Neuen Palais, 14415 Potsdam, Germany*  
**e-mail:** koppitz@rz.uni-potsdam.de

### Abstract

The class of all  $M$ -solid varieties of a given type  $\tau$  forms a complete sublattice of the lattice  $\mathcal{L}(\tau)$  of all varieties of algebras of type  $\tau$ . This gives a tool for a better description of the lattice  $\mathcal{L}(\tau)$  by characterization of complete sublattices. In particular, this was done for varieties of semigroups by L. Polák ([10]) as well as by Denecke and Koppitz ([4], [5]). Denecke and Hounnon characterized  $M$ -solid varieties of semirings ([3]) and  $M$ -solid varieties of groups were characterized by Koppitz ([9]). In the present paper we will do it for varieties of  $n$ -semigroups. An  $n$ -semigroup is an algebra of type  $(n)$ , where the operation satisfies the  $[i, j]$ -associative laws for  $1 \leq i < j \leq n$ , introduced by Dörtnie ([2]). It is clear that the notion of a 2-semigroup is the same as the notion of a semigroup. Here we will consider the case  $n \geq 3$ .

**Keywords:** hypersubstitution, presolid,  $n$ -semigroup.

**2000 Mathematics Subject Classification:** 08B15, 08B25.

## 1. INTRODUCTION

Let  $\tau$  be a fixed type of algebras, with fundamental operation symbols  $f_i$  of arity  $n_i$ , for  $i \in I$ . A hypersubstitution of type  $\tau$  is a mapping which associates to every operation symbol  $f_i$  an  $n_i$ -ary term  $\sigma(f_i)$  of type  $\tau$ . Let  $W_\tau(X)$  be the set of all terms of type  $\tau$  on an alphabet  $X := \{x_1, x_2, x_3, \dots\}$ . By  $W_\tau(X_n)$  ( $X_n := \{x_1, \dots, x_n\}$ ) we denote the set of all  $n$ -ary terms,  $n \geq 1$ . For  $1 \leq m, n \in \mathbb{N}$  we define an operation  $S_m^n : W_\tau(X_n) \times W_\tau(X_m)^n \rightarrow W_\tau(X_m)$  inductively as follows: For  $(t_1, \dots, t_n) \in W_\tau(X_n)^n$  we put:

- (i)  $S_m^n(x_i, t_1, \dots, t_n) := t_i$  for  $1 \leq i \leq n$ ;
- (ii)  $S_m^n(f_i(s_1, \dots, s_{n_i}), t_1, \dots, t_n) := f_i(S_m^n(s_1, t_1, \dots, t_n), \dots, S_m^n(s_{n_i}, t_1, \dots, t_n))$  for  $i \in I, s_1, \dots, s_{n_i} \in W_\tau(X_n)$  where  $S_m^n(s_1, t_1, \dots, t_n), \dots, S_m^n(s_{n_i}, t_1, \dots, t_n)$  will be assumed to be already defined.

Any hypersubstitution  $\sigma$  can be uniquely extended to a mapping  $\hat{\sigma}$  on  $W_\tau(X)$  inductively as follows:

- (i)  $\hat{\sigma}[w] := w$  for  $w \in X$ ;
- (ii)  $\hat{\sigma}[f_i(t_1, \dots, t_{n_i})] := S_m^{n_i}(\sigma(f_i), \hat{\sigma}[t_1], \dots, \hat{\sigma}[t_{n_i}])$  for  $i \in I, t_1, \dots, t_{n_i} \in W_\tau(X_m)$  where  $\hat{\sigma}[t_1], \dots, \hat{\sigma}[t_{n_i}]$  will be assumed to be already defined.

A binary operation  $\circ_h$  can be defined on the set  $Hyp(\tau)$  of all hypersubstitutions of type  $\tau$ , by letting  $\sigma_1 \circ_h \sigma_2 = \hat{\sigma}_1 \circ \sigma_2$ , where  $\circ$  is the usual composition of functions. The set  $Hyp(\tau)$  is closed under this associative operation. It also contains an identity element for  $\circ_h$ , namely the identity hypersubstitution  $\sigma_{id}$  which maps every  $f_i$  to  $f_i(x_1, \dots, x_{n_i})$ . Thus  $Hyp(\tau)$  is a monoid.

Now let  $M$  be any submonoid of  $Hyp(\tau)$ . A variety  $V$  is called  $M$ -solid if for every  $\sigma \in M$  and every identity  $u \approx v$  in  $V$ , the identity  $\hat{\sigma}[u] \approx \hat{\sigma}[v]$  holds in  $V$ . When  $M$  is the whole monoid  $Hyp(\tau)$ , an  $M$ -solid variety is called a solid variety. Two hypersubstitutions  $\sigma_1, \sigma_2$  are said to be  $V$ -equivalent if for every operation symbol  $f_i$  of type  $\tau$ ,  $\sigma_1(f_i) \approx \sigma_2(f_i)$

is an identity in  $V$ . In this case we write  $\sigma_1 \sim_V \sigma_2$ . In [11] it was proved that if  $\widehat{\sigma}_1[s] \approx \widehat{\sigma}_1[t]$  is an identity in  $V$  for given terms  $s, t \in W_\tau(X)$  and  $\sigma_1 \sim_V \sigma_2$  then  $\widehat{\sigma}_2[s] \approx \widehat{\sigma}_2[t]$  is an identity in  $V$ . Therefore, at most one element from each equivalence class of  $\sim_V$  is needed to test the  $M$ -solidity.

The motivation of studying  $M$ -solid varieties comes from following result of Denecke and Reichel in [6]. For each monoid  $M$  of  $Hyp(\tau)$ , the collection of all  $M$ -solid varieties of type  $\tau$  forms a complete lattice, which is a complete sublattice of the lattice  $\mathcal{L}(\tau)$  of all varieties of type  $\tau$ . This lattice  $\mathcal{L}(\tau)$  is in general large and complicated, and difficult to study, and the  $M$ -solid sublattices give us a way to study at least some of its sublattices. Thus it may be useful to study the monoid  $Hyp(\tau)$  and its submonoids  $M$  and the corresponding  $M$ -solid varieties, both in general and for specific type  $\tau$ , and the intersection of the lattice of all  $M$ -solid varieties with a fixed variety of type  $\tau$ . For specific types, much work has been done for type  $\tau = (2)$ , and in particular for varieties of semigroups. L. Polák ([10]) has given a characterization of the lattice of solid semigroup varieties, and various authors have studied  $M$ -solid semigroup varieties for various choices of  $M$ . Moreover, for type  $\tau = (2, 2)$ , in [3], all solid varieties of semirings are determined and, for type  $\tau = (2, 1, 0)$ , J. Koppitz ([9]) determined  $M$ -solid varieties of groups. More informations about hypersubstitutions, one can find in [8].

Our goal in this paper is a similar investigation for type  $(n)$ , for  $n \geq 3$ . Only a few solid varieties of type  $(n)$  have been known (see [1] and [7]). We will consider the concept of an  $n$ -semigroup, which is a natural extension of the concept of a semigroup. An  $n$ -semigroup is an algebra of type  $(n)$ , where the  $n$ -ary operation satisfies the  $[i, j]$ -associative laws

$$\begin{aligned} x_1 \dots x_{i-1} (x_i \dots x_{i+n-1}) x_{i+n} \dots x_{2n-1} &\approx \\ x_1 \dots x_{j-1} (x_j \dots x_{j+n-1}) x_{j+n} \dots x_{2n-1}, &\text{ for } 1 \leq i < j \leq n. \end{aligned}$$

Each  $n$ -group is an  $n$ -semigroup (see Dörnte [2]). Each semigroup  $(S; \cdot)$  induce an  $n$ -semigroup in the following way: Let  $f_n : S^n \rightarrow S$  be defined by  $f_n(a_1, a_2, \dots, a_n) := a_1 \cdot a_2 \cdot \dots \cdot a_n$  (we use the binary operation  $\cdot$  of the given semigroup). Since  $\cdot$  is associative,  $f_n$  satisfies the  $[i, j]$ -associative laws for  $1 \leq i < j \leq n$ , i.e.,  $(S; f_n)$  is an  $n$ -semigroup. Clearly, in the case  $n = 2$  we have the  $[1, 2]$ -associative law  $(x_1 x_2) x_3 \approx x_1 (x_2 x_3)$ . So the notion of a 2-semigroup is the same as the notion of a semigroup.

We also introduce the monoids  $NPer(n)$  and  $Pre(n)$  and give a characterization of all  $NPer(n)$ -solid as well as all  $Pre(n)$ -solid varieties of semigroups.

2. HYPERSUBSTITUTIONS OF TYPE  $(n)$ 

In this section we present some background information about hypersubstitutions and varieties of type  $(n)$ , and introduce the special monoids we shall be studying. We assume throughout a fixed type  $(n)$ , with  $n \geq 3$ , so we have one  $n$ -ary operation symbol which we shall denote by  $f$ . For  $\Sigma$  any set of identities of type  $(n)$ , we will denote by  $Mod(\Sigma)$  the variety determined by the set  $\Sigma$  and by  $IdV$  we denote the set of all identities which hold in a given variety  $V$ . Because of the  $[i, j]$ -associative laws,  $1 \leq i < j \leq n$ , a term over a variety of  $n$ -semigroups can be regarded as a word of the length  $(n-1)r+1$  for a suitable natural number  $r$ . By  $l(t)$  we denote the length of a given term  $t \in W_{(n)}(X)$  and  $var(t)$  means the set of variables occurring in  $t$ . By  $cv(t)$  we mean the cardinality of  $var(t)$ . For example, if  $t = f(x_1, \dots, x_1)$  then  $l(t) = n$ ,  $var(t) = \{x_1\}$ , and  $cv(t) = 1$ . An identity  $u \approx v$  is said to be normal if  $u = v$  or both terms  $u$  and  $v$  are different from a variable. Since any hypersubstitution  $\sigma$  in  $Hyp(n)$  is completely determined by what it does to  $f$ , we will denote by  $\sigma_t$  the hypersubstitution which maps  $f$  to the term  $t$ . For convenience, we list here some sets of terms and varieties of type  $(n)$  that we shall discuss later:

$W_{(n)}^{np}(X_n)$  be the set of all  $t \in W_{(n)}(X_n)$  containing a subword  $s$  with  $n = l(s) > cv(s)$ ;

$\widetilde{W}_{(n)}^{np}(X) := \{t \in W_{(n)}(X) \mid l(t) > cv(t)\}$ ;

$\widetilde{V}_n := Mod\{x_1 \dots x_{2n-1} \approx x_1 \dots x_{i-1} x_{i+1} x_{i+2} x_i x_{i+3} \dots x_{2n-1} \mid 1 \leq i \leq 2n-3\}$ ;

$\widetilde{W}_n := Mod\{t \approx x^n \mid t \in W_{(n)}(X_n), n = l(t) > cv(t)\}$ ;

$V_n := \widetilde{V}_n \cap \widetilde{W}_n$ .

It is easy to verify that there is no nontrivial solid variety of  $n$ -semigroups.

**Theorem 1.** *For each natural number  $n \geq 3$  there is not nontrivial solid variety of  $n$ -semigroups.*

**Proof.** Let  $V$  be a solid variety of  $n$ -semigroups. Then  $\widehat{\sigma}_{x_2}[(x_1 \dots x_n) x_{n+1} \dots x_{2n-1}] \approx \widehat{\sigma}_{x_2}[x_1 \dots x_{n-1} (x_n \dots x_{2n-1})] \in IdV$ , i.e.,  $x_{n+1} \approx x_2 \in IdV$  and  $V$  is the trivial variety of type  $(n)$ . ■

A hypersubstitution  $\sigma$  is called a pre-hypersubstitution if  $\sigma(f)$  is not a variable. The set  $Pre(n)$  of all pre-hypersubstitutions forms a submonoid of the monoid  $Hyp(n)$  of all hypersubstitutions of type  $(n)$ . A variety of  $n$ -semigroups is called presolid if it is  $M$ -solid for  $M = Pre(n)$ . Note that any solid variety is also presolid. By  $S_n$  we will denote the set of all bijections on the set  $\{1, \dots, n\}$ . For  $\pi \in S_n$ , the hypersubstitution  $\sigma$  with  $\sigma(f) = f(x_{\pi(1)}, \dots, x_{\pi(n)})$  will be denoted by  $\sigma_\pi$ . We will use the following notations of sets of hypersubstitutions:

$Pre(n) := Hyp(n) \setminus \{\sigma_{x_i} \mid 1 \leq i \leq n\}$  the set of all pre-hypersubstitutions;

$Per(n) := \{\sigma_\pi \mid \pi \in S_n\}$ ;

$Nper(n) := \{\sigma_t \mid t \in W_{(n)}^{np}(X_n)\} \cup \{\sigma_{id}\}$ .

**Proposition 2.** *For  $2 \leq n \in \mathbb{N}$ ,  $Nper(n)$  forms a monoid.*

**Proof.** We have to check that  $\sigma_1 \circ_h \sigma_2 \in Nper(n)$  for any  $\sigma_1, \sigma_2 \in Nper(n)$ . For this let  $\sigma_1, \sigma_2 \in Nper(n)$ . Then there are  $r, t \in W_{(n)}^{np}(X_n)$  such that  $\sigma_1(f) = r$  and  $\sigma_2(f) = t$ . In particular,  $r$  contains a subword  $s$  with  $n = l(s) > cv(s)$ . Further,  $\hat{\sigma}_1[t]$  contains a subterm  $S_n^n(r, x_{i_1}, \dots, x_{i_n})$ . Since  $r$  contains a subword  $s$  with  $n = l(s) > cv(s)$ , the term  $S_n^n(r, x_{i_1}, \dots, x_{i_n})$  contains a subword  $\tilde{s}$  with  $n = l(\tilde{s}) > cv(\tilde{s})$ . Consequently,  $\hat{\sigma}_1[t]$  contains the subword  $\tilde{s}$  with  $n = l(\tilde{s}) > cv(\tilde{s})$ , i.e.,  $\sigma_1 \circ_h \sigma_2(f) = \hat{\sigma}_1[t] \in W_{(n)}^{np}(X_n)$  and thus  $\sigma_1 \circ_h \sigma_2 \in Nper(n)$ . ■

### 3. PRESOLID VARIETIES OF $n$ -SEMIGROUPS

We begin the investigations of presolid varieties of  $n$ -semigroups by looking for a variety that contains all presolid varieties.

**Proposition 3.** *Let  $3 \leq n \in \mathbb{N}$  and  $V$  be any  $Pre(n)$ -solid variety of  $n$ -semigroups. Then  $V \subseteq \tilde{V}_n$ .*

**Proof.** Let  $\pi \in S_n$  with  $\pi(1) = 2$ ,  $\pi(2) = 1$  and  $\pi(k) = k$  for  $3 \leq k \leq n$ . If we apply  $\sigma_\pi$  to the  $[1, n]$ -associative law we get  $x_{n+1}x_2x_1x_3 \dots x_nx_{n+2} \dots x_{2n-1} \approx x_2x_1x_3 \dots x_{n+1}x_nx_{n+2}x_{n+3} \dots x_{2n-1} \in IdV$  since  $V$  is  $Pre(n)$ -solid. By suitable substitution we get  $x_1 \dots x_{2n-1} \approx x_2 \dots x_nx_1x_{n+1} \dots x_{2n-1} \in IdV$ . If  $n \geq 4$  then the application of  $\sigma_\pi$  to the  $[3, 4]$ -associative law gives  $x_2x_1x_4x_3x_5 \dots x_{2n-1} \approx x_2x_1x_3x_5x_4x_6 \dots x_{2n-1} \in IdV$ .

Both identities together provide  $x_1 \dots x_{2n-1} \approx x_1 \dots x_{i-1} x_{i+1} x_{i+2} x_i x_{i+3} \dots x_{2n-1} \in IdV$  for  $1 \leq i \leq n-2$ . Let  $\rho \in S_n$  with  $\rho(2n-1) = 2n-2$ ,  $\rho(2n-2) = 2n-1$  and  $\rho(k) = k$  for  $1 \leq k \leq 2n-3$ . Dually, then the application of  $\sigma_\rho$  to the  $[1, n]$ -associative law as well as to the  $[n-3, n-2]$ -associative law (if  $n \geq 4$ ) provides identities from which we can derive  $x_1 \dots x_{2n-1} \approx x_1 \dots x_{i-1} x_{i+1} x_{i+2} x_i x_{i+3} \dots x_{2n-1} \in IdV$  for  $n \leq i \leq 2n-3$ . Finally, we have

$$\begin{aligned}
& x_1 \dots x_{2n-1} \\
& \approx x_1 \dots x_{n-1} x_{n+1} x_{n+2} x_n x_{n+3} \dots x_{2n-1} \\
& \approx x_1 \dots x_{n+1} x_{n-2} x_{n-1} x_{n+2} x_n x_{n+3} \dots x_{2n-1} \\
& \approx x_1 \dots x_{n+1} x_{n-2} x_n x_{n-1} x_{n+2} x_{n+3} \dots x_{2n-1} \\
& \approx x_1 \dots x_{n-2} x_n x_{n+1} x_{n-1} x_{n+2} x_{n+3} \dots x_{2n-1}, \text{ i.e.,} \\
& x_1 \dots x_{2n-1} \approx x_1 \dots x_{n-2} x_n x_{n+1} x_{n-1} x_{n+2} x_{n+3} \dots x_{2n-1} \in IdV.
\end{aligned}$$

Altogether we have  $x_1 \dots x_{2n-1} \approx x_1 \dots x_{i-1} x_{i+1} x_{i+2} x_i x_{i+3} \dots x_{2n-1} \in IdV$  for  $1 \leq i \leq 2n-3$ . ■

Now we will determine identities satisfying by presolid varieties.

**Lemma 4.** *Let  $4 \leq n \in 2\mathbb{N}$  and  $V$  be any  $Pre(n)$ -solid variety of  $n$ -semigroups. Then  $x_1 \dots x_{2n-1} \approx x_{\pi(1)} \dots x_{\pi(2n-1)}$  for all  $\pi \in S_{2n-1}$ .*

**Proof.** Let  $\pi \in S_{2n-1}$  with  $\pi(1) = 2$ ,  $\pi(2) = 1$  and  $\pi(k) = k$  for  $3 \leq k \leq 2n-1$ . If we apply  $\sigma_\pi$  to the  $[1, n]$ -associative law we get  $x_{n+1} x_2 x_1 x_3 \dots x_n x_{n+2} \dots x_{2n-1} \approx x_2 x_1 x_3 \dots x_{n+1} x_n x_{n+2} \dots x_{2n-1} \in IdV$  since  $V$  is  $Pre(n)$ -solid and by suitable substitution we obtain

$$(1) \quad x_1 \dots x_{2n-1} \approx x_2 \dots x_n x_1 x_{n+1} \dots x_{2n-1} \in IdV.$$

By Proposition 3 we have  $V \subseteq \tilde{V}_n$ . Using the identities of  $\tilde{V}_n$  we get  $x_2 \dots x_n x_1 x_{n+1} \dots x_{2n-1} \approx x_2 x_1 x_3 \dots x_{2n-1} \in IdV$  (since  $n$  is a even number). Together with (1) we obtain  $x_1 \dots x_{2n-1} \approx x_2 x_1 x_3 \dots x_{2n-1} \in IdV$ . It is easy to see that one can derive  $x_1 \dots x_{2n-1} \approx x_{\pi(1)} \dots x_{\pi(2n-1)}$  for all  $\pi \in S_{2n-1}$  from  $x_1 \dots x_{2n-1} \approx x_2 x_1 x_3 \dots x_{2n-1}$  and the identities of  $\tilde{V}_n$ . ■

**Lemma 5.** *Let  $3 \leq n \in \mathbb{N}$ ,  $2n - 1 \leq p \in (n - 1)\mathbb{N} + 1$  and  $V$  be a variety of  $n$ -semigroups with  $V \subseteq \widetilde{V}_n$ . Then for each  $\pi \in S_p$  holds*

$$x_{\pi(1)} \dots x_{\pi(p)} \approx x_1 \dots x_p \in IdV \text{ or}$$

$$x_{\pi(1)} \dots x_{\pi(p)} \approx x_2 x_1 x_3 \dots x_p \in IdV.$$

**Proof.** Let  $\pi \in S_p$ . We consider the term  $x_{\pi(1)} \dots x_{\pi(p)}$  and move step by step  $x_p, x_{p-1}, \dots, x_3$  to the  $p^{th}, (p-1)^{th}, \dots, 3^{th}$  position using the identities of  $\widetilde{V}_n$ . Then we have on the first both positions  $x_1 x_2$  or  $x_2 x_1$ . This shows  $x_{\pi(1)} \dots x_{\pi(p)} \approx x_1 \dots x_p \in IdV$  or  $x_{\pi(1)} \dots x_{\pi(p)} \approx x_2 x_1 x_3 \dots x_p \in IdV$ . ■

It is easy to check that  $Nper(n) \subseteq Pre(n)$ . So, any presolid variety has to be  $Nper(n)$ -solid. Next we find the lattice of all  $Nper(n)$ -solid varieties of  $n$ -semigroups.

**Lemma 6.** *Let  $3 \leq n \in \mathbb{N}$  and  $V$  be any variety of  $n$ -semigroups with  $V \subseteq \widetilde{V}_n$ . Then for each  $t \in \widetilde{W}_{(n)}^{np}(X)$  holds  $t \approx z^n \in IdV$ .*

**Proof.** Let  $t \in \widetilde{W}_{(n)}^{np}(X)$ . Then there is a variable  $w \in X$  that occurs at least two times in  $t$ . If  $l(t) = n$  then  $l(t) > cv(t)$  and  $t \approx x^n \in IdV$  since  $V \subseteq \widetilde{W}_n$ . Suppose now that  $l(t) > n$ . Using the identities of  $\widetilde{V}_n$  we can move  $w$  on the first and the second position, respectively, i.e.,  $t \approx w w u_3 \dots u_{l(t)}$  with  $u_3, \dots, u_{l(t)} \in X$ . Since  $x_1 x_1 x_3 \dots x_n \approx z^n \in IdV$  we have  $w w u_3 \dots u_{n-1} (u_n \dots u_{l(t)}) \approx z^n \in IdV$ , i.e.,  $t \approx z^n \in IdV$ . ■

**Lemma 7.** *Let  $3 \leq n \in \mathbb{N}$  and  $V$  be any nontrivial variety of  $n$ -semigroups with  $V \subseteq \widetilde{W}_n$ . Then only normal identities hold in  $V$ .*

**Proof.** Assume that a non-normal identity  $u \approx v$  holds in  $V$ . Then  $u \neq v$  and one of the terms  $u$  and  $v$  is a variable. Without loss of generality let  $u$  be a variable. Since  $V$  is a nontrivial variety the term  $v$  ( $\neq u$ ) is not a variable. Then by substitution we get  $y \approx y^{l(v)} \in IdV$  where  $l(v) > 1$ . Clearly,  $l(v) = r(n-1) + 1$  for some natural number  $r \geq 1$ . From  $xy^{n-1} \approx z^n \in IdV$  it follows  $y^{r(n-1)+1} \approx z^n \in IdV$ , i.e.,  $y^{l(v)} \approx z^n \in IdV$ . But  $y \approx y^{l(v)}$  and  $y^{l(v)} \approx z^n$  provide  $x \approx y$ , and  $V$  is the trivial variety, a contradiction. ■

**Proposition 8.** *Let  $3 \leq n \in \mathbb{N}$ . A nontrivial variety  $V$  of  $n$ -semigroups is  $Nper(n)$ -solid iff  $V \subseteq \widetilde{W}_n$ .*

**Proof.** Assume that  $V$  is  $Nper(n)$ -solid. We have  $t_1 := x_1x_2^{n-1} \in W_{(n)}^{np}(X_n)$ , i.e.,  $\sigma_{t_1} \in Nper(n)$  and its application to the  $[1, 3]$ -associative law gives

$$(1) \quad x_1x_2^{n-1}x_{n+1}^{n-1} \approx x_1x_2^{n-1} \in IdV.$$

Further, we have  $t_2 := x_2x_3^{n-1} \in W_{(n)}^{np}(X_n)$ , i.e.,  $\sigma_{t_2} \in Nper(n)$  and its application to the  $[1, 2]$ -associative law gives

$$(2) \quad x_{n+1}x_{n+2}^{n-1} \approx x_3x_4^{n-1}x_{n+2}^{n-1} \in IdV.$$

Then one obtains  $xy^{n-1} \stackrel{(1)}{\approx} xy^{n-1}z^{n-1} \stackrel{(2)}{\approx} wz^{n-1} \in IdV$ , i.e., we have  $xy^{n-1} \approx z^n \in IdV$ . Dually, we can show that  $x^{n-1}y \approx z^n \in IdV$ . Let now  $t \in W_{(n)}(X_n)$  with  $n = l(t) > cv(t)$ . Then there are  $u_1, \dots, u_n \in X$  such that  $t = u_1 \dots u_n$ . Since  $l(t) > cv(t)$  there are  $i, j \in \{1, \dots, n\}$  with  $i < j$  such that  $u_i = u_j$ . Then the term  $s := x_1 \dots x_{j-1}x_ix_{j+1} \dots x_n$  belongs to  $W_{(n)}^{np}(X_n)$ , i.e.,  $\sigma_s \in Nper(n)$ . Without loss of generality let  $i \neq 1$ . Then the application of  $\sigma_s$  to the  $[1, j]$ -associative law gives  $x_1 \dots x_{j-1}x_ix_{j+1} \dots x_n x_{n+1} \dots x_{n+j-2}x_{n+i-1}x_{n+j} \dots x_{2n-1} \approx x_1 \dots x_{j-1}x_ix_{n+j} \dots x_{2n-1}$ . Then  $x_{n+1} \notin \{x_1, \dots, x_{j-1}, x_i, x_{n+j}, \dots, x_{2n-1}\}$  since  $1 < i < j \neq 1$ . So, we substitute  $x_{n+1}$  by  $x_{n+1}^n$  and get  $x_1 \dots x_{j-1}x_ix_{n+j} \dots x_{2n-1} \approx x_1 \dots x_{j-1}x_ix_{j+1} \dots x_n x_{n+1}^n \dots x_{n+j-2}x_{n+i-1}x_{n+j} \dots x_{2n-1}$ . It is easy to check that one can derive  $x_1 \dots x_{j-1}x_ix_{j+1} \dots x_n x_{n+1}^n \dots x_{n+j-2}x_{n+i-1}x_{n+j} \dots x_{2n-1} \approx z^n$  using  $xy^{n-1} \approx x^{n-1}y \approx z^n \in IdV$ , i.e., one gets  $x_1 \dots x_{j-1}x_ix_{n+j} \dots x_{2n-1} \approx z^n \in IdV$ . Consequently, if we substitute  $x_i$  by  $u_i$  for  $1 \leq i \leq n$  we get  $u_1 \dots u_n \approx z^n \in IdV$ , i.e.,  $t \approx z^n \in IdV$ . Altogether,  $V \subseteq \widetilde{W}_n$ .

Suppose now that  $V \subseteq \widetilde{W}_n$ . Let  $t \in W_{(n)}^{np}(X_n)$ . Then  $t$  contains a subterm  $s$  with  $n = l(s) > cv(s)$  and there are words  $u$  and  $v$  (the empty word  $\lambda$  is also possible for  $u$  as well as for  $v$ ) such that  $t = usv$ . Since  $s \approx z^n \in IdV$  we have  $t \approx uz^n v \in IdV$ . The repeated application of  $xy^{n-1} \approx x^{n-1}y \approx z^n \in IdV$  to  $uz^n v$  gives finally  $uz^n v \approx z^n$ , i.e.,  $t \approx z^n \in IdV$ . This shows that any  $\sigma \in Nper(n)$  is  $V$ -equivalent to  $\sigma_{x_1^n}$ .

Let  $u \approx v \in IdV$ . If  $u = v$  then clearly  $\widehat{\sigma}_{x_1^n}[u] \approx \widehat{\sigma}_{x_1^n}[v] \in IdV$ . If  $u \neq v$  and  $u \approx v$  is a normal identity of  $V$  then there are natural numbers  $r, s \geq 1$  such that  $\widehat{\sigma}_{x_1^n}[u] \approx u_1^{n^r}$  and  $\widehat{\sigma}_{x_1^n}[v] \approx v_1^{n^s}$  where  $u_1$  ( $v_1$ ) is the first letter in  $u$  (in  $v$ ). From  $xy^{n-1} \approx z^n \in IdV$  it follows  $x^n \approx z^n \in IdV$  and thus  $u_1^{n^r} \approx v_1^{n^s} \in IdV$ , i.e.,  $\widehat{\sigma}_{x_1^n}[u] \approx \widehat{\sigma}_{x_1^n}[v] \in IdV$ . Since only normal identities are satisfied in  $V$  by Lemma 7 we can conclude that  $V$  is  $Nper(n)$ -solid. ■



After the following lemma we are able to characterize all presolid varieties of  $n$ -semigroups.

**Lemma 9.** *Let  $3 \leq n \in 2\mathbb{N} + 1$ ,  $V$  be a variety of  $n$ -semigroups with  $V \subseteq \tilde{V}_n$ , and  $\sigma \in \text{Per}(n)$ . Then there holds*

$$\hat{\sigma}[x_1 \dots x_i (x_{i+1} \dots x_{i+n}) x_{i+n+1} \dots x_{2n-1}] \approx x_1 \dots x_{2n-1} \in \text{Id}V$$

for  $0 \leq i \leq n - 1$ .

**Proof.** Let  $\pi \in S_n$ . Without loss of generality let  $i = 0$ . Then

- (1)  $x_{\pi(1)} \dots x_{\pi(n)} x_{n+1} \dots x_{2n-1} \approx x_1 \dots x_{2n-1} \in \text{Id}V$  or  
 (2)  $x_{\pi(1)} \dots x_{\pi(n)} x_{n+1} \dots x_{2n-1} \approx x_2 x_1 x_3 \dots x_{2n-1} \in \text{Id}V$  by Lemma 5.  
 We put  $y_1 := x_1 \dots x_n$  in case (1) ( $y_1 := x_2 x_1 x_3 \dots x_n$  in case (2)) and  $y_j := x_{n+j-1}$  for  $2 \leq j \leq n$ . Using the identities of  $\tilde{V}_n$  it is easy to check that  $y_{\pi(1)} \dots y_{\pi(n)} \approx x_1 \dots x_{2n-1} \in \text{Id}V$  in case (1) and  $y_{\pi(1)} \dots y_{\pi(n)} \approx x_{n+1} x_2 x_1 x_3 \dots x_n x_{n+2} \dots x_{2n-1} \in \text{Id}V$  in case (2), respectively. Further, we have  $x_{n+1} x_2 x_1 x_3 \dots x_n x_{n+2} \dots x_{2n-1} \approx x_1 x_{n+1} x_2 x_3 \dots x_n x_{n+2} \dots x_{2n-1} \approx x_1 x_2 x_3 \dots x_n x_{n+1} x_{n+2} \dots x_{2n-1} \in \text{Id}V$  (since  $n$  is an odd number). This shows that  $\hat{\sigma}_\pi[(x_1 \dots x_n) x_{n+1} \dots x_{2n-1}] \approx S_{2n-1}^n(\sigma_\pi(f), S_{2n-1}^n(\sigma_\pi(f), x_1, \dots, x_n), x_{n+1}, \dots, x_{2n-1}) \approx x_1 \dots x_{2n-1} \in \text{Id}V$ . ■

**Theorem 10.** *Let  $n \geq 3$  be a natural number and  $V$  be a nontrivial variety of  $n$ -semigroups. Then  $V$  is  $\text{Pre}(n)$ -solid iff the following statements hold:*

- (i)  $V \subseteq V_n$ ;
- (ii) If  $x_{\pi(1)} \dots x_{\pi(n)} \approx x_1 \dots x_n \in \text{Id}V$  for some  $\pi \in S_n$  then  $x_{\pi \circ s(1)} \dots x_{\pi \circ s(n)} \approx x_{s(1)} \dots x_{s(n)} \in \text{Id}V$  for all  $s \in S_n$ ;
- (iii) If  $n \in 2\mathbb{N}$  then  $x_1 \dots x_{2n-1} \approx x_{\pi(1)} \dots x_{\pi(2n-1)}$  for all  $\pi \in S_{2n-1}$ .

**Proof.** Suppose that  $V$  is  $\text{Pre}(n)$ -solid. Then  $V \subseteq \tilde{V}_n$  by Proposition 3. Further,  $V$  is  $N\text{per}(n)$ -solid since  $N\text{per}(n) \subseteq \text{Pre}(n)$ . Then by Proposition 8 we get  $V \subseteq \tilde{W}_n$ . Therefore,  $V \subseteq \tilde{V}_n \cap \tilde{W}_n = V_n$  and it holds (i). Suppose that  $x_{\pi(1)} \dots x_{\pi(n)} \approx x_1 \dots x_n \in \text{Id}V$  for some  $\pi \in S_n$ . Further let  $\rho \in S_n$ . Then  $\sigma_\rho \in \text{Pre}(n)$ . Since  $V$  is  $\text{Pre}(n)$ -solid we have  $\hat{\sigma}_\rho[x_1 \dots x_n] \approx \hat{\sigma}_\rho[x_{\pi(1)} \dots x_{\pi(n)}] \in \text{Id}V$ , i.e.,  $x_{\pi \circ \rho(1)} \dots x_{\pi \circ \rho(n)} \approx x_{\rho(1)} \dots x_{\rho(n)} \in \text{Id}V$ . This shows (ii). Finally, (iii) it follows from Lemma 4.

Suppose that (i)–(iii) are satisfied. Let  $\sigma_t \in \text{Pre}(n)$ . If  $\sigma_t \notin \text{Per}(n)$  then  $t \in \widetilde{W}_{(n)}^{np}(X)$  and  $t \approx z^n \in \text{IdV}$  by Lemma 6, i.e.,  $\sigma_t$  is  $V$ -equivalent to  $\sigma_{x_1^n}$ , where  $\sigma_{x_1^n} \in \text{Nper}(n)$ . But (i) implies that  $V$  is  $\text{Nper}(n)$ -solid by Proposition 8. Thus  $\widehat{\sigma}_{x_1^n}[u] \approx \widehat{\sigma}_{x_1^n}[v] \in \text{IdV}$  for all  $u \approx v \in \text{IdV}$ , i.e.,  $\widehat{\sigma}_t[u] \approx \widehat{\sigma}_t[v] \in \text{IdV}$  for all  $u \approx v \in \text{IdV}$ . Let now  $\sigma_t \in \text{Per}(n)$  and  $u \approx v \in \text{IdV}$ . If  $\text{var}(u) \neq \text{var}(v)$  then without loss of generality there is a  $w \in \text{var}(u) \setminus \text{var}(v)$ . We substitute  $w$  by  $x^n$  and get  $\tilde{u} \approx v \in \text{IdV}$  from  $u \approx v \in \text{IdV}$  where  $x^n$  is a subterm of  $\tilde{u}$ , i.e.,  $\tilde{u} \in \widetilde{W}_{(n)}^{np}(X)$ . Then by Lemma 6 we have  $\tilde{u} \approx x^n \in \text{IdV}$ , i.e.,  $u \approx v \approx x^n \in \text{IdV}$ . If  $l(u) > cv(u)$  or  $l(v) > cv(v)$  then  $u \in \widetilde{W}_{(n)}^{np}(X)$  or  $v \in \widetilde{W}_{(n)}^{np}(X)$  and thus  $u \approx v \approx x^n \in \text{IdV}$  by Lemma 6. Consequently, if  $\text{var}(u) \neq \text{var}(v)$  or  $l(u) > cv(u)$  or  $l(v) > cv(v)$  then  $u \approx v \approx x^n \in \text{IdV}$ . If, in particular,  $l(u) = cv(u)$  then  $u = u_1 \dots u_{l(u)}$  with  $u_1, \dots, u_{l(u)} \in X$  and there is a  $\pi \in S_{l(u)}$  such that  $\widehat{\sigma}_t[u] \approx u_{\pi(1)} \dots u_{\pi(l(u))}$ . But from  $u \approx x^n \in \text{IdV}$  we get by the substitution  $u_i \mapsto u_{\pi(i)}$  for  $1 \leq i \leq l(u)$  that  $u_{\pi(1)} \dots u_{\pi(l(u))} \approx x^n \in \text{IdV}$ , i.e.,  $\widehat{\sigma}_t[u] \approx x^n \in \text{IdV}$ . If, in particular,  $l(v) = cv(v)$  then we get  $\widehat{\sigma}_t[v] \approx x^n \in \text{IdV}$  in the same matter. If  $l(u) > cv(u)$  ( $l(v) > cv(v)$ ) then  $u \in \widetilde{W}_{(n)}^{np}(X)$  ( $v \in \widetilde{W}_{(n)}^{np}(X)$ ) and it is easy to check that  $\widehat{\sigma}_t[u] \in \widetilde{W}_{(n)}^{np}(X)$  ( $\widehat{\sigma}_t[v] \in \widetilde{W}_{(n)}^{np}(X)$ ), too. Then  $\widehat{\sigma}_t[u] \approx x^n \in \text{IdV}$  ( $\widehat{\sigma}_t[v] \approx x^n \in \text{IdV}$ ) by Lemma 6. Consequently,  $\widehat{\sigma}_t[u] \approx x^n \approx \widehat{\sigma}_t[v] \in \text{IdV}$ . The remaining case is  $\text{var}(u) = \text{var}(v)$  and  $l(u) = cv(u)$  and  $l(v) = cv(v)$ . We put  $s := l(u)$  and  $\{u_1, \dots, u_s\} = \text{var}(u) = \text{var}(v)$ . Because of Lemma 9 (if  $n \in 2\mathbb{N} + 1$ ) and of (iii) (if  $n \in 2\mathbb{N}$ ), respectively, we have  $\widehat{\sigma}_t[x_1 \dots x_{i-1}(x_i \dots x_{i+n-1})x_{i+n} \dots x_{2n-1}] \approx \widehat{\sigma}_t[x_1 \dots x_{j-1}(x_j \dots x_{j+n-1})x_{j+n} \dots x_{2n-1}] \in \text{IdV}$  for  $1 \leq i < j \leq n$ . Therefore we can assume that

$$u = (\dots (u_1 \dots u_n)u_{n+1} \dots u_{2n-1}) \dots u_{s-1}u_s)$$

$$v = (\dots (u_{\pi(1)} \dots u_{\pi(n)})u_{\pi(n+1)} \dots u_{\pi(2n-1)}) \dots u_{\pi(s-1)}u_{\pi(s)})$$

for some permutation  $\pi \in S_s$ . Further there is a  $\rho \in S_n$  such that  $\sigma_t = \sigma_\rho$ . If  $s = 1$  we have obviously  $\widehat{\sigma}_\rho[u] \approx \widehat{\sigma}_\rho[v] \in \text{IdV}$ . If  $s = n$  then  $\widehat{\sigma}_\rho[u] \approx u_{\rho(1)} \dots u_{\rho(n)}$  and  $\widehat{\sigma}_\rho[v] \approx u_{\pi \circ \rho(1)} \dots u_{\pi \circ \rho(n)}$ . By (ii) from  $x_{\pi(1)} \dots x_{\pi(n)} \approx x_1 \dots x_n \in \text{IdV}$  it follows  $x_{\pi \circ \rho(1)} \dots x_{\pi \circ \rho(n)} \approx x_{\rho(1)} \dots x_{\rho(n)} \in \text{IdV}$ , i.e.,  $\widehat{\sigma}_\rho[u] \approx \widehat{\sigma}_\rho[v] \in \text{IdV}$ . Let now  $s > n$ . Then there is a  $\phi \in S_s$  such that  $\widehat{\sigma}_t[u] \approx u_{\phi(1)} \dots u_{\phi(s)}$  and  $\widehat{\sigma}_t[v] \approx u_{\pi \circ \phi(1)} \dots u_{\pi \circ \phi(s)}$ .

By Lemma 5 we have  $\hat{\sigma}_t[u] \approx u_1 \dots u_s$  or  $\hat{\sigma}_t[u] \approx u_2 u_1 u_3 \dots u_s =: \tilde{u}$ . If  $\hat{\sigma}_t[u] \approx u$ , i.e.,  $x_{\phi(1)} \dots x_{\phi(s)} \approx u_1 \dots u_s \in IdV$  then by the substitution  $u_i \mapsto u_{\pi(i)}$  for  $1 \leq i \leq s$  we get  $u_{\pi \circ \phi(1)} \dots u_{\pi \circ \phi(s)} \approx u_{\pi(1)} \dots u_{\pi(s)} \in IdV$ , i.e.,  $\hat{\sigma}_t[v] \approx v$ , and from  $u \approx v \in IdV$  it follows  $\hat{\sigma}_t[u] \approx \hat{\sigma}_t[v] \in IdV$ . If  $\hat{\sigma}_t[u] \approx \tilde{u}$ , i.e.,  $u_{\phi(1)} \dots u_{\phi(s)} \approx u_2 u_1 u_3 \dots u_s$  then by the same substitution we get  $u_{\pi \circ \phi(1)} \dots u_{\pi \circ \phi(s)} \approx u_{\pi(2)} u_{\pi(1)} u_{\pi(3)} \dots u_{\pi(s)} =: \tilde{v}$ , i.e.,  $\hat{\sigma}_t[v] \approx \tilde{v} \in IdV$ . Moreover, from Lemma 5 we get

$$u_{\pi(2)} u_{\pi(1)} u_{\pi(3)} \dots u_{\pi(s)} \approx u_1 \dots u_s \quad \text{or}$$

$$u_{\pi(2)} u_{\pi(1)} u_{\pi(3)} \dots u_{\pi(s)} \approx u_2 u_1 u_3 \dots u_s$$

as well as

$$u_{\pi^{-1}(2)} u_{\pi^{-1}(1)} u_{\pi^{-1}(3)} \dots u_{\pi^{-1}(s)} \approx u_1 \dots u_s \quad \text{or}$$

$$u_{\pi^{-1}(2)} u_{\pi^{-1}(1)} u_{\pi^{-1}(3)} \dots u_{\pi^{-1}(s)} \approx u_2 u_1 u_3 \dots u_s.$$

i.e.,

$$u_2 u_1 u_3 \dots u_s \approx u_{\pi(1)} \dots u_{\pi(s)} \quad \text{or}$$

$$u_2 u_1 u_3 \dots u_s \approx u_{\pi(2)} u_{\pi(1)} u_{\pi(3)} \dots u_{\pi(s)}.$$

This shows  $\tilde{v} \approx u$  or  $\tilde{v} \approx \tilde{u}$  as well as  $\tilde{u} \approx v$  or  $\tilde{u} \approx \tilde{v}$ . This implies  $\tilde{v} \approx \tilde{u}$  or both  $\tilde{v} \approx u$  and  $\tilde{u} \approx v$  hold in  $V$ . Since  $u \approx v \in IdV$  we have altogether  $\tilde{v} \approx \tilde{u} \in IdV$  and thus  $\hat{\sigma}_t[u] \approx \hat{\sigma}_t[v] \in IdV$  because of  $\hat{\sigma}_t[u] \approx \tilde{u} \in IdV$  and  $\hat{\sigma}_t[v] \approx \tilde{v} \in IdV$ .  $\blacksquare$

Let us apply Theorem 10 for the case  $n = 3$ . We obtain the following characterization of all presolid varieties of 3-semigroups.

**Corollary 11.** *A nontrivial variety of 3-semigroups is Pre(3)-solid iff  $V \subseteq Mod\{(xyz)wt \approx x(yzw)t \approx xy(zwt) \approx yzxwt \approx xzwy t \approx xywtz, xyx \approx x^2y \approx xy^2 \approx z^3\} =: W$  and it holds the following condition:*

(\*) *If  $x_1 x_2 x_3 \approx x_{\pi(1)} x_{\pi(2)} x_{\pi(3)} \in IdV$  for some  $\pi \in \{(12), (13), (23)\}$*

*then  $x_1 x_2 x_3 \approx x_{\rho(1)} x_{\rho(2)} x_{\rho(3)} \in IdV$  for all  $\rho \in S_3$ .*

**Proof.** Suppose that  $V$  is  $Pre(3)$ -solid. Then the conditions (i) and (ii) of Theorem 10 are satisfied. From (i) it follows that  $xyzwt \approx yzxtw \approx xzwy \approx xywtz \in IdV$  and  $xyx \approx x^2y \approx xy^2 \approx z^3 \in IdV$ . Hence  $V \subseteq W$ . Using (ii) we can verify condition (\*): If  $\pi = (13)$ , i.e.,  $x_1x_2x_3 \approx x_3x_2x_1 \in IdV$  then  $x_2x_1x_3 \approx x_2x_3x_1 \in IdV$  (for  $s = (12)$ ). Both identities provide  $x_1x_2x_3 \approx x_1x_3x_2 \approx x_2x_3x_1 \approx x_2x_1x_3 \approx x_2x_3x_1 \approx x_1x_3x_2 \in IdV$ . If  $\pi = (12)$ , i.e.,  $x_1x_2x_3 \approx x_2x_1x_3 \in IdV$  then  $x_1x_3x_2 \approx x_2x_3x_1 \in IdV$  (for  $s = (23)$ ). If  $\pi = (23)$ , i.e.,  $x_1x_2x_3 \approx x_1x_3x_2 \in IdV$  then  $x_2x_1x_3 \approx x_3x_1x_2 \in IdV$  (for  $s = (12)$ ). In the latter two cases, we conclude in the same matter as before.

Suppose now that  $V \subseteq W$  and (\*) is satisfied. Since  $V \subseteq W$ , the condition (i) of Theorem 10 holds. We have now to show that also condition (ii) is satisfied. For this let  $\pi \in S_3$ . If  $\pi \in \{(1), (12), (13), (23)\}$  then the condition is satisfied by (\*). If  $\pi = (123)$ , i.e.,  $x_1x_2x_3 \approx x_2x_3x_1 \in IdV$  then we have to check that also  $x_2x_1x_3 \approx x_3x_2x_1 \in IdV$ ,  $x_3x_2x_1 \approx x_1x_3x_2 \in IdV$ ,  $x_1x_3x_2 \approx x_2x_1x_3 \in IdV$ ,  $x_2x_3x_1 \approx x_3x_1x_2 \in IdV$ , and  $x_3x_1x_2 \approx x_1x_2x_3 \in IdV$ . Obviously, these five equations are consequences of the given identity  $x_1x_2x_3 \approx x_2x_3x_1 \in IdV$ . If  $\pi = (132)$  the we conclude in the same matter. This shows (ii). Condition (iii) can be neglected since 3 is odd. Altogether,  $V$  is  $Pre(3)$ -solid by Theorem 10. ■

#### REFERENCES

- [1] V. Budd, K. Denecke and S.L. Wismath, *Short-solid superassociative type  $(n)$  varieties*, East-West J. of Mathematics **3** (2) (2001), 129–145.
- [2] W. Dörnte, *Untersuchungen über einen verallgemeinerten Gruppenbegriff*, Math. Z. **29** (1928), 1–19.
- [3] K. Denecke and Hounnon, *All solid varieties of semirings*, Journal of Algebra **248** (2002), 107–117.
- [4] K. Denecke and J. Koppitz, *Pre-solid varieties of semigroups*, Archivum Mathematicum **31** (1995), 171–181.
- [5] K. Denecke and J. Koppitz, *Finite monoids of hypersubstitutions of type  $\tau = (2)$* , Semigroup Forum **56** (1998), 265–275.
- [6] K. Denecke and M. Reichel, *Monoids of hypersubstitutions and  $M$ -solid varieties*, Contributions to General Algebra **9** (1995), 117–126.
- [7] K. Denecke, J. Koppitz and S.L. Wismath, *Solid varieties of arbitrary type*, Algebra Universalis **48** (2002), 357–378.

- [8] K. Denecke and S.L. Wismath, *Hyperidentities and clones*, Gordon and Breach Scientific Publisher, 2000.
- [9] J. Koppitz, *Hypersubstitutions and groups*, Novi Sad J. Math. **34** (2) (2004), 127–139.
- [10] L. Polák, *All solid varieties of semigroups*, Journal of Algebra **219** (1999), 421–436.
- [11] J. Płonka, *Proper and inner hypersubstitutions of varieties*, p. 106–115 in: “*Proceedings of the International Conference: ‘Summer School on General Algebra and Ordered Sets’, Olomouc 1994*”, Palacký University, Olomouc 1994.

Received 15 July 2005