

BOGUSŁAWA BEDNAREK-KOZEK and A. KOZEK (Wrocław)

ON THE ROBUSTNESS OF PROPERTIES CHARACTERIZING THE NORMAL DISTRIBUTION

1. Introduction. In 1951 Sapogov [9] proved the robustness for Cramer's theorem. Certain types of extensions of this result were investigated in [4], [10]-[13].

In 1968 Hoang Huu Nhu [1] showed that, with some supplementary assumptions,

1. ε -independence of the random variables $X_1 + X_2$ and $X_1 - X_2$, where X_1 and X_2 are independent, implies $c/\sqrt{\ln 1/\varepsilon}$ -normality of X_1 and X_2 ;

2. ε -independence of

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \quad \text{and} \quad S^2 = \sum_{i=1}^n (X_i - \bar{X})^2,$$

where X_i are independent with the same distribution function F , implies $c/\sqrt{\ln 1/\varepsilon}$ -normality of X_i ($i = 1, \dots, n$).

The purpose of this paper is to prove analogous theorems:

1. for some ε -independent linear and quadratic statistics without the assumption that the random variables have finite moments,
2. for some quadratic statistics which have ε -polynomial regression on the sum of the random variables with finite variance.

We also obtain similar results in the case of Lévy's metric.

2. Notation and definitions. Let us denote by $F_X(x)$ the distribution function and by $f_X(t)$ the characteristic function of the random variable X . Further, let $E X$ and $\text{Var } X$ stand for the *mean* and *variance* of X , respectively. Moreover, let

$$M_\Theta X = E |X|^{2(1+\Theta)} \quad \text{for some } \Theta \in (0, \frac{1}{2}].$$

Let us write

$$(1) \quad \rho_{XY}(x, y) = G_{XY}(x, y) - F_X(x)F_Y(y),$$

where $G_{XY}(x, y)$ is the joint distribution function of (X, Y) , $q(\varepsilon, \alpha_0, f) = \sup\{c: |f_X(t)| > \varepsilon^{\alpha_0} \text{ for all } t \in (-c, c)\}$, with $1 > \varepsilon \geq 0$, $\alpha_0 > 0$ and let

$$(2) \quad h_\gamma(x) = \begin{cases} -\gamma & \text{if } x < -\gamma, \\ x & \text{if } -\gamma \leq x \leq \gamma, \\ \gamma & \text{if } x > \gamma, \end{cases}$$

where γ is a positive constant.

The definitions 1 and 2 may be found, for example, in [1]. However, for the sake of completeness, they should be included here.

Definition 1. A distribution function F is ε -normal if there exist m and $\sigma > 0$ such that

$$\sup_x \left| F(x) - \Phi\left(\frac{x-m}{\sigma}\right) \right| \leq \varepsilon,$$

where

$$\Phi(x) = (2\pi)^{-1/2} \int_{-\infty}^x \exp\left(-\frac{x^2}{2}\right) dx.$$

Definition 2. The random variables X and Y are said to be ε -independent if for every x and y

$$|\varphi_{XY}(x, y)| \leq \varepsilon,$$

where $\varphi_{XY}(x, y)$ is defined by (1).

Furthermore, let us introduce the following definitions.

Definition 3. We say that the random variable Y has ε -polynomial regression of order k on the random variable X if there exists a function $\eta(x)$ such that

$$\mathbf{E}(Y|X = x) = \sum_{j=1}^k \beta_j x^j + \eta(x),$$

where $\beta_k \neq 0$ and $\sup_x |\eta(x)| \leq \varepsilon$.

Definition 4. We say that the random variable X belongs to the family $F(M, \Theta; \varepsilon)$ if there exists $\gamma > 0$ such that γ and $-\gamma$ are the points of continuity of $F_X(x)$ and the following conditions hold:

$$\begin{aligned} F_X(-\gamma) &\leq \varepsilon, & F_X(\gamma) &\geq 1 - \varepsilon, \\ M^{-1} &\leq \text{Var} h_\gamma(X) \leq M, & M_\Theta(h_\gamma(X)) &\leq M, \end{aligned}$$

where M and Θ are constants, $M > 1$, $0 < \Theta \leq \frac{1}{2}$.

The constants C_i appearing in the theorems below depend on M , Θ , n but not on ε . This is the reason for considering the families $F(M, \Theta; \varepsilon)$ with M and Θ fixed.

3. Theorems. Let X_1, X_2 be random variables and let

$$(3) \quad S_1 = a_{11}X_1 + a_{12}X_2, \quad S_2 = a_{21}X_1 + a_{22}X_2$$

where

$$(4) \quad -\infty < \frac{a_{11}a_{22}}{a_{12}a_{21}} < 0.$$

THEOREM 1. Let $X_1, X_2 \in F(M, \Theta, \varepsilon)$ be independent random variables such that S_1 and S_2 given by (3) are ε -independent. If ε is sufficiently small, then the random variables X_1, X_2 are $\delta_i(\varepsilon)$ -normally distributed, where

$$(5) \quad \delta_i(\varepsilon) = \frac{C_1}{\sqrt{\ln 1/\varepsilon}}.$$

Remark. If X_1, X_2 are non-degenerate and independent random variables, then (4) is a necessary condition for the independence of S_1 and S_2 .

For the random variables X_1, \dots, X_n let us write

$$(6) \quad S = \sum_{j=1}^n X_j,$$

$$(7) \quad S_3 = \sum_{\substack{j=1 \\ k=1}}^n a_{jk} X_j X_k + \sum_{j=1}^n b_j X_j,$$

where

$$(8) \quad \sum_{j=1}^n a_{jj} \neq 0$$

and

$$(9) \quad \sum_{j,k=1}^n a_{jk} = 0, \quad \sum_{j=1}^n b_j = 0.$$

THEOREM 2. Let $X_1, \dots, X_n \in F(M, \Theta, \varepsilon)$ be independent and identically distributed random variables such that S and S_3 given by (6) and (7) are ε -independent. If ε is sufficiently small, then the random variables X_1, \dots, X_n are $\delta(\varepsilon)$ -normally distributed, where

$$(10) \quad \delta(\varepsilon) = \frac{C_2}{\sqrt{\ln 1/\varepsilon}}.$$

Remark. If non-degenerate independent random variables X_1, \dots, X_n are identically normally distributed, then by virtue of the theorem of Laha [3] $S(X_1, \dots, X_n)$ and $S_3(X_1, \dots, X_n)$ are independent if and only

if the statistics $S_3(X_1 - \lambda, \dots, X_n - \lambda)$ and $S_3(X_1, \dots, X_n)$ are identically distributed for each $\lambda \in R$. The assumptions (9) are the necessary condition for this case.

Let us remark that if the random variables U and V are ε -independent, then the random variables U and V^2 are 2ε -independent. Hence and from Theorem 2 we obtain

COROLLARY TO THEOREM 2. *Let*

$$S_4 = \sum_{j=1}^n a_j X_j,$$

where

$$\sum_{j=1}^n a_j^2 \neq 0, \quad \sum_{j=1}^n a_j = 0.$$

Let $X_1, \dots, X_n \in F(M, \Theta, 2\varepsilon)$ be independent and identically distributed random variables such that $S = \sum_{j=1}^n X_j$ and S_4 are ε -independent.

If ε is sufficiently small, then the random variables X_1, \dots, X_n are $\delta(\varepsilon)$ -normally distributed, where $\delta(\varepsilon)$ is given by (10) with the constant C_3 instead of C_2 .

THEOREM 3. *Let X_1, \dots, X_n be independent and identically distributed random variables with finite variance and let*

$$S_5 = \sum_{j,k=1}^n a_{jk} X_j X_k + \sum_{j=1}^n b_j X_j.$$

If the statistic S_5 has ε -polynomial regression of order ≤ 2 on S ($S = \sum_{j=1}^n X_j$) and if the regression coefficients satisfy conditions

$$(11) \quad n\beta_2 \neq \sum_{j=1}^n a_{jj}, \quad n^2\beta_2 = \sum_{j,k=1}^n a_{jk}, \quad n\beta_1 = \sum_{j=1}^n b_j,$$

$$\frac{\beta_0}{\sum_{j=1}^n a_{jj} - n\beta_2} > 0,$$

then, for ε sufficiently small, there exists a constant C_4 such that the random variables X_1, \dots, X_n are $C_4/\sqrt{\ln 1/\varepsilon}$ -normally distributed.

Remark. Lukacs and Laha [6] proved the following theorem:

Under the assumption that X_1, \dots, X_n are independently and identically distributed random variables, they are non-degenerate and normally distributed if and only if the statistic S_5 has polynomial regression of order ≤ 2 on S and the regression coefficients satisfy conditions (11).

4. Lemmas. In order to prove the theorems from Section 3 we use the well-known method [7] consisting in deduction of certain differential equations for characteristic functions and the method due to Hoang Huu Nhu [1] which allows us to use Esseen's theorem ([5], 20.3A).

To prove the theorems of Section 3 we need the following lemmas.

LEMMA 1. Let X_1, \dots, X_n be independently distributed random variables and let γ_i and $-\gamma_i$ ($\gamma_i > 0, i = 1, \dots, n$) be the points of continuity of $F_{X_i}(x)$ ($i = 1, \dots, n$) such that

$$F_{X_i}(-\gamma_i) \leq \varepsilon \quad \text{and} \quad F_{X_i}(\gamma_i) \geq 1 - \varepsilon.$$

If the statistics $T_1(X_1, \dots, X_n)$ and $T_2(X_1, \dots, X_n)$ are ε -independent and if

$$(12) \quad Y_i = h_{\gamma_i}(X_i) \quad (i = 1, \dots, n),$$

then

- I. Y_i ($i = 1, \dots, n$) are independent random variables,
- II. Y_i ($i = 1, \dots, n$) have finite moments of all orders,
- III. $\sup_x |F_{X_i}(x) - F_{Y_i}(x)| \leq \varepsilon$ ($i = 1, \dots, n$),

IV. the statistics $T_1(Y_1, \dots, Y_n)$ and $T_2(Y_1, \dots, Y_n)$ are $(6n + 1)\varepsilon$ -independent.

Proof. Properties I, II and III of the variables Y_i ($i = 1, \dots, n$) follow immediately from (12) and from the definition (2) of the function h_{γ_i} . Thus, in order to prove Lemma 1, it is enough to prove property IV.

Let

$$A = \{(a_1, \dots, a_n); -\gamma_i < a_i < \gamma_i, i = 1, \dots, n\}, \quad A' = R^n \setminus A,$$

$$B = B_x = T_1^{-1}[(-\infty, x)], \quad D = D_y = T_2^{-1}[(-\infty, y)],$$

where $T_i^{-1}[W]$ ($i = 1, 2$) denote the inverse image of the set $W \subset R$.

Let P_X and P_Y be the joint distributions of the variables X_1, \dots, X_n and Y_1, \dots, Y_n , respectively.

From the definition (2) of h_{γ_i} , we have

$$P_X(A') \leq 2n\varepsilon, \quad P_Y(A') \leq 2n\varepsilon.$$

Moreover, by virtue of the definitions of P_X and P_Y , for each Borelian set $W \subset R^n$ we have

$$P_Y(W) = P_Y(W \cap A) + P_Y(W \cap A'),$$

$$P_Y(W \cap A) = P_X(W) - P_X(W \cap A').$$

Hence we obtain

$$|\varphi_{T_1 T_2}(x, y)| = |P_Y(B \cap D) - P_Y(B)P_Y(D)| \leq (6n + 1)\varepsilon.$$

Thus the statistics T_1 and T_2 are $(6n+1)\varepsilon$ -independent, which completes the proof of Lemma 1.

LEMMA 2. *If the function $f(t)$ can be expressed by a formula*

$$f(t) = \exp \left\{ -\frac{1}{2} \sigma^2 t^2 + imt + \int_0^t \int_0^\tau K(u) du d\tau \right\}$$

for $t \in [-q, q]$, where $q = q(\varepsilon, \alpha_0, f)$, $\varepsilon > 0$, $\alpha_0 > 0$ and $|K(t)|$ is a function bounded by c for $t \in [-q, q]$, then

$$q \geq \sqrt{\frac{2\alpha_0}{\sigma^2 + c}} \sqrt{\ln \frac{1}{\varepsilon}}.$$

LEMMA 3. *Let U, V be random variables $a\varepsilon$ -independent and let $\varphi(u, v) = \varphi_{UV}(u, v)$ be the function defined by (1). If*

$$\begin{aligned} J_1 &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} v e^{itu} d\varphi(u, v), \\ J_2 &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} iuv e^{itu} d\varphi(u, v), \\ J_3 &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} v^2 e^{itu} d\varphi(u, v), \end{aligned}$$

then for each $N \geq 1$ and for each t

$$(13) \quad |J_1| \leq 8Na\varepsilon(1 + N|t|) + 2N^{-\theta}(M_\theta U + 2E|V|^{1+\theta}),$$

$$(14) \quad |J_2| \leq 8Na\varepsilon(1 + N|t|) + 2N^{-\theta}(M_\theta U + 2M_\theta V),$$

$$(15) \quad |J_2| \leq 8N^2a\varepsilon(2 + N|t|) + 2N^{-\theta}(M_\theta U + M_\theta V + \sqrt{(M_\theta U)(M_\theta V)}),$$

$$(16) \quad |J_3| \leq 8N^2a\varepsilon(1 + N|t|) + 2N^{-\theta}(M_\theta U + 2M_\theta V).$$

Integrating by parts we obtain after some calculations inequalities (13)-(16) (see [1], [8]).

LEMMA 4. *If Y_1 and Y_2 are independently distributed random variables, then*

$$(17) \quad M_\theta(Y_1 + \alpha Y_2) \leq 2(1 + |\alpha|)^3 \max_{j=1,2} (M_\theta Y_j).$$

If Y_1, \dots, Y_n are independent and identically distributed random variables and if

$$\max \{ |a_{11}|, |a_{12}|, \dots, |a_{nn-1}|, |a_{nn}|, |b_1|, \dots, |b_n| \} \leq 1,$$

then

$$(18) \quad \mathbf{E} \left| \sum_{j,k=1}^n a_{jk} Y_j Y_k + \sum_{j=1}^n b_j Y_j \right|^{1+\theta} \leq 4n^4 (M_\theta Y_1 + 1)$$

and

$$(19) \quad M_\theta \left(\sum_{j=1}^n Y_j \right) \leq n^4 M_\theta Y_1.$$

Proof. We shall prove (18). One can prove the remaining inequalities in an analogous way.

Let us remark that

$$\begin{aligned} \mathbf{E} \left| \sum_{j,k=1}^n a_{jk} Y_j Y_k + \sum_{j=1}^n b_j Y_j \right|^{1+\theta} &\leq n^3 \mathbf{E} \left| \max_{j=1,\dots,n} (|Y_j|^2) + \max_{j=1,\dots,n} (|Y_j|) \right|^{1+\theta} \\ &\leq 4n^3 \mathbf{E} (\max \{ |Y_1|^{2(1+\theta)}, \dots, |Y_n|^{2(1+\theta)}, |Y_1|^{1+\theta}, \dots, |Y_n|^{1+\theta} \}) \\ &\leq 4n^3 \mathbf{E} (\max \{ |Y_1|^{2(1+\theta)}, \dots, |Y_n|^{2(1+\theta)}, 1 \}) \leq 4n^4 (M_\theta Y_1 + 1), \end{aligned}$$

which completes the proof of Lemma 4.

LEMMA 5. *If $f(t)$ is the characteristic function of the distribution function $F(x)$ and if for each $t \in [-Q, Q]$*

$$(20) \quad f(t) = \exp \left\{ -\frac{1}{2} \sigma^2 t^2 + imt + \int_0^t \int_0^\tau K(u) du d\tau \right\}$$

and the function $K(t)$ satisfies the condition

$$(21) \quad |K(t)| \leq c < \sigma^2 \quad \text{for each } t \in [-Q, Q],$$

then

$$\sup_x \left| F(x) - \Phi \left(\frac{x-m}{\sigma} \right) \right| \leq \frac{1}{\pi} \frac{c}{\sigma^2 - c} + \frac{24}{\pi} \frac{1}{\sqrt{2\pi} \sigma Q}.$$

Proof. Let

$$L = L(t) = |f(t) - \exp(-\frac{1}{2}t^2\sigma^2 + imt)| \quad \text{for each } t \in [-Q, Q].$$

By the inequality $|e^z - 1| \leq |z| \exp|z|$ we obtain

$$\begin{aligned} L &= \exp(-\frac{1}{2}t^2\sigma^2) \left| \exp \left(\int_0^t \int_0^\tau K(u) du d\tau \right) - 1 \right| \\ &\leq \left| \int_0^t \int_0^\tau K(u) du d\tau \right| \exp \left(-\frac{1}{2}t^2\sigma^2 + \left| \int_0^t \int_0^\tau K(u) du d\tau \right| \right) \\ &\leq \frac{ct^2}{2} \exp(-\frac{1}{2}t^2(\sigma^2 - c)). \end{aligned}$$

Hence it follows that

$$\int_0^Q \left| \frac{f(t) - \exp(-\frac{1}{2}t^2\sigma^2 + imt)}{t} \right| dt \leq \int_0^Q \frac{1}{2} ct \exp(-\frac{1}{2}t^2(\sigma^2 - c)) dt \leq \frac{1}{2} \frac{c}{\sigma^2 - c}.$$

Thus, from Esseen's theorem ([5], 20.3.A) we have

$$\sup_x \left| F(x) - \Phi\left(\frac{x-m}{\sigma}\right) \right| \leq \frac{1}{\pi} \frac{c}{\sigma^2 - c} + \frac{24}{\pi} \frac{1}{\sqrt{2\pi}\sigma Q},$$

which completes the proof of Lemma 5.

5. Proofs of the theorems. By ε -independence of the statistics $S_1(X_1, X_2)$ and $S_2(X_1, X_2)$ and by virtue of Lemma 1 it follows that there exist independent random variables Y_i ($i = 1, 2$) such that Y_i have finite moments of all orders, $S_1(Y_1, Y_2)$ and $S_2(Y_1, Y_2)$ are 13ε -independent statistics and $\sup_x |F_{X_i}(x) - F_{Y_i}(x)| \leq \varepsilon$ ($i = 1, 2$).

If Y_i ($i = 1, 2$) are $c_5/\sqrt{\ln 1/\varepsilon}$ -normal, then there exists a constant C_1 such that X_1, X_2 are $C_1/\sqrt{\ln 1/\varepsilon}$ -normal.

Let us show that there exists a constant c_5 such that Y_1, Y_2 are $c_5/\sqrt{\ln 1/\varepsilon}$ -normal. Let us remark that multiplication of random variables by numbers not equal to zero preserves their properties to be ε -independent and ε -normally distributed. Therefore it is enough to show that if

$$(22) \quad U = Y_1 + Y_2, \quad V = Y_1 + \alpha Y_2,$$

where $\alpha < 0$ and U, V are 13ε -independently distributed statistics, then there exists a constant c_5 such that Y_1, Y_2 are $c_5/\sqrt{\ln 1/\varepsilon}$ -normal. First we need the following

LEMMA 6. *Let U, V given by (22) be 13ε -independently distributed statistics and let $f_i(t) = f_{Y_i}(t)$ ($i = 1, 2$). If*

$$q_1 = q\left(\varepsilon, \frac{\Theta}{20}, f_1 \cdot f_2\right)$$

and

$$(23) \quad Q_1 = \min(q_1, \varepsilon^{-1/5}),$$

then there exists a function $K_2(t)$ such that for each $t \in [-Q_1, Q_1]$

$$(24) \quad |K_2(t)| \leq c_6 \cdot \varepsilon^{\Theta/10}$$

and

$$f_2(t) = \exp\left(-\frac{1}{2}\sigma_2^2 t^2 + im_2 t + \int_0^t \int_0^\tau K_2(u) du d\tau\right),$$

where

$$\sigma_2^2 = \frac{\text{Var}(Y_1 + \alpha Y_2)}{\alpha(\alpha - 1)}, \quad m_2 = \mathbb{E} Y_2.$$

Proof. Let $\varphi(u, v) = \varphi_{U, V}(u, v)$ and

$$(25) \quad \begin{aligned} A_1 &= A_1(t) = \mathbb{E}[(Y_1 + \alpha Y_2)^2 \exp\{it(Y_1 + Y_2)\}], \\ A_2 &= A_2(t) = \mathbb{E}[(Y_1 + \alpha Y_2) \exp\{it(Y_1 + Y_2)\}]. \end{aligned}$$

By independence of the random variables Y_1, Y_2 we have

$$A_1 = -f_1''(t)f_2(t) - 2\alpha f_1'(t)f_2'(t) - \alpha^2 f_1(t)f_2''(t).$$

On the other hand, we have

$$A_1 = \mathbb{E}(V^2)f_1(t)f_2(t) + \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} v^2 e^{itu} d\varphi(u, v).$$

Hence we obtain

$$(26) \quad -R_2(t) = f_1''(t)f_2(t) + \alpha^2 f_1(t)f_2''(t) + 2\alpha f_1'(t)f_2'(t) + \mathbb{E}(V^2)f_1(t)f_2(t),$$

where

$$R_2(t) = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} v^2 e^{itu} d\varphi(u, v).$$

Similarly, it follows from (25) that

$$(27) \quad f_1'(t)f_2(t) + \alpha f_1(t)f_2'(t) = i\mathbb{E}(V)f_1(t)f_2(t) + \bar{R}_2(t),$$

where

$$\bar{R}_2(t) = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} ive^{itu} d\varphi(u, v).$$

After the differentiation of (27) we obtain from (26) that for every $t \in [-q_1, q_1]$

$$[\ln f_2(t)]'' = \frac{-\text{Var } V}{\alpha(\alpha - 1)} + \frac{H_2(t)}{f_1(t)f_2^2(t)},$$

where

$$(28) \quad H_2(t) = \frac{f_2(t)}{\alpha(\alpha - 1)} \left[\mathbb{E}(V)\bar{R}_2(t) - i\bar{R}_2'(t) - i(\alpha - 1) \frac{\bar{R}_2(t)f_2'(t)}{f_2(t)} - R_2(t) \right].$$

Hence, by using the facts that

$$\alpha < 0, \quad f_2'(0) = im_2, \quad \frac{\text{Var}(V)}{\alpha(\alpha - 1)} = \sigma_2^2,$$

we obtain by integration that

$$f_2(t) = \exp\left(-\frac{1}{2}\sigma_2^2 t^2 + im_2 t + \int_0^t \int_0^\tau \frac{H_2(u)}{f_1(u)f_2^2(u)} dud\tau\right)$$

for every $t \in [-q_1, q_1]$.

Let

$$K_2(t) = \frac{H_2(t)}{f_1(t)f_2^2(t)}.$$

We shall show that there exists a constant c'_6 such that

$$|K_2(t)| \leq c'_6 \max(E|Y_1|, E|Y_2|, 1) [1 + \max_{i=1,2} (M_\Theta(Y_i))] \varepsilon^{\Theta/10}$$

for every $t \in [-Q_1, Q_1]$.

Since

$$(30) \quad E|V| \leq E|Y_1| - \alpha E|Y_2| \quad \text{and} \quad |f'_2(t)| \leq E|Y_2|$$

we see from (28) that

$$|H_2(t)| \leq \frac{1}{\alpha(\alpha-1)} [(E|Y_1| + (1-2\alpha)E|Y_2|) |\bar{R}_2(t)| + |\bar{R}'_2(t)| + |R_2(t)|].$$

By 13ε -independence of U, V and by virtue of (30), Lemma 3 ((14), (16)) and Lemma 4 ((17)) we obtain

$$|H_2(t)| \leq \max(E|Y_1|, E|Y_2|, 1) \{104N\varepsilon(1-\alpha)[2+2N|t|+3N+2N^2|t|] + 4N^{-\Theta} [\max_{j=1,2} (M_\Theta Y_j)] [32(1-\alpha) + 11(1-\alpha)^4]\} \frac{1}{\alpha(\alpha-1)}$$

for every $t \in [-q_1, q_1]$.

Substituting N instead of $|t|$ and then $\varepsilon^{-1/5}$ instead of N we obtain

$$|H_2(t)| \leq \max(E|Y_1|, E|Y_2|, 1) [936(1-\alpha) + \max_{i=1,2} (M_\Theta Y_i) 172(1-\alpha)^4] \frac{1}{\alpha(\alpha-1)} \varepsilon^{\Theta/5}.$$

It follows from (23) that there exists a constant c'_6 such that for each $t \in [-Q_1, Q_1]$

$$|K_2(t)| \leq c'_6 \max(E|Y_1|, E|Y_2|, 1) [1 + \max_{i=1,2} (M_\Theta Y_i)] \cdot \varepsilon^{\Theta/10},$$

hence there exists a constant c_6 such that for $t \in [-Q_1, Q_1]$ formula (24) holds. This completes the proof of Lemma 6.

We turn now to the proof of Theorem 1. Let us exchange Y_1 for Y_2 and let us put α^{-1} instead of α . Then, by Lemma 6, there exists a function

$K_1(t)$ and a constant c_7 such that for every $t \in [-Q_1, Q_1]$

$$(31) \quad |K_1(t)| \leq c_7 \varepsilon^{\Theta/10}$$

and

$$(32) \quad f_1(t) = \exp\left(-\frac{1}{2}\sigma_1^2 t^2 + im_1 t + \int_0^t \int_0^\tau K_1(u) du d\tau\right),$$

where

$$\sigma_1^2 = \frac{\text{Var } V}{1 - \alpha}, \quad m_1 = \mathbf{E} Y_1.$$

It follows from Lemma 6 and from formulas (31) and (32) that the assumptions of Lemma 2 are satisfied if $f(t) = f_1(t)f_2(t)$, $\alpha_0 = \Theta/20$ and $q = Q_1$. Hence, we have

$$q_1 \geq \left[\frac{10}{\Theta} (\sigma_1^2 + \sigma_2^2 + (c_6 + c_7) \cdot \varepsilon^{\Theta/10}) \right]^{-1/2} \cdot \sqrt{\ln \frac{1}{\varepsilon}}.$$

By virtue of (20) and (24) we obtain that for ε sufficiently small condition (21) is satisfied. Thus, Lemmas 5 and 6, and formulas (31) and (32) imply that there exists a constant c_5 such that in both cases $Q_1 = q_1$ and $Q_1 = \varepsilon^{-1/5}$ the inequality

$$\left| F_{Y_i}(x) - \Phi\left(\frac{x - m_i}{\sigma_i}\right) \right| \leq \frac{c_5}{\sqrt{\ln 1/\varepsilon}} \quad (i = 1, 2)$$

holds for each $x \in R$.

This completes the proof of Theorem 1.

Proof of Theorem 2. It follows from Lemma 1 that in order to prove Theorem 2 it is enough to show that there exists a constant c_8 such that if $T_1 = S$, $T_2 = S_3$ and Y_i ($i = 1, 2, \dots, n$) are defined by (12), then Y_1, \dots, Y_n are $c_8/\sqrt{\ln 1/\varepsilon}$ -normally distributed random variables.

Let us assume $\varphi(u, v) = \varphi_{SS_3}(u, v)$, $f(t) = f_{Y_1}(t)$ and $A_3 = A_3(t) = \mathbf{E}(S_3 \exp(itS))$.

The independence of the random variables Y_i ($i = 1, \dots, n$) and properties (8) and (9) of S_3 imply that

$$A_3 = - \sum_{j=1}^n a_{jj} (\ln f(t))'' f^n(t) \quad \text{and} \quad A_3 = \text{Var}(Y_1) \sum_{j=1}^n a_{jj} f^n(t) + R(t),$$

where

$$R(t) = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} v e^{itv} d\varphi(u, v).$$

Hence if $\text{Var}(Y_1) = \sigma^2$, $E(Y_1) = m$ and $q_2 = (\varepsilon, \Theta/8n, f)$, then

$$[\ln f(t)]'' = -\sigma^2 - \frac{1}{\sum_{j=1}^n a_{jj}} \frac{R(t)}{f^n(t)} \quad \text{for each } t \in [-q_2, q_2].$$

Thus for each $t \in [-q_2, q_2]$

$$f(t) = \exp \left\{ -\frac{1}{2}t^2\sigma^2 + imt - \int_0^t \int_0^\tau \frac{R(u)}{\sum_{j=1}^n a_{jj}f^n(u)} du d\tau \right\}.$$

Without loss of generality we may assume that

$$\max\{|a_{11}|, \dots, |a_{nn}|, |b_1|, \dots, |b_n|\} \leq 1.$$

Since $S(Y_1, \dots, Y_n)$, $S_3(Y_1, \dots, Y_n)$ are $(6n+1)$ ε -independently distributed statistics, we see from Lemma 3 (13) and Lemma 4 ((18), (19)) that for every $t \in [-q_2, q_2]$

$$(35) \quad |R(t)| \leq 8N(6n+1)\varepsilon(1+N|t|) + 18n^4N^{-\Theta}M_\Theta(Y_1) + 16n^4N^{-\Theta}.$$

Let

$$(36) \quad N = \varepsilon^{-1/4}, \quad Q = \min(q_2, \varepsilon^{-1/4}), \quad K_3(t) = \frac{R(t)}{\sum_{j=1}^n a_{jj}f^n(t)},$$

$$c_9 = \frac{16(6n+1) + 18n^4}{\left| \sum_{j=1}^n a_{jj} \right|}.$$

It follows from (33)-(36) that for every $t \in [-Q_2, Q_2]$

$$(37) \quad |K_3(t)| \leq c_9(1 + M_\Theta(Y_1))\varepsilon^{\Theta/8}.$$

Moreover, (33), (34), (37) and Lemma 2 imply that

$$q_2 \geq \left[\frac{4n}{\Theta} \left(\sigma^2 + c_9(1 + M_\Theta(Y_1))\varepsilon^{\Theta/8} \right) \right]^{-1/2} \sqrt{\ln \frac{1}{\varepsilon}}.$$

Since condition (21) is satisfied for sufficiently small values of ε , we deduce from Lemma 5 that there exists a constant c_8 such that in both cases, $Q_2 = q_2$ and $Q_2 = \varepsilon^{-1/4}$, the inequality

$$\sup_x \left| F_{Y_1}(x) - \Phi \left(\frac{x-m}{\sigma} \right) \right| \leq \frac{c_8}{\sqrt{\ln 1/\varepsilon}}$$

holds.

This completes the proof of Theorem 2.

Proof of Theorem 3. To prove Theorem 3 it is enough to show that the assumptions of Lemma 5 are satisfied.

Since the statistics S_ε has ε -polynomial regression of order ≤ 2 on S , we have

$$\mathbb{E} \left(\sum_{j,k=1}^n a_{jk} X_j X_k + \sum_{j=1}^n b_j X_j \mid \sum_{j=1}^n X_j = x \right) = \beta_0 + \beta_1 x + \beta_2 x^2 + \eta(x)$$

and

$$\sup_x |\eta(x)| \leq \varepsilon.$$

If one multiplies the above expression by $\exp(itx)$ and takes the expectation, then

$$\begin{aligned} & \mathbb{E} \left[\left(\sum_{j,k=1}^n a_{jk} X_j X_k + \sum_{j=1}^n b_j X_j \right) \exp \left(it \sum_{j=1}^n X_j \right) \right] \\ &= \beta_0 \mathbb{E} \left[\exp \left(it \sum_{j=1}^n X_j \right) \right] + \beta_1 \mathbb{E} \left[\sum_{j=1}^n X_j \exp \left(it \sum_{j=1}^n X_j \right) \right] + \\ &+ \beta_2 \mathbb{E} \left[\left(\sum_{j=1}^n X_j \right)^2 \exp \left(it \sum_{j=1}^n X_j \right) \right] + \mathbb{E} \left[\eta \left(\sum_{j=1}^n X_j \right) \exp \left(it \sum_{j=1}^n X_j \right) \right]. \end{aligned}$$

We put $f(t) = f_{X_1}(t)$, $q_3 = q(\varepsilon, 1/2n, f)$,

$$\sigma^2 = \frac{\beta_0}{\sum_{j=1}^n a_{jj} - n\beta_2} \quad \text{and} \quad \eta_1(t) = \mathbb{E} \left[\eta \left(\sum_{j=1}^n X_j \right) \exp \left(it \sum_{j=1}^n X_j \right) \right].$$

It follows from (11) that for every $t \in [-q_3, q_3]$

$$[\ln f(t)]'' = -\sigma^2 - \frac{\eta_1(t)}{\left(\sum_{j=1}^n a_{jj} - n\beta_2 \right) f^n(t)}.$$

By integration we have for each $t \in [-q_3, q_3]$

$$f(t) = \exp \left(-\frac{1}{2} \sigma^2 t^2 + imt - \int_0^t \int_0^\tau \frac{\sigma^2}{\beta_0} \frac{\eta_1(u)}{f^n(u)} du d\tau \right).$$

Since for each $t \in [-q_3, q_3]$

$$\left| \frac{\sigma^2}{\beta_0} \frac{\eta_1(t)}{f^n(t)} \right| \leq c_{10} \varepsilon^{1/2},$$

where $c_{10} = |\sigma^2/\beta_0|$, the assumptions of Lemma 5 are satisfied. Thus, there exists a constant C_4 such that

$$\sup_x \left| F_{X_1}(x) - \Phi \left(\frac{x-m}{\sigma} \right) \right| \leq \frac{C_4}{\sqrt{\ln 1/\varepsilon}}.$$

The proof of Theorem 3 is therefore completed.

6. The case of Lévy's metric. $L(F_1, F_2)$ is said to be the *distance of the distribution functions* F_1, F_2 in Lévy's metric, i.e. the metric defined by the formula

$$L(F_1, F_2) = \inf \{h: F_1(x-h) - h \leq F_2(x) \leq F_1(x+h) + h \text{ for all } x\}.$$

It is well-known that the convergence of distribution functions in Lévy's metric is equivalent to the weak convergence.

In [13] Zolotarev proved that for any distribution functions F_1, F_2 and for every number $Q > e$

$$L(F_1, F_2) \leq \frac{1}{\pi} \int_0^Q \frac{|f_1(t) - f_2(t)|}{t} dt + 2e \frac{\log Q}{Q},$$

where $f_i(t)$ are the characteristic functions of F_i ($i = 1, 2$).

Using this theorem instead of Esseen's one we obtain Theorems 1'-3'.

Let K_N be the family of distribution functions of the normal non-degenerate law and let

$$e_N(X) = \inf_{F \in K_N} \{L(F_X, F)\}.$$

THEOREM 1'. *Under the assumptions of Theorem 1*

$$e_N(X_i) \leq C_{11} \frac{\log \log \frac{1}{\varepsilon}}{\sqrt{\log \frac{1}{\varepsilon}}} \quad (i = 1, 2).$$

THEOREM 2'. *If the assumptions of Theorem 2 are satisfied, then*

$$e_N(X_i) \leq C_{12} \frac{\log \log \frac{1}{\varepsilon}}{\sqrt{\log \frac{1}{\varepsilon}}} \quad (i = 1, \dots, n).$$

COROLLARY. *If the assumptions of the Corollary to Theorem 2 are satisfied, then*

$$e_N(X_i) \leq C_{13} \frac{\log \log \frac{1}{\varepsilon}}{\sqrt{\log \frac{1}{\varepsilon}}} \quad (i = 1, \dots, n).$$

THEOREM 3'. *If the assumptions of Theorem 3 are satisfied, then*

$$e_N(X_i) \leq c_{14} \frac{\log \log \frac{1}{\varepsilon}}{\sqrt{\log \frac{1}{\varepsilon}}} \quad (i = 1, \dots, n).$$

As the proofs of the above theorems are very similar to the proofs given in Section 5, they are omitted.

Acknowledgment. We wish to thank Dr. W. Klonecki for his helpful discussions.

References

- [1] Хоанг Хыу Ньы, *Об устойчивости некоторых характеристических свойств нормальной совокупности*, Теория вероятн. прим. 13 (1968), р. 308-314.
- [2] — *Оценка устойчивости одной характеристики экспоненциального закона*, Лит. мат. сборник 8 (1968), р. 175-177.
- [3] R. G. Laha, *On some properties of the normal and gamma distributions*, Proc. Amer. Math. Soc. 7 (1956), р. 172-174.
- [4] Ю. В. Линник, *Общие теоремы о разложении безгранично делимых законов*, Теория вероятн. прим. 3 (1958), р. 3-40.
- [5] M. Loève *Probability theory*, Van Nostrand, New York 1955.
- [6] E. Lukacs, R. G. Laha, *On some characterization problems connected with quadratic regression*, Biometrika 47 (1960), р. 335-343.
- [7] — *Applications of characteristic functions*, Griffin, London 1964.
- [8] L. D. Meshalkin, *On the robustness of some characterizations of the normal distribution*, Ann. Math. Statist. 39 (1968), р. 1747-1750.
- [9] Н. А. Сапогов, *Проблема устойчивости для теоремы Крамера*, Изв. АН СССР, сер. матем. 15 (1951), р. 205-218.
- [10] — *О независимых слагаемых суммы случайных величин распределенной приближенно нормально*, Вестник Ленингр. Чнив. 19 (1959), р. 78-105.
- [11] О. В. Шалаевский, *О устойчивости теоремы Д. А. Райкова*, Вестник Ленингр. Унив. 14 (1959), р. 41-49.
- [12] В. М. Золотарев, *К вопросу об устойчивости разложения нормального закона распределения на компоненты*, Теория вероятн. прим. 13 (1968), р. 738-742.
- [13] — *Несколько новых неравенств связанных с метрикой Леви*, Докл. АН СССР 190 (1970), р. 1019-1021.

MATHEMATICAL INSTITUTE
POLISH ACADEMY OF SCIENCES

*Received on 15. 6. 1971;
revised version on 10. 2. 1972*

BOGUSŁAWA BEDNAREK-KOZEK i A. KOZEK (Wrocław)

O STABILNOŚCI WŁASNOŚCI CHARAKTERYZUJĄCYCH ROZKŁAD NORMALNY

STRESZCZENIE

Jeśli X_1, \dots, X_n są niezależnymi zmiennymi losowymi, to ze stochastycznej niezależności lub z wielomianowej regresji jednej statystyki względem drugiej w wielu wypadkach wynika, że rozkłady zmiennych losowych X_1, \dots, X_n są już określone.

W pracy tej badamy stabilność pewnych własności charakteryzujących rozkład normalny. Przez zastąpienie niezależności statystyk słabszym założeniem ich ε -niezależności otrzymujemy charakteryzację rozkładów wyjściowych zmiennych losowych, jako „bliskich” odpowiednim rozkładom normalnym. Jako miary odległości między dwoma dystrybuantami F_1, F_2 używamy metryki $\rho(F_1, F_2) = \sup_x |F_1(x) - F_2(x)|$ oraz metryki Lévy'ego.
