

A PROOF OF MENGER'S THEOREM BY CONTRACTION

FRANK GÖRING

Department of Mathematics
Technical University of Ilmenau
D-98684 Ilmenau Germany

Abstract

A short proof of the classical theorem of Menger concerning the number of disjoint AB -paths of a finite graph for two subsets A and B of its vertex set is given. The main idea of the proof is to contract an edge of the graph.

Keywords: connectivity, disjoint paths, digraph, Menger.

2000 Mathematics Subject Classifications: 05C40.

Proofs of Menger's Theorem are given in [7, 6, 4, 8, 2]. A short proof is given by T. Böhme, F. Göring and J. Harant in [1]; another short proof based on edge deletion is given by the author in [5]. The new idea here is to get a short proof by contracting an arbitrary edge of the original graph.

For terminology and notation not defined here we refer to [3]. A graph with no edges is denoted by its vertex set. Let G be a finite graph (loops and multiple edges being allowed). For an edge e of G let $G - e$ and G/e denote the graphs obtained from G by removing e and contracting e to one vertex v_e , respectively. For (possibly empty) sets of vertices A and B of G let an AB -separator be a set of vertices of G such that the graph obtained from G by deleting these vertices contains no path from A to B . Note that a single vertex of $A \cap B$ is considered as a path from A to B , too. An AB -connector is a subgraph of G such that each of its components is a path from A to B having only one vertex in common with A and B , respectively. In particular the empty graph is also an AB -connector. If we contract an edge incident with a vertex of A or B then the resulting vertex is considered to be a vertex of A or B , respectively.

Theorem (Menger, 1927). *Let G be a finite graph, A and B sets of vertices of G , and s the minimum number of vertices forming an AB -separator. Then there is an AB -connector C with $|C \cap A| = s$.*

Proof. If G is edgeless then set $C = A \cap B$. Suppose, G is a counterexample with $|E(G)|$ minimal. Then G contains an edge e from x to y and G/e has an AB -separator S with $|S| < s$, otherwise we are done. Obviously, $v_e \in S$. Then $P = (S \setminus \{v_e\}) \cup \{x, y\}$ is an AB -separator of G with $|P| = |S| + 1 = s$. An AP -separator (as well as an PB -separator) of $G - e$ is an AB -separator of G . Consequently, $G - e$ has an AP -connector X and a PB -connector Y containing P . Since $X \cap Y = P$, the set $C = (X \cup Y)$ is an AB -connector of G with $|C \cap A| = s$, a contradiction. ■

References

- [1] T. Böhme, F. Göring and J. Harant, *Menger's Theorem*, J. Graph Theory **37** (2001) 35–36.
- [2] W. McCuaig, *A simple proof of Menger's theorem*, J. Graph Theory **8** (1984) 427–429.
- [3] R. Diestel, *Graph Theory* (2nd edition), (Springer-Verlag, New York, 2000).
- [4] G.A. Dirac, *Short proof of Menger's graph theorem*, Mathematika **13** (1966) 42–44.
- [5] F. Goering, *Short Proof of Menger's Theorem*, to appear in Discrete Math.
- [6] T. Grünwald (later Gallai), *Ein neuer Beweis eines Mengerschen Satzes*, J. London Math. Soc. **13** (1938) 188–192.
- [7] K. Menger, *Zur allgemeinen Kurventheorie*, Fund. Math. **10** (1927) 96–115.
- [8] J.S. Pym, *A proof of Menger's theorem*, Monatshefte Math. **73** (1969) 81–88.

Received 8 June 2000

Revised 21 May 2001