

A NOTE ON STRONGLY MULTIPLICATIVE GRAPHS

CHANDRASHEKAR ADIGA

H.N. RAMASWAMY

AND

D.D. SOMASHEKARA

Department of Studies in Mathematics
University of Mysore
Manasagangotri
Mysore – 570 006, INDIA

Abstract

In this note we give an upper bound for $\lambda(n)$, the maximum number of edges in a strongly multiplicative graph of order n , which is sharper than the upper bound obtained by Beineke and Hegde [1].

Keywords and phrases: graph labeling, strongly multiplicative graphs.

2000 Mathematics Subject Classification: 05C78.

1. INTRODUCTION

In an interesting paper [1] L.W. Beineke and S.M. Hegde have studied strongly multiplicative graphs. In fact, they showed that certain graphs like trees, wheels and grids are strongly multiplicative. They also obtained an upper bound for the maximum number of edges $\lambda(n)$ for a given strongly multiplicative graph of order n . Erdős [2] has earlier obtained an asymptotic formula for $\lambda(n)$.

In this note we obtain an upper bound for $\lambda(n)$ which is sharper than that upper bound given in [1], for large values of n .

2. MAIN RESULT

We recall the definition of strongly multiplicative graphs given in [1].

Definition. A graph with n vertices is said to be strongly multiplicative if its vertices can be labeled $1, 2, \dots, n$ so that the values on the edges, obtained as the product of the labels of their end vertices, are all distinct.

Let $\lambda(n)$ denote the maximum number of edges in a strongly multiplicative graph of order n . Thus

$$\lambda(n) = |\{rs | 1 \leq r < s \leq n\}|.$$

Also, let

$$\delta(n) = \lambda(n) - \lambda(n-1).$$

The number theoretic functions λ and δ were studied in [1]. We first give an upper bound for $\delta(n)$ which leads to an improvement of Theorem 3.1 [1].

Lemma. If $p(n)$ denotes the least prime divisor of n , then

$$\delta(n) \leq \begin{cases} n - \frac{n}{p(n)} + 1 & \text{for } n \equiv 0, 1, 3 \pmod{4}, \\ n - \frac{n}{p(n)} & \text{for } n \equiv 2 \pmod{4}. \end{cases}$$

Proof. Let $p(n) = p$. We observe that $n \cdot 1$ can be written as $p\left(\frac{n}{p}\right)$, $n \cdot 2$ can be written as $2p\left(\frac{n}{p}\right), \dots, n\left(\frac{n}{p} - 1\right)$ can be written as $\left\{\left(\frac{n}{p} - 1\right)p\right\} \frac{n}{p}$. This shows that $\delta(n) \leq n - \frac{n}{p} + 1$. Note that if $n \equiv 2 \pmod{4}$, then $p = 2$, so that $\frac{n}{p}$ is odd and kp is even. Therefore $\delta(n) \leq n - \frac{n}{p(n)}$.

Now we obtain an upper bound for $\lambda(n)$ using the above Lemma.

Theorem. $\lambda(n) \leq \frac{n(n+1)}{2} + (n-2) - \left[\frac{n+2}{4}\right] - \sum_{i=2}^n \frac{i}{p(i)}$, where $[x]$ denotes the largest integer less than or equal to x .

Proof. Since $\delta(n) = \lambda(n) - \lambda(n-1)$, we have

$$\lambda(n) = \sum_{i=2}^n \delta(i).$$

On using the above Lemma, it follows that,

$$\begin{aligned}\lambda(n) &\leq \sum_{i=2}^n \left(i - \frac{i}{p(i)} + 1 \right) - \left\lceil \frac{n+2}{4} \right\rceil \\ &= \left(\frac{n(n+1)}{2} - 1 \right) - \sum_{i=2}^n \frac{i}{p(i)} + (n-1) - \left\lceil \frac{n+2}{4} \right\rceil,\end{aligned}$$

and the result follows.

Remark. The following table shows that the upper bound for $\lambda(n)$ given in the above Theorem is sharper than the upper bound given by L.W. Beineke and S.M. Hegde [1] for large values of n .

n	Upper bound for $\lambda(n)$ using our Theorem	Upper bound for $\lambda(n)$ given by Beineke and Hegde
25	236	240
27	268	279
42	629	661
60	1,263	1,350

Acknowledgement

We thank the referee for his valuable comments and suggestions.

REFERENCES

- [1] L.W. Beineke and S.M. Hegde, *Strongly multiplicative graphs*, Discuss. Math. Graph Theory **21** (2001) 63–76.
- [2] P. Erdős, *An asymptotic inequality in the theory of numbers*, Vestnik Leningrad, Univ. **15** (1960) 41–49.

Received 24 October 2001

Revised 6 July 2002