

MARIAN NOWAK

Bochner representable operators on Köthe-Bochner spaces

Abstract. Let E be a Banach function space and X be a real Banach space. We study Bochner representable operators from a Köthe-Bochner space $E(X)$ to a Banach space Y . We consider the problem of compactness and weak compactness of Bochner representable operators from $E(X)$ (provided with the natural mixed topology) to Y .

2000 Mathematics Subject Classification: 47B38, 47B05, 46E40, 46A70.

Key words and phrases: Köthe-Bochner spaces, Generalized DF-spaces, Bochner representable operators, Weakly compact operators, Compact operators, Mixed topologies, Mackey topologies.

1. Introduction and preliminaries. The integral representation of linear operators on function spaces (in particular, Lebesgue spaces and Orlicz spaces) has been the object of much study (see [Ph], [DP], [G₁], [G₂], [DS], [D], [U₁], [U₂], [A₁], [A₂], [Na₁], [Na₂]). In particular, K. Andrews [A₁, Theorems 2 and 5], [A₂, Theorem 3] and V.G. Navodnov [Na₂, Corollary] obtained a Dunford-Pettis-Phillips type theorem for compact operators (resp. weakly compact operators) from the space $L^1(X)$ of Bochner integrable functions to a Banach space Y . Moreover, V.G. Navodnov ([Na₂], [Na₁]) has considered Bochner representable operators from a Köthe-Bochner space $E(X)$ to a Banach space Y .

In Section 2 we study the problem of compactness and weak compactness of Bochner representable operators from a Köthe-Bochner space $E(X)$ (provided with the natural mixed topology $\gamma_{E(X)}$) to a Banach space Y . The space $(E(X), \gamma_{E(X)})$ is a generalized DF-space, so we can apply the Grothendieck's DF techniques (see [G₁], [G₂], [Ru]).

Throughout the paper $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are real Banach spaces with the Banach duals X^* and Y^* . Let B_X and B_Y denote the closed unit balls in X and Y . Let $\mathcal{L}(X, Y)$ stand for the space of all bounded linear operators

from X to Y provided with the uniform convergence norm $\|\cdot\|_{X \rightarrow Y}$. The *strong operator topology* (briefly SOT) is the topology on $\mathcal{L}(X, Y)$ defined by the family of seminorms $\{p_x : x \in X\}$, where $p_x(U) := \|U(x)\|_Y$ for $U \in \mathcal{L}(X, Y)$ (see [DS, p. 475–477]). Let \mathbb{N} stand for the set of natural numbers.

Let (Ω, Σ, μ) be a complete finite measure space. Assume that $(E, \|\cdot\|_E)$ is a Banach function space and let E' stand for the Köthe dual of E . Then the associated norm $\|\cdot\|_{E'}$ on E' can be defined for $v \in E'$ by

$$\|v\|_{E'} = \sup \left\{ \left| \int_{\Omega} u(\omega) v(\omega) d\mu \right| : u \in E, \|u\|_E \leq 1 \right\}.$$

A Banach function space $(E, \|\cdot\|_E)$ is said to be *perfect*, if $E = E''$ and $\|u\|_{E'} = \|u\|_E$ for $u \in E$. It is well known that $(E, \|\cdot\|_E)$ is perfect if and only if $\|\cdot\|_E$ satisfies both the σ -Fatou property and the σ -Levy property (see [KA, Theorem 6.1.7]).

From now we will assume that $L^\infty \subset E \subset L^1$, where the inclusion maps are continuous. Moreover, we will assume that $\|\cdot\|_{E'}$ on E' is order continuous. Since $(L^1)' = L^\infty$, the space L^1 is excluded.

By $L^0(X)$ we denote the set of μ -equivalence classes of all strongly Σ -measurable functions $f : \Omega \rightarrow X$. Let $\mathcal{T}_0(X)$ stand for the topology on $L^0(X)$ of the F -norm $\|\cdot\|_{L^0(X)}$ that generates convergence in measure on sets of finite measure.

For $f \in L^0(X)$ let $\tilde{f}(\omega) = \|f(\omega)\|_X$ for $\omega \in \Omega$. Then the space

$$E(X) = \{f \in L^0(X) : \tilde{f} \in E\}$$

equipped with the norm $\|f\|_{E(X)} := \|\tilde{f}\|_E$ is a Banach space and is usually called a *Köthe-Bochner space* (see [CM], [L] for more details). We will denote by $\mathcal{T}_E(X)$ the topology of the norm $\|\cdot\|_{E(X)}$. Recall that the algebraic tensor product $E \otimes X$ is the subspace of $E(X)$ spanned by the functions of the form $u \otimes x$, $(u \otimes x)(\omega) = u(\omega)x$, where $u \in E$, $x \in X$ and $\omega \in \Omega$. For $r > 0$ denote

$$B_{E(X)}(r) = \{f \in E(X) : \|f\|_{E(X)} \leq r\}.$$

Let τ be a linear topology on $E(X)$. A linear operator $T : E(X) \rightarrow Y$ is said to be $(\tau, \|\cdot\|_Y)$ -compact (resp. $(\tau, \|\cdot\|_Y)$ -weakly compact) if there exists a neighbourhood U of 0 for τ such that $T(U)$ is relatively norm compact (resp. relatively weakly compact) in Y . By $Bd(E(X), \tau)$ we will denote the collection of all τ -bounded subsets of $E(X)$.

2. Bochner representable operators on Köthe-Bochner spaces. We start by recalling terminology concerning Bochner representable operators $T : E(X) \rightarrow Y$ (see [A₁], [A₂], [Na₁], [Na₂] for more details).

A function $\mathcal{K} : \Omega \rightarrow \mathcal{L}(X, Y)$ is said to be SOT-measurable if for every $x \in X$ the function $\mathcal{K}_x : \Omega \ni \omega \mapsto \mathcal{K}(\omega)(x) \in Y$ is strongly Σ -measurable. We

say that two SOT-measurable functions \mathcal{K}_1 and \mathcal{K}_2 are *SOT-equivalent* (briefly, $\mathcal{K}_1 \approx \mathcal{K}_2$) if $\mathcal{K}_1(\omega)(x) = \mathcal{K}_2(\omega)(x)$ for all $x \in X$ and μ -almost all $\omega \in \Omega$ (see [Na₁], [Na₂]).

Recall that a bounded linear operator $T : E(X) \rightarrow Y$ is said to be *Bochner representable* if there exists a SOT-measurable function $\mathcal{K} : \Omega \rightarrow \mathcal{L}(X, Y)$ (called the *representing kernel* for T) such that for each $f \in E(X)$ the function $\langle f, \mathcal{K} \rangle : \Omega \ni \omega \mapsto \langle f(\omega), \mathcal{K}(\omega) \rangle \in Y$ is Bochner integrable and

$$T(f) = \int_{\Omega} \langle f(\omega), \mathcal{K}(\omega) \rangle d\mu \quad \text{for all } f \in E(X).$$

The following theorem will be of importance (see [Na₁, Theorem 1]).

THEOREM 2.1 *For a SOT-measurable function $\mathcal{K} : \Omega \rightarrow \mathcal{L}(X, Y)$ the following statements are equivalent:*

- (i) \mathcal{K} is the representing kernel for a Bochner representable operator $T : E(X) \rightarrow Y$.
- (ii) There exists a SOT-measurable function $\mathcal{K}_0 : \Omega \rightarrow \mathcal{L}(X, Y)$ such that $\mathcal{K}_0 \approx \mathcal{K}$ and the μ -equivalence class of the function $\|\mathcal{K}_0(\cdot)\|_{X \rightarrow Y}$ belongs to E' .

Recall that the *mixed topology* $\gamma[\mathcal{T}_E(X), \mathcal{T}_0(X) \upharpoonright_{E(X)}]$ (briefly $\gamma_{E(X)}$) on $E(X)$ is the finest Hausdorff locally convex topology on $E(X)$ which agrees with $\mathcal{T}_0(X)$ on $\|\cdot\|_{E(X)}$ -bounded subsets of $E(X)$ (see [C, Chap. III], [N₂], [F, §3] for more details). Then

$$\mathcal{T}_0(X) \upharpoonright_{E(X)} \subset \gamma_{E(X)} \subset \mathcal{T}_E(X).$$

Since $B_{E(X)}(1)$ is closed in $(E(X), \mathcal{T}_0(X) \upharpoonright_{E(X)})$ (see [KA, Lemma 4.3.4]), by [W, Theorem 2.4.1] we get

$$(1) \quad Bd(E(X), \gamma_{E(X)}) = Bd(E(X), \|\cdot\|_{E(X)}).$$

This means that $(E(X), \gamma_{E(X)})$ is a generalized DF-space (see [Ru]). Hence using the Grothendieck classical results (see [Ru, p. 429], [G₁, Corollary 1 of Theorem 11], [G₂, Chap. IV, 4.3, Corollary 1 of Theorem 2]) we get:

THEOREM 2.2 *Let $T : E(X) \rightarrow Y$ be a $(\gamma_{E(X)}, \|\cdot\|_Y)$ -continuous linear operator which transforms $\gamma_{E(X)}$ -bounded sets into relatively norm compact (resp. relatively weakly compact) sets in Y . Then T is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -compact (resp. $(\gamma_{E(X)}, \|\cdot\|_Y)$ -weakly compact).*

A linear operator $T : E(X) \rightarrow Y$ is called $(\gamma, \|\cdot\|_Y)$ -linear if $\|T(f_n)\|_Y \rightarrow 0$ whenever $\|f_n\|_{L^0(X)} \rightarrow 0$ and $\sup_n \|f_n\|_{E(X)} < \infty$ (see [W]). It is known that a linear operator $T : E(X) \rightarrow Y$ is $(\gamma, \|\cdot\|_Y)$ -linear if and only if T is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -continuous (see [W, Theorem 2.6.1(iii)]).

PROPOSITION 2.3 *Assume that $(E, \|\cdot\|_E)$ is a perfect Banach function space such that the associated norm $\|\cdot\|_{E'}$ on E' is order continuous. Then every Bochner representable operator $T : E(X) \rightarrow Y$ is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -continuous.*

PROOF Assume that $T : E(X) \rightarrow Y$ is a Bochner representable operator. Then, by Theorem 2.1, there exists a SOT-measurable function $\mathcal{K}_0 : \Omega \rightarrow \mathcal{L}(X, Y)$ such that the μ -equivalence class $v_0 = [\|\mathcal{K}_0(\cdot)\|_{X \rightarrow Y}]$ belongs to E' and

$$T(f) = \int_{\Omega} \langle f(\omega), \mathcal{K}_0(\omega) \rangle d\mu \quad \text{for all } f \in E(X).$$

Hence for $f \in E(X)$ we get

$$\begin{aligned} \|T(f)\|_Y &= \left\| \int_{\Omega} \langle f(\omega), \mathcal{K}_0(\omega) \rangle d\mu \right\|_Y \\ &\leq \int_{\Omega} \|\langle f(\omega), \mathcal{K}_0(\omega) \rangle\|_Y d\mu \\ &\leq \int_{\Omega} \|f(\omega)\|_X \cdot \|\mathcal{K}_0(\omega)\|_{X \rightarrow Y} d\mu. \end{aligned}$$

Putting

$$\varphi_{v_0}(u) = \int_{\Omega} u(\omega) v_0(\omega) d\mu \quad \text{for } u \in E,$$

we see that φ_{v_0} is a γ -linear functional on E (see [N₁, Theorem 3.1]). It follows that T is a $(\gamma, \|\cdot\|_Y)$ -linear operator, i.e., T is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -continuous. ■

Before stating our main result we require a preliminary definition (see [A₁, p. 258]). A SOT-measurable function $\mathcal{K} : \Omega \rightarrow \mathcal{L}(X, Y)$ is said to have its essential range in the *uniformly norm compact operators* (resp. *uniformly weakly compact operators*) if there exists a relatively norm compact (resp. relatively weakly compact) set C in Y such that $\langle x, \mathcal{K}(\omega) \rangle \in C$ for μ -almost all $\omega \in \Omega$ and all $x \in B_X$.

First we recall the well known Dunford-Pettis-Phillips type theorem for compact operators (resp. weakly compact operators) from the space $L^1(X)$ to a Banach space Y (see [Na₂, Corollary]).

THEOREM 2.4 *For a bounded linear operator $T : L^1(X) \rightarrow Y$ the following statements are equivalent:*

- (i) T is $(\|\cdot\|_{L^1(X)}, \|\cdot\|_Y)$ -compact (resp. $(\|\cdot\|_{L^1(X)}, \|\cdot\|_Y)$ -weakly compact).
- (ii) T is Bochner representable operator with the representing kernel \mathcal{K} having its essential range in the uniformly norm compact operators (resp. uniformly weakly compact operators).

Now we are in position to extend and strengthen the implication (ii) \implies (i) of Theorem 2.4 for the case of linear operators from a Köthe-Bochner space $E(X)$ to a Banach space Y .

THEOREM 2.5 *Assume that $(E, \|\cdot\|_E)$ is a perfect Banach function space such that the associated norm $\|\cdot\|_{E'}$ on E' is order continuous. Let $T : E(X) \rightarrow Y$ be a Bochner representable operator with the representing kernel \mathcal{K} having its essential range in the uniformly norm compact operators (resp. uniformly weakly compact operators). Then T is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -compact (resp. $(\gamma_{E(X)}, \|\cdot\|_Y)$ -weakly compact).*

PROOF In view of Proposition 2.3 T is $(\gamma_{E(X)}, \|\cdot\|_Y)$ -continuous. Since $Bd(E(X), \gamma_{E(X)}) = Bd(E(X), \|\cdot\|_{E(X)})$ (see (1)), by Theorem 2.2 it is enough to show that for every $r > 0$, the set $T(B_{E(X)}(r))$ is relatively norm compact (resp. relatively weakly compact) in Y . Indeed, let $r > 0$. Since the inclusion map $(E, \|\cdot\|_E) \hookrightarrow (L^1, \|\cdot\|_{L^1})$ is supposed to be continuous, there exists $r_0 > 0$ such that $B_{E(X)}(r_0) \subset B_{L^1(X)}(\frac{1}{2})$.

Moreover, by our assumption there exists a relatively norm compact (resp. relatively weakly compact) set C in Y such that $\langle x, \mathcal{K}(\omega) \rangle \in C$ for μ -almost all $\omega \in \Omega$ and all $x \in B_X$. Now we shall show that $T(f) \in \overline{\text{conv}} C$ for all $f \in B_{E(X)}(r_0)$, where $\overline{\text{conv}} C$ stands for the norm closed (=weakly closed) convex hull of C in Y . Indeed, let $f \in B_{E(X)}(r_0)$. Then by [DS, p. 117] there exists a sequence (s_n) of X -valued Σ -simple functions such that $\|s_n(\omega)\|_X \leq 2\|f(\omega)\|_X$ μ -a.e. on Ω and $s_n \rightarrow f$ in μ -measure. Then $\|s_n - f\|_{L^0(X)} \rightarrow 0$ and $\sup_n \|s_n\|_{E(X)} \leq 2\|f\|_{E(X)} \leq 2r_0$. Hence $\|T(s_n) - T(f)\|_Y \rightarrow 0$, because T is $(\gamma, \|\cdot\|_Y)$ -linear.

Since $s_n = \sum_{i=1}^{k_n} \mathbb{1}_{A_{n,i}} \otimes x_{n,i}$ and $s_n \in B_{E(X)}(2r_0)$ for $n \in \mathbb{N}$, we get $s_n \in B_{L^1(X)}(1)$, i.e., $\sum_{i=1}^{k_n} \|x_{n,i}\|_X \mu(A_{n,i}) \leq 1$ for $n \in \mathbb{N}$. Hence, using [DU, Corollary 2.2.8, p. 48] we get

$$\begin{aligned} T(s_n) &= \sum_{i=1}^{k_n} T(\mathbb{1}_{A_{n,i}} \otimes x_{n,i}) \\ &= \sum_{i=1}^{k_n} \int_{A_{n,i}} \langle x_{n,i}, \mathcal{K}(\omega) \rangle d\mu \\ &= \sum_{i=1}^{k_n} \|x_{n,i}\|_X \int_{A_{n,i}} \left\langle \frac{x_{n,i}}{\|x_{n,i}\|_X}, \mathcal{K}(\omega) \right\rangle d\mu \\ &\in \left(\sum_{i=1}^{k_n} \|x_{n,i}\|_X \mu(A_{n,i}) \right) \overline{\text{conv}} C \subset \overline{\text{conv}} C. \end{aligned}$$

Hence $T(f) \in \overline{\text{conv}} C$, that is, $T(B_{E(X)}(r_0))$ is relatively norm compact (resp. relatively weakly compact) in Y , because $\overline{\text{conv}} C$ is norm compact in Y by Mazur's theorem (resp. $\overline{\text{conv}} C$ is weakly compact in Y by Krein-Šmulian's theorem). Since $T(B_{E(X)}(r)) \subset \frac{r}{r_0} \overline{\text{conv}} C$, we obtain that $T(B_{E(X)}(r))$ is relatively norm compact (resp. relatively weakly compact) in Y , as desired. \blacksquare

Now we will consider Bochner representable operators on the space $L^\infty(X)$. By a Young function we mean here a continuous convex mapping $\Phi : [0, \infty) \rightarrow [0, \infty)$ that vanishes only at 0 and $\Phi(t)/t \rightarrow 0$ as $t \rightarrow 0$ and $\Phi(t)/t \rightarrow \infty$ as $t \rightarrow \infty$.

The Orlicz-Bochner space $L^\Phi(X) := \{f \in L^0(X) : \int_\Omega \Phi(\lambda \|f(\omega)\|_X) d\mu < \infty$ for some $\lambda > 0\}$ is a Banach space under the norm $\|f\|_{L^\Phi(X)} := \inf\{\lambda > 0 : \int_\Omega \Phi(\|f(\omega)\|_X/\lambda) d\mu \leq 1\}$ (see [RR] for more details).

We will need the following characterization of the mixed topology $\gamma_{L^\infty(X)}$ on $L^\infty(X)$ (see [N₂, Theorem 4.5]).

THEOREM 2.6 *Then mixed topology $\gamma_{L^\infty(X)}$ on $L^\infty(X)$ is generated by the family of norms $\|\cdot\|_{L^\Phi(X)}$, where Φ runs over the family of all Young functions.*

As a consequence of Theorems 2.5 and 2.6 we get:

COROLLARY 2.7 *Let $T : L^\infty(X) \rightarrow Y$ be a Bochner representable operator and assume that the representing kernel \mathcal{K} for T has its range in the uniformly norm compact operators (resp. uniformly weakly compact operators). Then there exists a Young function Φ such that the set*

$$\left\{ \int_\Omega \langle f(\omega), \mathcal{K}(\omega) \rangle d\mu : f \in L^\infty(X), \|f\|_{L^\Phi(X)} \leq 1 \right\}$$

is relatively norm compact (resp. relatively weakly compact) in Y .

REFERENCES

- [A₁] K. Andrews, *Representation of compact and weakly compact operators on the space of Bochner integrable functions*, Pacific J. Math., **92**, no. 2 (1981), 257–267.
- [A₂] K. Andrews, *The Radon-Nikodym property for spaces of operators*, J. London Math. Soc., (2), **28** (1982), 113–122.
- [C] J.B. Cooper, *Saks Spaces and Applications to Functional Analysis*, North-Holland Publ. Co., Amsterdam, New York, 1978.
- [CM] P. Cembranos and J. Mendoza, *Banach spaces of vector-valued functions*, Lectures Notes in Math., **1676**, Springer Verlag, Berlin, Heidelberg, 1997.
- [D] N. Dinculeanu, *Vector Measures*, Pergamon Press, New York, 1967.
- DP, N. Dunford and J. Pettis, *Linear operators on summable functions*, Trans. Amer. Math. Soc., **47** (1940), 323–392.
- [DS] N. Dunford and J. Schwartz, *Linear Operators, Part. I, General Theory*, Interscience, Publ. Inc., New York, 1958.
- [DU] J. Diestel and J.J. Uhl, *Vector Measures*, Amer. Math. Soc., Math. Surveys **15**, Providence, 1977.
- [F] K. Feledziak, Comment. Math. Prace Mat. **37** (1997), 81–98.

-
- [G₁] A. Grothendieck, *Sur les espaces (F) and (DF)*, Summa Brasil. Math. **3** (1954), 57–122.
- [G₂] A. Grothendieck, *Espaces vectoriels topologiques*, Sao Paulo, 1954.
- [KA] L.V. Kantorovitch and A.V. Akilov, *Functional Analysis* (in Russian), Nauka Moscow, 1984 (3rd edition).
- [L] Pei-Kee Lin, *Köthe-Bochner Function Spaces*, Birkhäuser Verlag, Boston, Basel, Berlin, 2003.
- [Na₁] V.G. Navodnov, *Integral representation of operators acting from a Banach space of measurable vector-valued functions into a Banach space*, Izv. Vyssh. Uchebn. Zaved. Mat., no. **3** (1983), 82–84 (in Russian).
- [Na₂] V.G. Navodnov, *On the theory of integral operators in spaces of measurable vector functions* (Russian), Issled. Prikl. Math., **12** (1984), 162–174. Translated in J. Soviet Mat. 45, no. 2 (1989), 1093–1100.
- [N₁] M. Nowak, *Mixed topology on normed function spaces, I*, Bull. Polish Acad. Sci. Math., **36**, no. 5–6 (1988), 251–262.
- [N₂] M. Nowak, *Lebesgue topologies on vector-valued function spaces*, Math. Japonica **52** no. 2 (2000), 171–182.
- [Ph] R.S. Phillips, *On linear transformations*, Trans. Amer. Math. Soc. **48** (1940), 516–541.
- [RR] M.M. Rao and Z.D. Ren, *Theory of Orlicz spaces*, Marcel Dekker, New York, Basel, Hong Kong, 1991.
- [Ru] W. Ruess, *[Weakly] compact operators and DF spaces*, Pacific J. Math., **98**, no. 2 (1982), 419–441.
- [U₁] J.J. Uhl, *Compact operators on Orlicz spaces*, Rend. Semin. Math. Univ. Padova **42** (1969), 209–219.
- [U₂] J.J. Uhl, *On a class of operators on Orlicz spaces*, Studia Math. **40** (1971), 17–22.
- [W] A. Wiweger, *Linear spaces with mixed topology*, Studia Math., **20** (1961), 47–68.

MARIAN NOWAK
FACULTY OF MATHEMATICS, COMPUTER SCIENCE AND ECONOMETRICS
UL. SZAFRANA 4A, 65–516 ZIELONA GÓRA, POLAND
E-mail: M.Nowak@wmie.uz.zgora.pl

(Received: 5.02.2008)
