

ON FOUR-DIMENSIONAL RIEMANNIAN WARPED PRODUCT  
MANIFOLDS SATISFYING CERTAIN PSEUDO-SYMMETRY  
CURVATURE CONDITIONS

BY

RYSZARD DESZCZ (WROCLAW)

**1. Introduction.** Let  $(M, g)$  be a connected  $n$ -dimensional,  $n \geq 3$ , smooth Riemannian manifold with a not necessarily definite metric  $g$ . The manifold  $(M, g)$  is said to be pseudo-symmetric ([11]) if its curvature tensor  $R$  satisfies at every point of  $M$  the following condition:

(\*) the tensors  $R \cdot R$  and  $Q(g, R)$  are linearly dependent.

It is easy to see that if (\*) holds at a point of  $M$  then the Weyl conformal curvature tensor  $C$  satisfies at this point the condition

(\*\*) the tensors  $R \cdot C$  and  $Q(g, C)$  are linearly dependent.

A manifold  $(M, g)$  fulfilling (\*\*) at each point of  $M$  is called Weyl-pseudo-symmetric ([8]).

As was proved in ([12]), if  $n \geq 5$  then (\*) and (\*\*) are equivalent at each point at which  $C$  is non-zero. In particular, from this result it follows (see also [16]) that for  $n \geq 5$  the conditions  $R \cdot C = 0$  and  $R \cdot R = 0$  are equivalent at each point of  $(M, g)$  at which  $C \neq 0$ . On 4-manifolds, this is not always the case. A suitable example was given in [5] (Lemme 1.1). That example, by a certain modification, also gives rise to an example of a non-pseudo-symmetric manifold satisfying (\*\*) with  $R \cdot C$  non-zero (see [10]). Moreover, in [2] an example of a non-pseudo-symmetric conformally flat manifold of dimension  $n \geq 4$  was described.

In the present paper we shall prove (Section 4) that (\*) and (\*\*) are equivalent at every point of a 4-dimensional warped product manifold at which  $C$  does not vanish. From this it follows immediately that the above-mentioned Riemannian manifold obtained in [5] is a non-warped product manifold satisfying  $R \cdot C = 0$ . It is known that (\*) and (\*\*) are equivalent on manifolds isometrically immersed as hypersurfaces of a Euclidean space  $\mathbb{E}^{n+1}$ ,  $n \geq 4$  (see [3], Corollary).

If (\*) holds at a point of  $M$  then at this point the following condition is

fulfilled:

(\*\*\*) the tensors  $R \cdot S$  and  $Q(g, S)$  are linearly dependent,

where  $S$  denotes the Ricci tensor. A manifold  $(M, g)$  satisfying (\*\*\*) at every point of  $M$  is said to be Ricci-pseudo-symmetric ([14]). So, any pseudo-symmetric manifold is Ricci-pseudo-symmetric. However, the converse fails in general (see [14], [7]). We shall prove (Section 5) that (\*) and (\*\*\*) are equivalent at every point of a 4-dimensional warped product manifold at which the tensor  $S - (K/n)g$  does not vanish, where  $K$  is the scalar curvature.

Section 2 is concerned with some facts on pseudo-symmetric tensors. We recapitulate the basic formulas about warped products in Section 3. Finally, an analogue of Theorem 1 from [19] is mentioned at the end of that section.

**2. Pseudo-symmetric tensors.** Let  $(M, g)$  be an  $n$ -dimensional,  $n \geq 3$ , Riemannian manifold with a not necessarily definite metric  $g$ . We denote by  $\nabla$ ,  $R$ ,  $S$ ,  $C$  and  $K$  the Levi-Civita connection, the curvature tensor, the Ricci tensor, the Weyl conformal curvature tensor and the scalar curvature of  $(M, g)$  respectively. For a  $(0, k)$ -tensor field  $T$  on  $M$ ,  $k \geq 1$ , we define the tensor fields  $R \cdot T$  and  $Q(g, T)$  by

$$\begin{aligned} (R \cdot T)(X_1, \dots, X_k; X, Y) &= (R(X, Y) \cdot T)(X_1, \dots, X_k) \\ &= -T(R(X, Y)X_1, X_2, \dots, X_k) - \dots - T(X_1, \dots, X_{k-1}, R(X, Y)X_k), \\ Q(g, T)(X_1, \dots, X_k; X, Y) &= -((X \wedge Y) \cdot T)(X_1, \dots, X_k) \\ &= T((X \wedge Y)X_1, X_2, \dots, X_k) + \dots + T(X_1, \dots, X_{k-1}, (X \wedge Y)X_k) \end{aligned}$$

respectively, where  $R(X, Y)$  and  $X \wedge Y$  are derivations of the algebra of tensor fields on  $M$  and  $X_1, \dots, X_k, X, Y \in \mathfrak{X}(M)$ ,  $\mathfrak{X}(M)$  being the Lie algebra of vector fields on  $M$ . These derivations are extensions of the endomorphisms  $R(X, Y)$  and  $X \wedge Y$  of  $\mathfrak{X}(M)$  defined by

$$\begin{aligned} R(X, Y)Z &= \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, \\ (X \wedge Y)Z &= g(Z, Y)X - g(Z, X)Y \end{aligned}$$

respectively. A  $(0, k)$ -tensor field  $T$  is said to be *pseudo-symmetric* if the tensors  $R \cdot T$  and  $Q(g, T)$  are linearly dependent at every point of  $M$ . In the special case when  $R \cdot T$  vanishes on  $M$ , the tensor  $T$  is called *semi-symmetric*. A  $(0, 4)$ -tensor field  $T$  on  $M$  is said to be a *generalized curvature tensor* [18] if

$$\begin{aligned} T(X_1, X_2, X_3, X_4) + T(X_1, X_3, X_4, X_2) + T(X_1, X_4, X_2, X_3) &= 0, \\ T(X_1, X_2, X_3, X_4) &= -T(X_2, X_1, X_3, X_4), \\ T(X_1, X_2, X_3, X_4) &= T(X_3, X_4, X_1, X_2), \end{aligned}$$

for all  $X_i \in \mathfrak{X}(M)$ . For a generalized curvature tensor field  $T$  we define the *concircular curvature tensor*  $Z(T)$  by

$$Z(T) = T - \frac{K(T)}{n(n-1)}G,$$

where  $K(T)$  is the scalar curvature of  $T$  and  $G$  is the generalized curvature tensor defined by

$$G(X_1, X_2, X_3, X_4) = g((X_1 \wedge X_2)X_3, X_4).$$

A generalized curvature tensor  $T$  is called *trivial* at  $x \in M$  ([8]) if  $Z(T)$  vanishes at  $x$ . Similarly, for a symmetric  $(0, 2)$ -tensor field  $A$  we define the tensor  $Z(A)$  by

$$Z(A) = A - \frac{\text{tr}(A)}{n}g.$$

A symmetric  $(0, 2)$ -tensor field  $A$  is said to be *trivial* at  $x \in M$  if  $Z(A)$  vanishes at  $x$ .

**Remark 1** ([2], Lemma 1.1(iii)). Let  $T$  be a generalized curvature tensor (resp. a  $(0, 2)$ -symmetric tensor) at a point  $x$  of a manifold  $(M, g)$ . Then the equalities  $Z(T) = 0$  and  $Q(g, T) = 0$  are equivalent at this point.

If a generalized curvature tensor  $T$  (resp. a  $(0, 2)$ -symmetric tensor  $A$ ) is pseudo-symmetric then  $R \cdot T = L_T Q(g, T)$  (resp.  $R \cdot A = L_A Q(g, A)$ ) on  $U_T = \{x \in M : Z(T)(x) \neq 0\}$  (resp. on  $U_A = \{x \in M : Z(A)(x) \neq 0\}$ ), where  $L_T$  is a function defined on  $U_T$  (resp.  $L_A$  is a function defined on  $U_A$ ). The functions  $L_T$  and  $L_A$  are uniquely determined and called the *associated functions* of the pseudo-symmetric tensors  $T$  and  $A$  respectively ([8]).

A Riemannian manifold  $(M, g)$  is said to be *pseudo-symmetric* if its curvature tensor  $R$  is pseudo-symmetric [11]; then

$$(1) \quad R \cdot R = L_R Q(g, R)$$

on  $U_R$ . Any semi-symmetric manifold ( $R \cdot R = 0$ , [20]) is pseudo-symmetric. Examples of non-semi-symmetric pseudo-symmetric manifolds are given in [2], [3], [6] and [11].

$(M, g)$  is said to be *Weyl-pseudo-symmetric* if its Weyl conformal curvature tensor  $C$  is pseudo-symmetric [8]; then

$$(2) \quad R \cdot C = L_C Q(g, C)$$

on  $U_C$ . Any pseudo-symmetric manifold is Weyl-pseudo-symmetric. The converse fails in general (see Section 1). Note that  $U_C = \{x \in M : C(x) \neq 0\}$ .

$(M, g)$  is said to be *Ricci-pseudo-symmetric* if its Ricci tensor  $S$  is pseudo-symmetric ([14], [7]); then

$$(3) \quad R \cdot S = L_S Q(g, S)$$

on  $U_S$ . Of course, any pseudo-symmetric manifold is Ricci-pseudo-symmetric. The converse fails in general (see [14], [7]). The conditions (1) and (3) are equivalent on manifolds with vanishing Weyl conformal curvature tensor  $C$ . Namely, we have

LEMMA 1 ([2], Lemma 1.2, [13], Lemma 2). *If  $C$  vanishes at  $x \in M$  and  $n \geq 3$ , then at  $x$  the following three identities are equivalent to each other:*

$$\left( (n-2)\alpha + \frac{K}{n-1} \right) \left( S - \frac{k}{n}g \right) = S^2 - \frac{\text{tr}(S^2)}{n}g,$$

$$R \cdot S = \alpha Q(g, S), \quad R \cdot R = \alpha Q(g, R),$$

where  $\alpha \in \mathbb{R}$ ,  $S^2(X, Y) = S(\tilde{S}(X), Y)$ ,  $S(X, Y) = g(\tilde{S}(X), Y)$  and  $x, y \in \mathfrak{X}(M)$ .

LEMMA 2. (i) *If  $(M, g)$  is 3-dimensional then  $C$  vanishes identically.*

(ii) (cf. [15], p. 48) *Any generalized curvature tensor  $T$  at a point  $x$  of a 3-dimensional Riemannian manifold  $(M, g)$  satisfies*

$$T(X_1, X_2, X_3, X_4) = g(X_1, X_4)A(X_2, X_3) + g(X_2, X_3)A(X_1, X_4) \\ - g(X_1, X_3)A(X_2, X_4) - g(X_2, X_4)A(X_1, X_3),$$

for all  $X_i \in \mathfrak{X}(M)$ , where  $A$  is the  $(0, 2)$ -tensor defined by

$$A(X_1, X_2) = \text{Ricc}(T)(X_1, X_2) - \frac{K(T)}{4}g(X_1, X_2),$$

$\text{Ricc}(T)$  and  $K(T)$  being the Ricci tensor and the scalar curvature of  $T$  respectively.

(iii) *Let  $A$  be a symmetric  $(0, 2)$ -tensor on a 2-dimensional Riemannian manifold  $(M, g)$ . Then*

$$g(X_1, X_4)A(X_2, X_3) + g(X_2, X_3)A(X_1, X_4) - g(X_2, X_4)A(X_1, X_3) \\ - g(X_1, X_3)A(X_2, X_4) = \text{tr}(A)G(X_1, X_2, X_3, X_4)$$

on  $M$ .

LEMMA 3. *Let  $A$  and  $B$  be non-zero symmetric  $(0, 2)$ -tensors at a point  $x$  of a manifold  $(M, g)$ . If  $Q(A, B) = 0$  at  $x$  then  $A = \lambda B$ ,  $\lambda \in \mathbb{R} - \{0\}$ , at  $x$ .*

The proof of this lemma was given in [9] (see the proof of Lemma 3.4).

**3. Warped products.** Let  $(M, \bar{g})$  and  $(N, \tilde{g})$ ,  $\dim M = p$ ,  $\dim N = n - p$ ,  $1 \leq p \leq n$ , be Riemannian manifolds covered by systems of charts  $\{\bar{V}; x^a\}$  and  $\{\tilde{V}; y^\alpha\}$  respectively. Let  $F$  be a positive smooth function on  $M$ . The *warped product*  $M \times_F N$  of  $(M, \bar{g})$  and  $(N, \tilde{g})$  (see [17], [1]) is the Cartesian product  $M \times N$  with the metric  $g = \bar{g} \oplus F\tilde{g}$  (more precisely,  $\bar{g} \oplus F\tilde{g} = \Pi_1^*\bar{g} + (F \circ \Pi_1)\Pi_2^*\tilde{g}$ ,  $\Pi_1 : M \times N \rightarrow M$  and  $\Pi_2 : M \times N \rightarrow N$

being the natural projections). Let  $\{\bar{V} \times \tilde{V}; u^1 = x^1, \dots, u^p = x^p, u^{p+1} = y^1, \dots, u^n = y^{n-p}\}$  be a product chart for  $M \times N$ . The local components of the metric  $\bar{g} \oplus F\tilde{g}$  with respect to this chart are

$$(4) \quad g_{rs} = \begin{cases} \bar{g}_{ab} & \text{if } r = a, s = b, \\ F\tilde{g}_{\alpha\beta} & \text{if } r = \alpha, s = \beta, \\ 0 & \text{otherwise.} \end{cases}$$

Here and below,  $a, b, c, d, e, f \in \{1, \dots, p\}$ ,  $\alpha, \beta, \gamma, \delta, \lambda, \mu \in \{p+1, \dots, n\}$  and  $r, s, t, u, v, w \in \{1, \dots, n\}$ . The local components of the tensors  $R$  and  $S$  of the metric  $\bar{g} \oplus F\tilde{g}$  which may not vanish identically are

$$(5) \quad R_{abcd} = \bar{R}_{abcd},$$

$$(6) \quad R_{\alpha ab\beta} = -\frac{1}{2}T_{ab}\tilde{g}_{\alpha\beta},$$

$$(7) \quad R_{\alpha\beta\gamma\delta} = F\tilde{R}_{\alpha\beta\gamma\delta} - \frac{\Delta F}{4}\tilde{G}_{\alpha\beta\gamma\delta},$$

$$(8) \quad S_{ab} = \bar{S}_{ab} - \frac{n-p}{2F}T_{ab},$$

$$(9) \quad S_{\alpha\beta} = \tilde{S}_{\alpha\beta} - \frac{1}{2}\left(\text{tr}(T) + \frac{(n-p-1)\Delta F}{2F}\right)\tilde{g}_{\alpha\beta},$$

$$(10) \quad T_{ab} = \nabla_b F_a - \frac{1}{2F}F_a F_b, \quad \text{tr}(T) = \bar{g}^{ab}T_{ab},$$

$$\Delta F = \bar{g}^{ab}F_a F_b, \quad F_a = \frac{\partial}{\partial x^a}(F).$$

The scalar curvature  $K$  of  $\bar{g} \oplus F\tilde{g}$  satisfies

$$(11) \quad K = \bar{K} + \frac{1}{F}\tilde{K} - \frac{n-p}{F}\left(\text{tr}(T) + \frac{(n-p-1)\Delta F}{4F}\right).$$

Using (5)–(9) and (11), we obtain the following relations for the local components  $C_{rstu}$  of the tensor  $C$  of  $\bar{g} \oplus F\tilde{g}$ :

$$(12) \quad C_{abcd} = \bar{R}_{abcd} - \frac{1}{n-2}(\bar{g}_{ad}\bar{S}_{bc} - \bar{g}_{ac}\bar{S}_{bd} + \bar{g}_{bc}\bar{S}_{ad} - \bar{g}_{bd}\bar{S}_{ac}) \\ + \frac{n-p}{2(n-2)F}(\bar{g}_{ad}T_{bc} - \bar{g}_{ac}T_{bd} + \bar{g}_{bc}T_{ad} - \bar{g}_{bd}T_{ac}) \\ + \frac{K}{(n-1)(n-2)}\bar{G}_{abcd},$$

$$(13) \quad C_{\alpha ab\beta} = -\frac{1}{n-2}\left(\frac{p-2}{2}T_{ab} + F\bar{S}_{ab}\right)\tilde{g}_{\alpha\beta} - \frac{1}{n-2}\bar{g}_{ab}\tilde{S}_{\alpha\beta} \\ + \frac{1}{(n-1)(n-2)} \\ \times \left(F\bar{K} + \tilde{K} - \frac{(n-2p+1)\text{tr}(T)}{2} + \frac{(p-1)(n-p-1)\Delta F}{4F}\right)\bar{g}_{ab}\tilde{g}_{\alpha\beta},$$

$$(14) \quad C_{\alpha\beta\gamma\delta} = F\tilde{R}_{\alpha\beta\gamma\delta} - \frac{F}{n-2}(\tilde{g}_{\alpha\delta}\tilde{S}_{\beta\gamma} - \tilde{g}_{\alpha\gamma}\tilde{S}_{\beta\delta} + \tilde{g}_{\beta\gamma}\tilde{S}_{\alpha\delta} - \tilde{g}_{\beta\delta}\tilde{S}_{\alpha\gamma}) \\ + FP\tilde{G}_{\alpha\beta\gamma\delta},$$

$$(15) \quad C_{abc\alpha} = C_{ab\alpha\beta} = C_{a\alpha\beta\gamma} = 0,$$

$$(16) \quad P = \frac{1}{n-2} \left( \frac{FK}{n-1} + \text{tr}(T) + \frac{(n-2p)\Delta F}{4F} \right).$$

LEMMA 4 ([6], Theorem 1). *The curvature tensor  $R$  of a warped product  $M \times_F N$  satisfies  $R \cdot R = LQ(g, R)$  if and only if*

$$(17) \quad (\bar{R} \cdot \bar{R})_{abcdef} = LQ(\bar{g}, \bar{R})_{abcdef},$$

$$(18) \quad H^f{}_d \bar{R}_{fabc} = \frac{1}{2F}(T_{ac}H_{bd} - T_{ab}H_{cd}),$$

$$(19) \quad H_{ad} \left( \tilde{R}_{\delta\alpha\beta\gamma} - \frac{\Delta F}{4F} \tilde{G}_{\delta\alpha\beta\gamma} \right) = -\frac{1}{2} T_{fd} H^f{}_a \tilde{G}_{\delta\alpha\beta\gamma},$$

$$(20) \quad (\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} = \left( LF + \frac{\Delta F}{4F} \right) Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu},$$

where

$$(21) \quad H_{ad} = \frac{1}{2} T_{ad} + FL\bar{g}_{ad}.$$

LEMMA 5 ([6], Corollary 1). *Let  $(M, \bar{g})$ ,  $\dim M \geq 2$  and  $(N, \tilde{g})$ ,  $\dim N \geq 2$ , be manifolds of constant curvature. The curvature tensor  $R$  of the warped product  $M \times_F N$  satisfies  $R \cdot R = LQ(g, R)$  if and only if*

$$(22) \quad \frac{2\bar{K}}{p(p-1)} (\bar{g}_{ab}H_{cd} - \bar{g}_{ac}H_{bd}) = \frac{1}{F}(T_{ac}H_{bd} - T_{ab}H_{cd}),$$

$$(23) \quad H_{ad} \left( \frac{\tilde{K}}{(n-p)(n-p-1)} - \frac{\Delta F}{4F} \right) = -\frac{1}{2} T_{fd} H^f{}_a,$$

Using (4)–(16), (21) and Lemma 2(iii), we obtain

LEMMA 6. *The only local components of the Weyl conformal curvature tensor  $C$  of a 4-dimensional warped product  $M \times_F N$  which are not identically zero are*

$$(24) \quad C_{\alpha 11\beta} = -\frac{1}{2} \bar{g}_{11} \left( \tilde{S}_{\alpha\beta} - \frac{\tilde{K}}{3} \tilde{g}_{\alpha\beta} \right),$$

$$(25) \quad C_{\alpha\beta\gamma\delta} = \frac{F}{2} (\tilde{g}_{\alpha\delta}\tilde{S}_{\beta\gamma} - \tilde{g}_{\alpha\gamma}\tilde{S}_{\beta\delta} + \tilde{g}_{\beta\gamma}\tilde{S}_{\alpha\delta} - \tilde{g}_{\beta\delta}\tilde{S}_{\alpha\gamma}) - \frac{F\tilde{K}}{3} \tilde{G}_{\alpha\beta\gamma\delta},$$

provided that  $\dim M = 1$ ;

$$(26) \quad C_{abcd} = \frac{P}{F} G_{abcd}, \quad P = \frac{1}{6} \left( F\bar{K} + \tilde{K} + \text{tr}(T) - \frac{\Delta F}{2F} \right),$$

$$(27) \quad C_{\alpha\beta\gamma\delta} = \frac{P}{F} G_{\alpha\beta\gamma\delta},$$

$$(28) \quad C_{\alpha ab\beta} = -\frac{P}{2F} G_{\alpha ab\beta},$$

provided that  $\dim M = 2$ ;

$$(29) \quad C_{4ab4} = -\frac{1}{2} \tilde{g}_{44} W_{ab}, \quad W_{ab} = V_{ab} - \frac{\text{tr}(V)}{3} \bar{g}_{ab}, \quad V_{ab} = F \bar{S}_{ab} + \frac{1}{2} T_{ab},$$

$$(30) \quad C_{abcd} = \frac{1}{2F} (\bar{g}_{ad} W_{bc} + \bar{g}_{bc} W_{ad} - \bar{g}_{ac} W_{bd} - \bar{g}_{bd} W_{ac}),$$

provided that  $\dim M = 3$ .

From the above lemma the following theorem follows immediately:

**THEOREM 1.** *Suppose  $\dim M \times_F N = 4$ .*

(i) *If  $\dim M = 1$ , then  $M \times_F N$  is conformally flat if and only if*

$$\tilde{S} = \frac{\tilde{K}}{3} \tilde{g}.$$

(ii) *If  $\dim M = 2$ , then  $M \times_F N$  is conformally flat if and only if*

$$\text{tr}(T) = -\frac{FK}{3}.$$

(iii) *If  $\dim M = 3$ , then  $M \times_F N$  is conformally flat if and only if*

$$F\bar{S} + \frac{T}{2} = \frac{1}{3} \left( F\bar{K} + \frac{\text{tr}(T)}{2} \right) \bar{g}.$$

**Remark 2.** (i) Necessary and sufficient conditions for  $M \times_F N$ ,  $\dim M \times_F N \geq 4$  and  $\dim N \geq 2$ , to be conformally flat are given in [19] (Theorem 1).

(ii) An example of a 4-dimensional conformally flat warped product  $M \times_F N$ ,  $\dim N = 1$ , is described in [2] (Lemma 4.3). The manifold  $(M, \bar{g})$  considered in that example is non-semi-symmetric, conformally flat and pseudo-symmetric, but  $M \times_F N$  is not pseudo-symmetric.

(iii) The assertion (iii) of Theorem 1 can be easily generalized (by making use of (12)–(16)) as follows: The manifold  $M \times_F N$ ,  $\dim M = n - 1$ ,  $n \geq 4$ , is conformally flat if and only if

$$\bar{C} = 0 \quad \text{and} \quad F\bar{S} + \frac{(n-3)T}{2} = \frac{1}{n-1} \left( F\bar{K} + \frac{(n-3)\text{tr}(T)}{2} \right) \bar{g}$$

on  $M$ .

Another consequence of Lemma 6 is

**THEOREM 2.** *Suppose  $\dim M \times_F N = 4$  and  $\dim M = 2$ . Then  $C \cdot C = -\frac{P}{2F} Q(g, C)$  on  $M \times_F N$ .*

The tensor  $C \cdot C$  is defined analogously to the tensor  $R \cdot T$  in Section 2. Riemannian manifolds satisfying the condition  $C \cdot C = 0$  were considered in [4] (see also [13], Corollary 1).

LEMMA 7 ([7], Theorem 1). *The Ricci tensor  $S$  of  $M \times_F N$  satisfies  $R \cdot S = LQ(g, S)$  if and only if*

$$(31) \quad (\bar{R} \cdot \bar{S})_{abcd} - LQ(\bar{g}, \bar{S})_{abcd} = \frac{n-p}{F} ((\bar{R} \cdot H)_{abcd} - LQ(\bar{g}, H)_{abcd}),$$

$$(32) \quad H_{ab} \left( \tilde{S}_{\alpha\beta} - \frac{1}{2F} \left( \text{tr}(T) + \frac{(n-p-1)\Delta F}{2F} \right) g_{\alpha\beta} \right) \\ = H_{cb} \left( \bar{S}^c{}_a - \frac{n-p}{2F} T^c{}_a \right) g_{\alpha\beta},$$

$$(33) \quad (\tilde{R} \cdot \tilde{S})_{\alpha\beta\gamma\delta} = \left( LF + \frac{\Delta F}{4F} \right) Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta}.$$

LEMMA 8 ([7], Corollary 1). *Let  $(M, \bar{g})$ ,  $\dim M \geq 2$  and  $(N, \tilde{g})$ ,  $\dim N \geq 2$ , be Einstein manifolds. Then the Ricci tensor  $S$  of  $M \times_F N$  satisfies  $R \cdot S = LQ(g, S)$  if and only if*

$$(34) \quad (\bar{R} \cdot H)_{abcd} = LQ(\bar{g}, H)_{abcd},$$

$$(35) \quad \frac{F}{n-p} \left( \frac{\bar{K}}{p} - \frac{\tilde{K}}{(n-p)F} + (n-p)L \right. \\ \left. + \frac{1}{2F} \left( \text{tr}(T) + \frac{(n-p-1)\Delta F}{2F} \right) \right) H_{ab} = H_{ac} H^c{}_b.$$

#### 4. Not conformally flat 4-dimensional warped products satisfying $R \cdot C = LQ(g, C)$

PROPOSITION 1. *Suppose  $R \cdot C = LQ(g, C)$  on  $M \times_F N$ , where  $\dim M = 1$  and  $\dim N = 3$ . If  $C$  is non-zero at  $x \in M \times_F N$  then  $R \cdot R = LQ(g, R)$  at  $x$ .*

Proof. From (2) we have

$$(R \cdot C)_{\alpha\beta\gamma\delta\lambda\mu} = LQ(g, C)_{\alpha\beta\gamma\delta\lambda\mu},$$

whence, by (7), it follows that

$$(\tilde{R} \cdot C)_{\alpha\beta\gamma\delta\lambda\mu} = \left( LF + \frac{\Delta F}{4F} \right) Q(\tilde{g}, C)_{\alpha\beta\gamma\delta\lambda\mu}.$$

This, by an application of (25) and contraction with  $\tilde{g}^{\beta\gamma}$ , yields

$$(\tilde{R} \cdot \tilde{S})_{\alpha\delta\lambda\mu} = \left( LF + \frac{\Delta F}{4F} \right) Q(\tilde{g}, \tilde{S})_{\alpha\delta\lambda\mu},$$

which, in view of Lemma 2(i) and Lemma 1, implies (20). Further, the relation  $(R \cdot C)_{1\alpha\beta\gamma1\delta} = LQ(g, C)_{1\alpha\beta\gamma1\delta}$ , in virtue of (6), (15), (24) and (21), turns into

$$H_{11} \left( \frac{1}{F} C_{\delta\alpha\beta\gamma} - \frac{\tilde{K}}{6} \tilde{G}_{\delta\alpha\beta\gamma} + \frac{1}{2} (\tilde{g}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{g}_{\beta\delta} \tilde{S}_{\alpha\gamma}) \right) = 0.$$

Applying (25) and contracting the resulting equality with  $\tilde{g}^{\gamma\delta}$ , we get  $H_{11}(\tilde{S}_{\alpha\beta} - \frac{\tilde{K}}{3} \tilde{g}_{\alpha\beta}) = 0$ , which, by (24), (25) and the assumption  $C(x) \neq 0$ , gives  $H_{11}(x) = 0$ . Now Lemma 4 completes the proof.

**PROPOSITION 2.** *Suppose  $R \cdot C = LQ(g, C)$  on  $M \times_F N$ , where  $\dim M = \dim N = 2$ . If  $C$  is non-zero at  $x \in M \times_F N$  then  $R \cdot R = LQ(g, R)$  and  $H = 0$  at  $x$ .*

*Proof.* The relation  $(R \cdot C)_{a\alpha\beta\gamma d\delta} = LQ(g, C)_{a\alpha\beta\gamma d\delta}$ , by making use of (15), (27), (28), (6) and (21), gives  $PH = 0$ . Since  $C(x) \neq 0$ , it follows that  $H(x) = 0$ . But this, in view of Lemma 5, completes the proof.

**PROPOSITION 3.** *Suppose  $R \cdot C = LQ(g, C)$  on  $M \times_F N$ , where  $\dim M = 3$  and  $\dim N = 1$ . If  $C$  is non-zero at  $x \in M \times_F N$  then  $R \cdot R = LQ(g, R)$  at  $x$ .*

*Proof.* From the equality  $(R \cdot C)_{4ab4cd} = LQ(g, C)_{4ab4cd}$ , by making use of (15) and (29), it follows that

$$(36) \quad (\bar{R} \cdot W)_{abcd} = LQ(\bar{g}, W)_{abcd}.$$

Furthermore, the equality  $(R \cdot C)_{4abcd4} = LQ(g, C)_{4abcd4}$ , by an application of (6), (15), (29) and (21), yields

$$(37) \quad H^e{}_d C_{eabc} = \frac{1}{2F} (H_{bd} W_{ac} - H_{cd} W_{ab}),$$

whence, by (30), we get

$$(38) \quad \bar{g}_{ab} H^e{}_d W_{ec} - \bar{g}_{ac} H^e{}_d W_{eb} + 2(W_{ab} H_{cd} - W_{ac} H_{bd}) = 0.$$

Contracting this with  $\bar{g}^{ad}$  and  $\bar{g}^{cd}$  respectively, we obtain

$$(39) \quad H^e{}_a W_{eb} = H^e{}_b W_{ea},$$

$$(40) \quad H^e{}_b W_{ea} = \frac{2}{3} \tau W_{ab} + \frac{1}{3} \rho \bar{g}_{ab}, \quad \rho = H^{ef} W_{ef}, \quad \tau = \text{tr}(H),$$

respectively. Now (38) takes the form

$$(41) \quad (W_{ab} H_{cd} - W_{ac} H_{bd}) + \frac{1}{3} \tau (\bar{g}_{ab} W_{cd} - \bar{g}_{ac} W_{bd}) + \frac{1}{6} \rho \bar{G}_{dabc} = 0.$$

Transvecting the above equality with  $H^{ab}$  and using (39) and (40) we find

$$(42) \quad H_{cd} = \frac{2}{3} \tau^2 W_{cd} + \frac{1}{3} \tau \rho \bar{g}_{cd}.$$

From this we have  $\rho\bar{R} \cdot H = \frac{2}{3}\tau^2\bar{R} \cdot W$  and, by (36),

$$\bar{R} \cdot H = \frac{2}{3}\tau^2 LQ(\bar{g}, W).$$

Applying (42) in the last equality, we get

$$(43) \quad \rho(\bar{R} \cdot H - LQ(\bar{g}, H)) = 0,$$

which, by (21), implies

$$(44) \quad \rho(\bar{R} \cdot T - LQ(\bar{g}, T)) = 0.$$

Thus (36), in virtue of (29) and (44), turns into

$$(45) \quad \rho(\bar{R} \cdot \bar{S} - LQ(\bar{g}, \bar{S})) = 0.$$

We have now the two possibilities: (a)  $\rho(x) \neq 0$  and (b)  $\rho(x) = 0$ .

(a) In this case  $\bar{R} \cdot \bar{S} = LQ(\bar{g}, \bar{S})$ . Thus, in view of Lemma 1, (17) holds at  $x$ . Now, we prove that (18) also holds at  $x$ . (42) shows that this is trivial if  $\tau(x) = 0$ . Suppose that  $\tau(x) \neq 0$ . First of all, we note that from (41), by transvection with  $H^a_f$  and an application of (40) and (42) the following relation can be obtained at  $x$ :

$$(46) \quad H_{bf}H_{cd} - H_{cf}H_{bd} = 0,$$

whence

$$(47) \quad H_{bf}H^f_d = \tau H_{bd}.$$

Next, transvecting the equality  $\bar{C}_{eabc} = 0$  with  $H^e_d$  we obtain

$$(48) \quad H^e_d \bar{R}_{eabc} = H_{cd} \bar{S}_{ab} - H_{bd} \bar{S}_{ac} + \bar{g}_{ab} \bar{S}_{ec} H^e_d - \bar{g}_{ac} \bar{S}_{eb} H^e_d \\ - \frac{\bar{K}}{2} (\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}).$$

The formula (42), by making use of (29), can be rewritten in the form

$$(49) \quad F \bar{S}_{ab} = \left( \frac{3}{2} \frac{1}{\tau^2} \rho - 1 \right) H_{ab} + \left( FL - \frac{1}{2\tau} \rho \right) \bar{g}_{ab},$$

which, by transvection with  $H^b_d$  and the use of (47), yields

$$F \bar{S}_{ea} H^e_d = \left( \frac{1}{\tau} \rho - \tau + FL \right) H_{ad}.$$

Applying the last two equalities in (48) we get

$$H^e_d \bar{R}_{eabc} = \frac{1}{F} \left( \frac{3}{2\tau^2} \rho - 1 \right) (H_{cd} H_{ab} - H_{bd} H_{ac}) \\ + \frac{1}{F} \left( \frac{1}{2\tau} \rho - \tau + 2FL - \frac{\bar{K}F}{2} \right) (\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}),$$

which, by (46), reduces to

$$(50) \quad H^e {}_d \bar{R}_{eabc} = \frac{1}{F} \left( \frac{1}{2\tau} \rho - \tau + 2FL - \frac{\bar{K}F}{2} \right) (\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}).$$

This, by the Ricci identity, yields

$$(51) \quad \bar{R} \cdot H = \frac{1}{F} \left( \frac{1}{2\tau} \rho - \tau + 2FL - \frac{\bar{K}F}{2} \right) Q(\bar{g}, H).$$

Comparing (51) with (43) we obtain

$$\left( \frac{1}{F} \left( \frac{1}{2\tau} \rho - \tau + 2FL - \frac{\bar{K}F}{2} \right) - L \right) Q(\bar{g}, H) = 0,$$

whence

$$(52) \quad \left( \frac{1}{F} \left( \frac{1}{2\tau} \rho - \tau + 2FL - \frac{\bar{K}F}{2} \right) - L \right) \left( H - \frac{\tau}{3} \bar{g} \right) = 0.$$

If  $(H - (\tau/3)\bar{g})(x) = 0$ , then (49) gives  $\bar{S} = (\bar{K}/3)\bar{g}$ . But this, by (29), gives  $W = 0$  and, consequently,  $C(x) = 0$ , which is a contradiction. So  $(H - (\tau/3)\bar{g})(x) \neq 0$ . Applying now (52) in (50) we get

$$(53) \quad H^e {}_d \bar{R}_{eabc} = L(\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}).$$

Note that (46) can be expressed in the following form:

$$L(\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}) = \frac{1}{2F} (T_{ac} H_{bd} - T_{ab} H_{cd}).$$

Thus (53) turns into (18).

(b) Since  $\rho$  vanishes at  $x$ , the formula (42) takes the form  $\tau W = 0$ , whence, by (29), (30) and the assumption  $C(x) \neq 0$ , we obtain the equality

$$(54) \quad \tau = \frac{\text{tr}(T)}{2} + 3FL = 0$$

at this point. The tensor  $W$  now takes the form

$$(55) \quad W_{ab} = F\bar{S}_{ab} - \frac{F\bar{K}}{3}\bar{g}_{ab} + H_{ab}.$$

The formula (40), by (54), gives

$$(56) \quad H^e {}_a W_{eb} = 0.$$

Thus (38) turns into

$$(57) \quad W_{ab} H_{cd} - W_{ac} H_{bd} = 0,$$

which can be rewritten in the form

$$(58) \quad F(\bar{S}_{ab} H_{cd} - \bar{S}_{ac} H_{bd}) = H_{ac} H_{bd} - H_{ab} H_{cd} + \frac{F\bar{K}}{3} (\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}).$$

From (57), by transvection with  $H^c_e$  and making use of (56), we find

$$(59) \quad H_{cd}H^c_e = 0.$$

Further, transvecting (55) with  $H^a_d$  and applying (56) and (59) we get

$$(60) \quad H^c_a \bar{S}_{cd} = \frac{\bar{K}}{3} H_{ad}.$$

Next, transvecting (58) with  $\bar{S}_e^b$  and using (60) we obtain

$$\begin{aligned} F \left( \bar{S}_e^b \bar{S}_{ab} - \frac{\bar{K}}{3} \bar{S}_{ea} \right) H_{cd} \\ = \frac{\bar{K}}{3} (H_{ac} H_{ed} - H_{ea} H_{cd}) + \frac{F\bar{K}}{3} \left( \bar{S}_{ac} - \frac{\bar{K}}{3} \bar{g}_{ac} \right) H_{ed}, \end{aligned}$$

which, by (58), turns into

$$\begin{aligned} \left( \bar{S}^2 - \frac{2\bar{K}\bar{S}}{3} + \frac{\bar{K}^2}{9} \bar{g} \right) H = 0, \quad \text{or} \\ \left( \left( \bar{S}^2 - \frac{\text{tr}(\bar{S}^2)}{3} \bar{g} \right) - \left( \frac{\bar{K}}{6} + \frac{\bar{K}}{2} \right) \left( \bar{S} - \frac{\bar{K}}{3} \bar{g} \right) + \frac{1}{3} \left( \text{tr}(\bar{S}^2) - \frac{\bar{K}^2}{3} \right) \bar{g} \right) H = 0 \end{aligned}$$

and

$$(61) \quad \left( \left( \bar{S}^2 - \frac{\text{tr}(\bar{S}^2)}{3} \bar{g} \right) - \left( \frac{\bar{K}}{6} + \frac{\bar{K}}{2} \right) \left( \bar{S} - \frac{\bar{K}}{3} \bar{g} \right) \right) H = 0.$$

Suppose that  $H(x) = 0$ . Then, of course, (18) holds at  $x$ . The formula (36) turns into  $\bar{R} \cdot \bar{S} = LQ(\bar{g}, \bar{S})$ . But this, in view of Lemma 2(i) and Lemma 1, implies (17). Consider now the case  $H(x) \neq 0$ . Then (61) gives

$$\bar{S}^2 - \frac{\text{tr}(\bar{S}^2)}{3} \bar{g} = \left( \frac{\bar{K}}{6} + \frac{\bar{K}}{2} \right) \left( \bar{S} - \frac{\bar{K}}{3} \bar{g} \right).$$

But this, in view of Lemma 2(i) and Lemma 1, yields  $\bar{R} \cdot \bar{R} = LQ(\bar{g}, \bar{R})$  and

$$(62) \quad L = \bar{K}/6.$$

Thus the condition (17) is fulfilled. Finally, the identity (48), in virtue of (60), gives

$$H^e_d \bar{R}_{eabc} = H_{cd} \bar{S}_{ab} - H_{bd} \bar{S}_{ac} - \frac{\bar{K}}{6} (\bar{g}_{ab} H_{cd} - \bar{g}_{ac} H_{bd}),$$

which, by (58), turns into

$$FH^e_d \bar{R}_{eabc} = H_{ac} H_{bd} - H_{ab} H_{cd} + \frac{F\bar{K}}{6} (\bar{g}_{ab} H_{ac} - \bar{g}_{ac} H_{bd}).$$

This, by making use of (61) and (21), leads to (18). Our proposition is thus proved.

Combining Propositions 1–3 we obtain

**THEOREM 3.** *Suppose  $\dim M \times_F N = 4$ . If at a point of  $M \times_F N$  the tensor  $C$  is non-zero, then the relations  $R \cdot C = LQ(g, C)$  and  $R \cdot R = LQ(g, R)$  are equivalent at this point.*

The following corollary is a consequence of Theorem 2.

**COROLLARY 1.** *Let  $M \times_F N$  be an analytic not conformally flat 4-dimensional warped product. Then the relations  $R \cdot C = LQ(g, C)$  and  $R \cdot R = LQ(g, R)$  are equivalent on  $M \times_F N$ .*

**5. Non-Einstein 4-dimensional warped product satisfying  $R \cdot S = LQ(g, S)$**

**PROPOSITION 4.** *Suppose  $R \cdot S = LQ(g, S)$  on  $M \times_F N$ , where  $\dim M = 1$  and  $\dim N = 3$ . If  $S - (K/4)g$  is non-zero at  $x \in M \times_F N$ , then  $R \cdot R = LQ(g, R)$  at  $x$ .*

**Proof.** The equality (33), in view of Lemma 2(i) and Lemma 1, turns into (20). Further, (32) yields

$$H_{11} \left( \tilde{S}_{\alpha\beta} - \left( \frac{1}{2F} \Delta F - \text{tr}(T) \right) \tilde{g}_{\alpha\beta} \right) = 0,$$

whence  $H(\tilde{S} - (\tilde{K}/3)\tilde{g}) = 0$ . If  $H = 0$ , then (18) and (19) are satisfied and Lemma 4 completes the proof. If  $\tilde{S} - (\tilde{K}/3)\tilde{g} = 0$ , then  $C = 0$ , and our assertion, by Lemma 1, is also true.

**PROPOSITION 5.** *Suppose  $R \cdot S = LQ(g, S)$  on  $M \times_F N$ , where  $\dim M = \dim N = 2$ . If  $S - (K/4)g$  is non-zero at  $x \in M \times_F N$  then  $R \cdot R = LQ(g, R)$  at  $x$ .*

**Proof.** The relations (34) and (35) take the forms

$$(63) \quad \left( L - \frac{\bar{K}}{2} \right) \left( H - \frac{\text{tr}(H)}{2} \bar{g} \right) = 0,$$

$$(64) \quad H_{ca} H^c_b = \rho H_{ab}$$

respectively, where

$$(65) \quad \rho = \frac{F}{2} \left( \frac{\bar{K}}{2} - \frac{\tilde{K}}{2F} + 2L + \frac{\text{tr}(T)}{2F} + \frac{1}{4F^2} \Delta F \right).$$

In view of Lemma 2(iii),  $H$  satisfies the following identity at  $x$ :

$$(66) \quad \bar{g}_{bc} H_{ad} + \bar{g}_{ad} H_{bc} - \bar{g}_{ac} H_{bd} - \bar{g}_{bd} H_{ac} = \text{tr}(H)(\bar{g}_{ad} \bar{g}_{bc} - \bar{g}_{ac} \bar{g}_{bd}).$$

Transvecting this with  $H^b_f$  and using (64) we get

$$(67) \quad H_{cf} H_{ad} - H_{df} H_{ac} = (\text{tr}(H) - \rho)(\bar{g}_{ad} H_{cf} - \bar{g}_{ac} H_{df}),$$

whence, by transvection with  $H^{cf}$  and making use of (64) we obtain

$$(68) \quad \rho(\operatorname{tr}(H) - \rho) \left( H - \frac{\operatorname{tr}(H)}{2} \bar{g} \right) = 0.$$

Consider two possibilities:

$$(a) \quad \left( H - \frac{\operatorname{tr}(H)}{2} \bar{g} \right) (x) \neq 0,$$

$$(b) \quad \left( H - \frac{\operatorname{tr}(H)}{2} \bar{g} \right) (x) = 0.$$

(a) In this case we have

$$(69) \quad L = \bar{K}/2,$$

$$(70) \quad \rho(\operatorname{tr}(H) - \rho) = 0.$$

(a1) Suppose additionally that  $\rho(x) = 0$ . Then from (64) and (67) it follows that  $\operatorname{tr}(H)H_{ae} = 0$  and, in consequence,

$$(71) \quad \operatorname{tr}(H) = 0.$$

Now (67), by (21), (69) and (71), turns into (22). Further, (71) gives  $L = -\operatorname{tr}(T)/(4F)$ . Thus, from (65) we obtain  $FL = \tilde{K}/2 - \Delta F/(4F)$ . But this, together with (64) and (21), yields (23). Now Lemma 5 completes the proof.

(a2) Let  $\rho(x) \neq 0$ . Then, in virtue of (70), we get

$$(72) \quad \operatorname{tr}(H) = \rho.$$

Applying this, (21) and (69) in (67) we obtain (22). Further, (72) yields

$$\frac{\operatorname{tr}(T)}{4} + FL = \frac{FL}{2} - \frac{\tilde{K}}{4} + \frac{\Delta F}{8F},$$

whence

$$\operatorname{tr}(H) = FL - \frac{\tilde{K}}{2} + \frac{\Delta F}{4F}.$$

But this, together with (72) and (64), yields (23). Now Lemma 5 completes the proof in this case.

(b) Now (64) takes the form

$$(73) \quad \operatorname{tr}(H) \left( \frac{\operatorname{tr}(H)}{2} - \rho \right) = 0.$$

If  $\operatorname{tr}(H) = 0$ , then also  $H = 0$  and Lemma 5 completes the proof. If  $\operatorname{tr}(H) \neq 0$ , then (73) gives  $\operatorname{tr}(H) = 2\rho$ , whence we get  $F\bar{K}/2 = \tilde{K}/2 - \Delta F/(4F)$ . Moreover, the equality  $H = \frac{1}{2}\operatorname{tr}(H)\bar{g}$  yields  $T = \frac{1}{2}\operatorname{tr}(T)\bar{g}$ . Now (8) and (9) lead to  $S = (K/4)g$ , a contradiction. This completes the proof.

PROPOSITION 6. *Suppose  $R \cdot S = LQ(g, S)$  on  $M \times_F N$ , where  $\dim M = 3$  and  $\dim N = 1$ . If  $S - (K/4)g$  and  $C$  are non-zero at  $x \in M \times_F N$ , then  $R \cdot R = LQ(g, R)$  at  $x$ .*

Proof. We consider two cases:

- (a)  $(\bar{S} - (\bar{K}/3)\bar{g})(x) = 0$ ,
- (b)  $(\bar{S} - (\bar{K}/3)\bar{g})(x) \neq 0$ .

(a) The relation (31) turns into  $(L - \bar{K}/6)Q(\bar{g}, H) = 0$ , which, by Remark 1, yields

$$(74) \quad \left(L - \frac{\bar{K}}{6}\right) \left(H - \frac{\text{tr}(H)}{3}\bar{g}\right) = 0.$$

If  $H - (\text{tr}(H)/3)\bar{g} = 0$ , then (29) and (30) yield  $C = 0$ , a contradiction. If  $H - (\text{tr}(H)/3)\bar{g} \neq 0$ , then from (74) we obtain  $L = \bar{K}/6$ . Further, (32) takes the form  $H_{ca}H^c_b = \frac{1}{2}(F\bar{K} + \text{tr}(T))H_{ab}$ . Let  $B$  be a  $(0, 4)$ -tensor with local components  $B_{abcd} = H_{ad}H_{bc} - H_{ac}H_{bd}$ . We note that  $\text{Ricc}(B) = 0$  and  $K(B) = 0$ . Thus Lemma 2(ii) implies  $B_{abcd} = 0$ , whence

$$\frac{\bar{K}}{3}(\bar{g}_{ab}H_{cd} - \bar{g}_{ac}H_{bd}) = \frac{1}{F}(T_{ac}H_{bd} - T_{ab}H_{cd}).$$

But this turns into (18). Now Lemma 4 completes the proof.

(b) We rewrite (32) in the form

$$(75) \quad \begin{aligned} H^2_{ab} &= \left(FL + \frac{\text{tr}(T)}{2}\right)H_{ab} + FA_{ab}, \\ H^2_{ab} &= H_{ca}H^c_b, \quad A_{ab} = \bar{S}_{ca}H^c_b. \end{aligned}$$

Now, transvecting the identity  $\bar{C}_{ebcd} = 0$  with  $H_a^e$  and  $\bar{S}_a^e$  respectively, we obtain

$$(76) \quad \begin{aligned} H_a^e \bar{R}_{ebcd} &= H_{ad}\bar{S}_{bc} - H_{ac}\bar{S}_{bd} + \bar{g}_{bc}A_{ad} \\ &\quad - \bar{g}_{bd}A_{ac} - \frac{\bar{K}}{2}(\bar{g}_{bc}H_{ad} - \bar{g}_{bd}H_{ac}), \\ \bar{S}_a^e \bar{R}_{ebcd} &= \bar{S}_{ad}\bar{S}_{bc} - \bar{S}_{ac}\bar{S}_{bd} + \bar{g}_{bc}\bar{S}^2_{ad} \\ &\quad - \bar{g}_{bd}\bar{S}^2_{ac} - \frac{\bar{K}}{2}(\bar{g}_{bc}\bar{S}_{ad} - \bar{g}_{bd}\bar{S}_{ac}) \end{aligned}$$

respectively, where  $\bar{S}^2_{ad} = \bar{S}_{ea}\bar{S}^e_d$ . From the last two relations, by symmetrization in  $a, b$  and making use of the Ricci identity and (31), it follows that

$$(77) \quad Q \left( \bar{g}, F\bar{S}^2 - F \left( \frac{\bar{K}}{2} + L \right) \bar{S} - A + \left( \frac{\bar{K}}{2} + L \right) H \right)_{abcd} = -Q(H, \bar{S})_{abcd}.$$

Contracting (77) with  $\bar{g}^{ad}$  we get

$$(78) \quad F\bar{S}^2 = \frac{F \operatorname{tr}(\bar{S}^2)}{3}\bar{g} + F\left(\frac{\bar{K}}{2} + L\right)\left(\bar{S} - \frac{\bar{K}}{3}\bar{g}\right) - \frac{\operatorname{tr}(H)}{3}\bar{S} + \frac{\bar{K}}{3}H \\ + \left(A - \frac{\operatorname{tr}(A)}{3}\bar{g}\right) - \left(\frac{\bar{K}}{2} + L\right)\left(H - \frac{\operatorname{tr}(H)}{3}\bar{g}\right).$$

Substituting this in (77) we find

$$(79) \quad Q\left(\bar{S} - \frac{\bar{K}}{3}\bar{g}, H - \frac{\operatorname{tr}(H)}{3}\bar{g}\right) = 0.$$

We may assume that  $H - (\operatorname{tr}(H)/3)\bar{g} \neq 0$ . Of course, if  $H = 0$ , then  $R \cdot R = LQ(g, R)$  at  $x$ . If  $H \neq 0$  and  $H - (\operatorname{tr}(H)/3)\bar{g} = 0$ , then (75) yields  $\bar{S} = (\bar{K}/3)\bar{g}$ , a contradiction. Now (79), in view of Lemma 3, implies

$$(80) \quad \bar{S} - \frac{\bar{K}}{3}\bar{g} = \lambda\left(H - \frac{\operatorname{tr}(H)}{3}\bar{g}\right), \quad \lambda \in \mathbb{R} - \{0\}.$$

Thus (31) yields  $(\lambda F - 1)(\bar{R} \cdot \bar{S} - LQ(\bar{g}, \bar{S})) = 0$ . Assume that

$$(81) \quad \lambda F = 1;$$

then (80) gives  $A = \frac{1}{3}(\bar{K} - \operatorname{tr}(H)/F)H + H^2/F$ . Substituting this into (75) we obtain  $(\bar{K} + \operatorname{tr}(T)/F)H = 0$ , whence  $K + \operatorname{tr}(T)/F = 0$ . But the last relation, together with (80), (81), (8), (9) and (11), leads to  $S = (K/4)g$ , a contradiction. Thus  $\lambda F - 1 \neq 0$  and

$$(82) \quad \bar{R} \cdot \bar{S} = LQ(\bar{g}, \bar{S}).$$

Now, in view of Lemma 1, we obtain (17). Furthermore, the equality (31), in virtue of (82), (76)–(78), turns into

$$Q\left(\bar{g}, \left(L + \frac{\bar{K}}{6}\right)H + \frac{\operatorname{tr}(H)}{3}\bar{S} - A\right) = 0,$$

whence, by (80), it follows that

$$A = \frac{\operatorname{tr}(A)}{3}\bar{g} + \left(\frac{\lambda \operatorname{tr}(H)}{3} + L + \frac{\bar{K}}{6}\right)\left(H - \frac{\operatorname{tr}(H)}{3}\bar{g}\right).$$

On the other hand, combining (75) and (80), we find

$$(83) \quad \left(\frac{1}{\lambda} - F\right)A = \frac{1}{3}\left(\operatorname{tr}(T) + \frac{\bar{K}}{\lambda}\right)H.$$

But the last two relations yield

$$(\lambda \operatorname{tr}(T) - 6L(1 - \lambda F) + \bar{K})(1 + \lambda F)\left(H - \frac{\operatorname{tr}(H)}{3}\bar{g}\right) = 0,$$

whence

$$(\lambda \operatorname{tr}(T) - 6L(1 - \lambda F) + \bar{K})(1 + \lambda F) = 0.$$

Suppose that  $1 + \lambda F = 0$ . Then (80), (29) and (30) give  $C = 0$ , a contradiction. Thus we have  $\lambda \operatorname{tr}(T) - 6L(1 - \lambda F) + \bar{K} = 0$ , or, equivalently,

$$(84) \quad \frac{1}{\lambda} \left( L - \frac{\bar{K}}{6} \right) = \frac{\operatorname{tr}(H)}{3}.$$

Thus (83) takes the form

$$(85) \quad A = 2LH.$$

Substituting this into (75) we obtain  $H^2 = \operatorname{tr}(H)H$ . Let  $B$  be a  $(0, 4)$ -tensor with local components  $B_{abcd} = H_{ad}H_{bc} - H_{ac}H_{bd}$ . Evidently,  $\operatorname{Ricc}(B) = 0$  and  $K(B) = 0$ . Thus, Lemma 2(ii) implies

$$(86) \quad B_{abcd} = 0.$$

From (80), by (86), we obtain

$$H_{ad}\bar{S}_{bc} - H_{ac}\bar{S}_{bd} = \frac{1}{3}(\bar{K} - \lambda \operatorname{tr}(H))(\bar{g}_{bc}H_{ad} - \bar{g}_{bd}H_{ac}).$$

Substituting this and (85) into (76) we find

$$H_a{}^e \bar{R}_{ebcd} = \left( 2L - \frac{\bar{K}}{6} - \frac{\lambda \operatorname{tr}(H)}{3} \right) (\bar{g}_{bc}H_{ad} - \bar{g}_{bd}H_{ac}),$$

whence, by (84), we get  $H_a{}^e \bar{R}_{ebcd} = L(\bar{g}_{bc}H_{ad} - \bar{g}_{bd}H_{ac})$ . But this, by (86), turns into (18). Now Lemma 4 completes the proof.

Combining Propositions 4–6 and Lemma 1 we obtain

**THEOREM 4.** *Suppose  $\dim M \times_F N = 4$ . If  $S - (K/4)g$  is non-zero at a point of  $M \times_F N$ , then the relations  $R \cdot S = LQ(g, S)$  and  $R \cdot R = LQ(g, R)$  are equivalent at this point.*

This theorem yields

**COROLLARY 2.** *Let  $M \times_F N$  be an analytic non-Einstein 4-dimensional warped product. Then the relations  $R \cdot S = LQ(g, S)$  and  $R \cdot R = LQ(g, R)$  are equivalent on  $M \times_F N$ .*

#### REFERENCES

- [1] R. L. Bishop and B. O'Neill, *Manifolds of negative curvature*, Trans. Amer. Math. Soc. 145 (1969), 1–49.
- [2] J. Deprez, R. Deszcz and L. Verstraelen, *Examples of pseudo-symmetric conformally flat warped products*, Chinese J. Math. 17 (1989), 51–65.
- [3] —, —, —, *Pseudo-symmetry curvature conditions on hypersurfaces of Euclidean spaces and on Kählerian manifolds*, Ann. Fac. Sci. Univ. Paul Sabatier Toulouse 9 (1988), 183–192.

- [4] J. Deprez, P. Verheyen and L. Verstraelen, *Characterization of conformally flat hypersurfaces*, Czechoslovak Math. J. 35 (110) (1985), 140–145.
- [5] A. Derdziński, *Exemples de métriques de Kaehler et d'Einstein autoduales sur le plan complexe*, in: Géométrie riemannienne en dimension 4 (Séminaire Arthur Besse 1978/79), Cedic/Fernand Nathan, Paris 1981, 334–346.
- [6] R. Deszcz, *On pseudo-symmetric warped product manifolds*, to appear.
- [7] —, *On Ricci-pseudo-symmetric warped products*, Demonstratio Math. 22 (1989), 1053–1065.
- [8] —, *On pseudo-symmetric totally umbilical submanifolds of Riemannian manifolds admitting some types of generalized curvature tensors*, Zeszyty Naukowe Politech. Śląsk., in print.
- [9] —, *On conformally flat Riemannian manifolds satisfying certain curvature conditions*, Tensor (N.S.), in print.
- [10] —, *Examples of four-dimensional Riemannian manifolds satisfying some pseudo-symmetry curvature conditions*, in: Geometry and Topology of Submanifolds, II, Avignon, May 1988, World Sci. Publ., Singapore 1990, 134–143.
- [11] R. Deszcz and W. Grycak, *On some class of warped product manifolds*, Bull. Inst. Math. Acad. Sinica 15 (1987), 311–322.
- [12] —, —, *On manifolds satisfying some curvature conditions*, Colloq. Math. 57 (1989), 89–92.
- [13] —, —, *On certain curvature conditions on Riemannian manifolds*, ibid. 58 (1990), 259–268.
- [14] R. Deszcz and M. Hotłoś, *Remarks on Riemannian manifolds satisfying certain curvature condition imposed on the Ricci tensor*, Prace Nauk. Politech. Szczec. 11 (1989), 23–34.
- [15] L. P. Eisenhart, *Riemannian Geometry*, Princeton Univ. Press, Princeton 1966.
- [16] W. Grycak, *Riemannian manifolds with a symmetry condition imposed on the 2-nd derivative of the conformal curvature tensor*, Tensor (N.S.) 46 (1987), 287–290.
- [17] G. I. Kruchkovich, *On semi-reducible Riemannian spaces*, Dokl. Akad. Nauk SSSR 115 (1957), 862–865 (in Russian).
- [18] K. Nomizu, *On the decomposition of generalized curvature tensor fields*, in: Differential Geometry in honor of K. Yano, Kinokuniya, Tokyo 1972, 335–345.
- [19] Y. Ogawa, *On conformally flat spaces with warped product Riemannian metric*, Nat. Sci. Rep. Ochanomizu Univ. 29 (1978), 117–127.
- [20] Z. I. Szabó, *Structure theorems on Riemannian spaces satisfying  $R(X, Y) \cdot R = 0$* , J. Differential Geom. 17 (1982), 531–582.

DEPARTMENT OF MATHEMATICS  
 AGRICULTURAL ACADEMY  
 C. NORWIDA 25  
 50-375 WROCLAW, POLAND

*Reçu par la Rédaction le 21.3.1989;  
 en version modifiée le 9.3.1990*