

## A NOTE ON CAREFUL PACKING OF A GRAPH

M. WOŹNIAK<sup>1</sup>

*Instytut Matematyki AGH  
al. Mickiewicza 30, 30-059 Kraków, Poland*

### Abstract

Let  $G$  be a simple graph of order  $n$  and size  $e(G)$ . It is well known that if  $e(G) \leq n - 2$ , then there is an edge-disjoint placement of two copies of  $G$  into  $K_n$ . We prove that with the same condition on size of  $G$  we have actually (with few exceptions) a careful packing of  $G$ , that is an edge-disjoint placement of two copies of  $G$  into  $K_n \setminus C_n$ .

**Keywords:** packing of graphs

**1991 Mathematics Subject Classification:** 05C70

### 1. INTRODUCTION

We shall use standard graph theory notation. We consider only finite, undirected graphs of order  $n = |V(G)|$  and size  $e(G) = |E(G)|$ . All graphs will be assumed to have neither loops nor multiple edges.

For graphs  $G$  and  $H$  we denote by  $G \cup H$  the *vertex disjoint union* of graphs  $G$  and  $H$  and  $kG$  stands for the disjoint union of  $k$  copies of the graph  $G$ .

Suppose  $G_1, \dots, G_k$  are graphs of order  $n$ . We say that there is a *packing* of  $G_1, \dots, G_k$  (into the complete graph  $K_n$ ) if there exist injections  $\alpha_i : V(G_i) \rightarrow V(K_n)$ ,  $i = 1, \dots, k$ , such that  $\alpha_i^*(E(G_i)) \cap \alpha_j^*(E(G_j)) = \emptyset$  for  $i \neq j$ , where the map  $\alpha_i^* : E(G_i) \rightarrow E(K_n)$  is induced by  $\alpha_i$ .

A packing of  $k$  copies of a graph  $G$  will be called a *k-placement* of  $G$ . A packing of two copies of  $G$  i.e. a 2-placement is an *embedding* of  $G$  (in its complement  $\overline{G}$ ). So, an embedding of a graph  $G$  is a permutation

---

<sup>1</sup>This paper was partially supported by Polish Research Grant KBN 2 P 301 050 031

$\sigma$  on  $V(G)$  such that if an edge  $xy$  belongs to  $E(G)$  then  $\sigma(x)\sigma(y)$  does not belong to  $E(G)$ .

A *careful packing* of a graph  $G$  is a packing of  $C_n$  and two copies of  $G$  into the complete graph. In others words this is an edge-disjoint placement of two copies of  $G$  into  $K_n \setminus C_n$ . Geometrically speaking, if we identify the cycle  $C_n$  with a convex  $n$ -gon on the plane, the careful packing of  $G$  means the possibility to draw (edge-disjointly) two copies of  $G$  using only the internal edges.

The following theorem was proved, independently, in [2], [4] and [7].

**Theorem 1.** *Let  $G = (V, E)$  be a graph of order  $n$ . If  $|E(G)| \leq n - 2$ , then  $G$  can be embedded in its complement  $\overline{G}$ .*

The example of the star  $K_{1,n-1}$  shows that Theorem 1 cannot be improved by increasing the size of  $G$ .

This result have been improved in many ways. For instance, the following theorem completely characterizes those graphs with  $n$  vertices and  $n - 1$  edges which are embeddable ([5], [6]).

**Theorem 2.** *Let  $G = (V, E)$  be a graph of order  $n$ . If  $|E(G)| \leq n - 1$ , then either  $G$  is embeddable or  $G$  is isomorphic to one of the following graphs :  $K_{1,n-1}$ ,  $K_{1,n-4} \cup K_3$  for  $n \geq 8$ ,  $K_1 \cup 2K_3$ ,  $K_1 \cup C_4$ ,  $K_1 \cup K_3$  and  $K_2 \cup K_3$ .*

**Remark .** For other generalization and improvements of Theorem 2 see for instance [8], [9] or [10]. The general references here are [11] and [1] (see also [12]).

Our purpose is to prove the following

**Theorem 3.** *Let  $G$  be a graph of order  $n$ ,  $n \geq 6$ . If  $e(G) \leq n - 2$ , then there exists a careful packing of  $G$  except for two graphs of order 6:  $K_3 \cup K_2 \cup K_1$  and  $C_4 \cup 2K_1$ , and for two families of graphs:  $K_{1,n-2} \cup K_1$  and  $K_{1,n-3} \cup K_2$ .*

The proof the theorem is given in the next section.

**Corollary 4.** *Let  $G$  be a graph of order  $n$ ,  $n \geq 3$ . If  $e(G) \leq n - 3$ , then there exists a careful packing of  $G$ .*

**Proof.** The corollary is evident for  $n = 3$  and  $4$  and easy to verify for  $n = 5$ . For  $n \geq 6$  it follows from Theorem 3. ■

We finish this section with some remarks.

Observe first that if we want to pack two copies of a graph  $G$  together with the cycle  $C_n$ , then the following necessary condition must hold:

$$\Delta(G) + \delta(G) \leq n - 3.$$

For, the vertex  $u$  with  $d(u) = \Delta(G)$  must be placed with another vertex of  $G$  and with a vertex of  $C_n$  of degree 2. Another evident, necessary condition is determined by the number of edges in the complete graph  $K_n$ . We must have  $2(n - 2) + n \leq \binom{n}{2}$  which implies  $n \geq 6$ .

So, from this point of view, there are only two “small” exceptional graphs in Theorem 3.

Since it is very easy to find a 2-placement for exceptional graphs of Theorem 3, so this theorem is an improvement of Theorem 1. On the other hand, Corollary 4 can also be considered as an improvement of the following theorem of Ore (cf.[3]).

**Theorem 5.** *If  $G$  is a simple graph of order  $n \geq 3$  and  $e(G) > \binom{n-1}{2} + 1$ , then  $G$  is Hamiltonian.*

Indeed, restated in terms of packing, Theorem 5 states that if  $G$  is a graph of order  $n$ ,  $n \geq 3$ , and  $e(G) \leq n - 3$ , then there is a packing of  $G$  into  $K_n \setminus C_n$ , whereas Corollary 4 ensures a packing of two copies of  $G$  into  $K_n \setminus C_n$ .

## 2. PROOF

We start with some simple observations formulated as lemmas.

**Lemma 6.** *Let  $G$  be a graph composed of the cycle  $C_k$  and one vertex, say  $u$ , not on the cycle. Denote by  $|N(u, C_k)|$  the number of edges connecting  $u$  with  $C_k$ . If  $|N(u, C_k)| > \frac{k}{2}$ , then the cycle  $C_k$  can be extended to a cycle of length  $k + 1$  passing through  $u$ . ■*

**Lemma 7.** *Let  $G$  be a graph composed of the cycle  $C_k$  and two vertices, say  $u, v$ , not on the cycle. If*

1.  $uv \in E(G)$ ,
2.  $|N(u, C_k)| \geq 1$ ,  $|N(v, C_k)| \geq 1$ ,
3.  $|N(u, C_k)| + |N(v, C_k)| \geq k + 1$ ,

then the cycle  $C_k$  can be extended to a cycle of length  $k+2$  passing through  $u$  and  $v$ .

**Proof.** It is easy to see that at least one of the neighbours of the vertex  $v$  on the cycle  $C_k$  has as its neighbour on the cycle  $C_k$ , a vertex connected by an edge with the vertex  $u$ . The possibility to extend the cycle  $C_k$  to the cycle  $C_{k+2}$  is now evident. ■

**Lemma 8.** *If the graph  $G$  has an end-vertex, say  $x$ , adjacent to the vertex, say  $y$ , of degree  $d(y) \geq \frac{n-1}{2}$  and there is a careful packing of  $G' = G \setminus \{x\}$ , then there is a careful packing of the graph  $G$ .*

**Proof.** Observe first that in the careful packing of  $G'$  the image of  $y$  is distinct from  $y$ . Indeed, otherwise we would have too many edges adjacent to  $y$  in  $K_{n-1}$  (two edges of  $C_{n-1}$  and at least  $n-2$  edges belonging to two copies of  $G'$ ).

Thus it is easy to extend the packing of  $G'$  (by putting  $x$  on  $x$ ) and then to extend  $C_{n-1}$  by applying Lemma 6 to the complement of the graph  $G$ . ■

**Proof of Theorem 3.** In the remainder of this section we adopt the following convention: Given a careful packing of a graph  $G$ , we say that an edge  $e$  of  $K_n$  is *black* or *blue* if it belongs to the first or second copy of  $G$ , respectively, and that an edge  $e$  of  $K_n$  is *red* if it belongs to the corresponding cycle  $C_n$ .

The proof is by induction on  $n$ . Without loss of generality we may assume that all the graphs under consideration are of maximum size  $n-2$ . Let us start with small values of  $n$  i.e.  $n=6$  and  $n=7$ . It is easy to see that there are five graphs of order 6 and size 4 which are not exceptional:  $K_1 \cup P_5$ ,  $K_1 \cup S'_5$ ,  $K_2 \cup P_4$ ,  $2P_3$  and  $2K_1 \cup (S_3 + e)$ . The careful packings of these graphs are depicted in Figure 1 (the edges of  $C_6$  are not marked). Observe that they can be used to obtain the careful packings of  $(n, n-2)$ -graphs for  $n=7$ . We can also use Lemma 8. The details are left to the reader.




Figure 1. Carefull packing of graphs of order 6

Suppose now that the theorem is true for all  $n' < n$  and let  $G$  be an  $(n, n-2)$ -graph. Assume also that  $G$  is not one of the exceptional graphs. We shall consider two main cases.

*Case 1.*  $G$  has two independent end-edges.

Denote the independent end-edges of  $G$  by  $uu'$  and  $vv'$ ,  $u, v$  being the corresponding end-vertices of  $G$ . Consider now the graph  $G' = G \setminus \{u, v\}$ . Suppose that there exists a careful packing for  $G'$ , say  $\sigma'$ . It is easy to extend the bijection  $\sigma'$  to a packing of  $G$ . Moreover, since the edge  $uv$  is neither black nor blue, we can consider it as a red one. We assign the red colour also to  $n-4$  edges connecting  $u$  with  $C_{n-2}$  and to  $n-4$  edges connecting  $v$  with  $C_{n-2}$ . By Lemma 7 (with  $k = n-2$ ) the careful packing of  $G$  exists. The case where  $G'$  is an exceptional graph will be considered below as *Case 3*.

*Case 2.*  $G$  has not two independent end-edges.

Since  $G$  has at least two tree components, the above condition implies that at least one of them is trivial and the other is a star. Let  $u$  be an isolated vertex of  $G$  and let  $x$  be a vertex defined by

$$d_G(x) = \min\{d_G(y) : y \in V(G), d_G(y) \geq 2\}$$

We consider the graph  $G' = G \setminus \{u, x\}$ . Suppose that  $G'$  is not one of the exceptional graphs; other cases are considered below as *Case 3*. Then there exists a careful packing for  $G'$ , say  $\sigma'$ . It is evident that by putting  $x$  on  $u$  and  $u$  on  $x$  we extend  $\sigma'$  to a packing of  $G$ . We may assume that the vertices  $x$  and  $u$  send  $n - 2 - d(x)$  red edges to the red cycle  $C_{n-2}$  contained in  $G'$ . We can apply Lemma 7 and obtain a careful packing of  $G$  if  $2(n - 2 - d_G(x)) \geq n - 1$ . Hence  $n - 3 \geq 2d_G(x)$ .

Thus, we may assume that

$$(*) \quad d_G(x) \geq \frac{n-2}{2}$$

So, for  $n \geq 7$ ,  $d_G(x) \geq 3$ . Consider first the case where  $G$  has two trivial components.

*Case 2 (a)*  $G$  has two isolated vertices, say  $u, v$ .

Consider first the case  $n = 8$ . The case by case examination shows that: either  $G$  contains an end-vertex such that we can apply Lemma 8, or  $G$  is such that the graph  $G' = G \setminus \{u, x\}$  is exceptional (see *Case 3*). So, we may assume that  $n \geq 9$ . Consider now the graph  $G_1 = G \setminus \{u, v, x\}$ . If  $G_1$  is not one of the exceptional graphs, we can apply the induction hypothesis. Let  $\sigma_1$  be a careful packing of  $G_1$ . Denote by  $y_1$  a vertex of  $G_1$  non adjacent to  $x$  (such a vertex exists by the definition of  $x$ ). Without loss of generality we may assume that  $y_1$  is the first vertex on the red cycle  $y_1, y_2, \dots, y_{n-3}$  corresponding to the careful packing of  $G_1$ . Then the cycle  $xy_1y_2 \dots y_{n-3}uvx$  can be considered as a red cycle of the careful packing of  $G$ , say  $\sigma$ , obtained from  $\sigma'$  by putting  $\sigma(x) = v$ ,  $\sigma(v) = x$ ,  $\sigma(u) = u$  and  $\sigma(w) = \sigma'(w)$  for  $w \in V(G) \setminus \{u, v, x\}$ .

*Case 2 (b)*  $G$  has only one isolated vertex.

Hence  $G$  is of the form  $K_1 \cup K_{1,r} \cup R$  where  $r \geq 1$  and the graph  $R$  has no isolated vertices. Moreover, since by *Case 1*,  $R$  contains no end-vertices we may assume, by (\*), that either all vertices of  $R$  are of a degree greater than or equal to  $\frac{n-2}{2}$ , or  $R$  is empty. In the first case, for  $n > 6$ , this contradicts the fact that the average degree of  $R$  is equal to 2. In the second case  $G$  is exceptional, a contradiction.

*Case 3.*  $G'$  is one of the exceptional graphs, where  $G'$  denotes one of the graphs defined in *Cases 1* or *2* ( $n \geq 8$ ).

We shall need some additional notations. Namely, by  $S'_p$  we denote a tree of order  $p$  obtained by subdividing one of the edges of the star  $K_{1,p-2}$  and by  $(K_{1,p-1} + e)$  we denote, as usually, the graph of order  $p$  obtained by adding one edge to the edge-set of the star  $K_{1,p-1}$ .




Figure 2. Carefull packing of  $K_2 \cup P_3 \cup C_3$  and  $K_1 \cup K_{1,3} \cup C_3$

Without loss of generality we may assume that every other choice of two or three (for  $n \geq 9$ ) vertices in a way described in *Cases 1* and *2* leads also to one of the exceptional graphs. Of course, we can proceed as in *Case 2* also in the case where the graph  $G$  has two independent end-edges.

Recall that  $G$  itself is not an exceptional graph.

The case by case examination shows that then  $G$  belongs to one of the following families of graphs:  $P_3 \cup K_{1,n-4}$ ,  $K_1 \cup S'_{n-1}$ ,  $2K_1 \cup (K_{1,n-3} + e)$ ,  $K_1 \cup K_3 \cup K_{1,n-5}$ , or  $n = 8$  and  $G$  is isomorphic to  $4K_1 \cup K_4$ ,  $2K_1 \cup 2K_3$ ,  $2K_2 \cup C_4$ ,  $K_2 \cup P_3 \cup C_3$  or  $3K_1 \cup K_{2,3}$ .

Observe that in all graphs belonging to the above mentioned families, except for  $K_1 \cup K_3 \cup K_{1,3}$ , there is a vertex of a degree greater than or equal to  $n - 4$ , so we can apply Lemma 8 (since  $n \geq 8$ ).

The careful packings of  $4K_1 \cup K_4$ ,  $2K_2 \cup C_4$  or  $2K_1 \cup 2K_3$  are very symmetric and easy to find.

The careful packing of  $K_2 \cup P_3 \cup C_3$  as well as the careful packing of  $K_1 \cup K_3 \cup K_{1,3}$  are depicted in Fig. 2.

Finally, the careful packing of  $3K_1 \cup K_{2,3}$  can be easily obtained from the careful packing of  $2K_1 \cup K_{1,3}$  into  $K_6$ .

This completes the proof of the theorem. ■

#### REFERENCES

- [1] B. Bollobás, *Extremal Graph Theory* (Academic Press, London, 1978).
- [2] B. Bollobás and S.E. Eldridge, *Packings of graphs and applications to computational complexity*, J. Combin Theory (B) **25** (1978) 105–124.
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, (North-Holland, New York, 1976).

- [4] D. Burns and S. Schuster, *Every  $(p, p-2)$ -graph is contained in its complement*, J. Graph Theory **1** (1977) 277–279.
- [5] D. Burns and S. Schuster, *Embedding  $(n, n-1)$ -graphs in their complements*, Israel J. Math. **30** (1978) 313–320.
- [6] B. Ganter, J. Pelikan and L. Teirlinck, *Small sprawling systems of equicardinal sets*, Ars Combinatoria **4** (1977) 133–142.
- [7] N. Sauer and J. Spencer, *Edge disjoint placement of graphs*, J. Combin Theory (B) **25** (1978) 295–302.
- [8] M. Woźniak, *Embedding of graphs in the complements of their squares*, in: J. Nesselrill and M. Fiedler, eds, Fourth Czechoslovakian Symp. on Combinatorics, Graphs and Complexity, (Elsevier Science Publishers B.V., 1992) 345–349.
- [9] M. Woźniak, *Embedding graphs of small size*, Discrete Applied Math. **51** (1994) 233–241.
- [10] M. Woźniak and A.P. Wojda, *Triple placement of graphs*, Graphs and Combinatorics **9** (1993) 85–91.
- [11] H.P. Yap, *Some Topics In Graph Theory* (London Mathematical Society, Lectures Notes Series 108, Cambridge University Press, Cambridge 1986).
- [12] H.P. Yap, *Packing of graphs – a survey*, Discrete Math. **72** (1988) 395–404.

Received 26 April 1994

Revised 14 July 1994